

# Use of Duckweed Growing on Sewage Water as Poultry Feed

Shazia Iram\*, Sumera Abrar\*, Iftikhar Ahmad\*\*, Tassawar Khanam\*\*, Atiya Azim\*\*, Mukhtar Ahmad Nadeem\*\*

\*Department of Environmental Sciences, Fatima Jinnah Women University, Rawalpindi, Pakistan.

\*\*Institute of Animal Sciences at National Agricultural Research Centre (NARC), Islamabad, Pakistan.

**Abstract-** Present study was focused on intake of duckweed by chicks and to evaluate the growth performance and carcass characteristics of chicks. A total number of 106 chicks were allocated to 3 treatments with 3 replicates. The treatments were categorized into group A (25% duckweed and 75% commercial feed), group B (50% duckweed and 50% commercial feed) and group C (100% commercial feed). Results showed that crude protein was higher in group B followed by group A while in group C crude protein was lower but total feed intake, average weight, live weight, meat and bone weight, skin weight, head weight, liver weight, intestine weight and shank weight were higher as compared to group A and group B and this was due to the fluffy nature of duckweed and small size of crop. It was concluded that sewage grown duckweed can be successfully utilized as poultry feed. Chicks can survive on duckweed and farmer can save money. This is the first study in Pakistan.

**Index Terms-** Bioremediation; duckweed; poultry feed; sewage water.

## I. INTRODUCTION

Poultry farming has now become one of the most dynamic associated parts of agriculture throughout the world. Poultry culture in South Asian countries is expanding rapidly and the rate of growth of commercial layer and broiler (meat producing) farms is phenomenal to meet the ever increasing demand for proteins through poultry meat and eggs. In Pakistan, poultry industry had made considerable contribution to food production and plays a vital role in the economy of the country. Recently duckweed system was introduced to solve the feed problems of farmer. Duckweed plays an important role in the extraction and accumulation of metals from water (Susarla *et al.*, 2002). Duckweed accumulates excess nutrients into their own biomass (Boyt *et al.*, 1977) and later be processed into a variety of products such as poultry feed and fishmeal thereby bringing a cost recovery aspect into the waste management process (Hillman and Culley, 1978). The duckweeds (*Lemna* sp., *Spirodella* sp.) have received the most attention because of their rapid growth to high biomass, ease of handling and high protein content as compared to the other macrophytes.


Numerous studies have demonstrated the value of duckweed as a feed for poultry, fish and other animals (Skillicorn *et al.*,

1993). Duckweed closely resembles to the animal protein and has a better array of essential amino acids than the vegetable protein (Hillman and Culley, 1978). The amino acid profile of duckweed and soybean is very similar (Rusoff *et al.*, 1980) and also has a rich concentration of beta carotene and xanthophylls (Journey *et al.*, 1993). Dried duckweed contains crude protein up to 40% (Vandyke *et al.*, 1977) and can be compared with soybean meal as a source of plant protein. Duckweed was used as feed for poultry because of its high nutritional value (Abdullaev, 1969). Duckweed when dehydrated has been used to replace alfalfa (Lucerne) meal as a protein source in conventional poultry diets as it has high protein content 20-40%, low fiber content, high mineral content, non toxicity and few known pest. The protein content of duckweed responds quickly to the availability of nutrients in the water (Leng *et al.*, 1995). Duckweed has been known for a long period of time as a potential source of food for humans and animals and as a source of natural products it is the only source of supplementary protein for fish, chickens ducks and pigs (Rodriguez *et al.*, 1996). Village pigs, horses or ruminants could be fed on freshly harvested duckweed. For most applications with poultry, dried duckweed would be preferable as it could be expected that duckweed would provide an ideal wet supplement to any high energy diet (Leng *et al.*, 1995). Aim of the current study was to utilize the sewage grown duckweed as feed for poultry for saving the farmers' money.

## II. MATERIALS AND METHODS

### *Collection of Duckweed*

The duckweed (*Lemna minor*) was harvested from the wastewater treatment ponds of National Institute of Bioremediation at National Agricultural Research Center (NARC), Islamabad, Pakistan. Purpose was to reclaim the sewage water of main NARC offices building and hostels through bioremediation for irrigation purposes (Fig. 1). The used water treatment garden project was started in October 2008 and finalized in February 2009. Total area was about 0.38 acre. The used water treatment garden facilitated to treat 35,000 US gallons per day for irrigation. The toxicity analysis report of the output water from treatment garden indicated that water was fit for irrigation.



**Used-Water treatment management at National Agriculture research Center**

**Fig. 1 Bioremediation ponds for treatment of sewage water.**

After collection of duckweed from pond 4 and 5, the samples were brought in laboratory for analysis and rinsed in distilled water to remove dust and stored in refrigerator for further procedure (freezing and drying). There were two types of duckweed, small leaves duckweed and large leaves duckweed. Duckweed (*Lemna minor*) was dried in the air for 30 days without sun-drying. Small and large leaves were mixed together and then milled to fine particles after mixed with the commercial feed (standard diet) (table 1) according to the required percentages (Khanum *et al.*, 2005).

Soyabean meal	10.30	Phyzyme	0.00
Oil	1.260	Allzyme	0.00
Premix	0.50		

In composition of feed, crude protein, dry matter, ether extract, fiber, ash, gross energy, nitrogen free extract (NFE), K, P, Na Ca and aflatoxin were examined and in addition small leaves, large leaves, mixture of small and large leaves and composition of duckweed and commercial feed after mixing were also analyzed (AOAC, 1994).

**Chicks rearing and dietary treatment**

A total number of 106 chicks were procured from commercial hatchery. Experiment was designed with 3 treatments using 3 replicates each having 11 to13 chicks. These chicks were raised indoors on deep litter system. The chicks were kept up to 4 weeks of age. The experiment was conducted during June, 2010.

In housing, the chicks were provided with 1 square feet space per bird. Lighting schedule was practiced throughout the experiment as well as standard brooding and rearing temperature was maintained.

**Table 1. Ingredients of commercial feed of chicks.**

Ingredients	Percentage	Ingredients	Percentage
Corn	44.00	Sunflower meal	6.00
Rice	10.00	Marbel chips	0.70
Rice polish	7.00	Dicalcium phos	1.57
Wheat bran	3.90	L-lysine	0.20
C. gluten meal 60%	0.70	Lysine sulphate	0.27
Rapeseed meal	3.00	DL-methionine	0.13
Canola meal	10.30	Sod.Chlorine	0.35
Guar meal	1.00	Threonine	0.08



**Fig. 2 Housing system of chicks.**

Water for birds was obtained from a clean source and boiled to eradicate the contaminants. Water was accessible to the birds all the time in the pens as well as vaccination was also provided to the chicks (Table 2).

**Table 2. Chick's vaccination.**

Days	Vaccination	Nature
Day-01	IB	Eye-Drop
Day-06	ND	Eye-Drop
Day-10	IBD	Eye-Drop
Day-15	HP	Injection
Day-20	IBD	Eye-Drop
Day-24	ND + IB	Eye-Drop

**Statistical analyses**

Data regarding weight of bird, weight gain per bird, weight gain per bird per day, feed intake, intake per bird per day and feed conversion ratio was recorded analyzed with Complete Randomized Design (CRD) with two factor factorial analysis of variance by using Minitab software. For significant P-value, least significant difference (LSD) was determined for mean comparison at 5% level.

**III. RESULTS AND DISCUSSION**

Duckweed closely resembles to the animal protein and has a better array of essential amino acids than the vegetable protein (Hillman *et al.*, 1978). The amino acid profile of duckweed and

soybean is very similar (Rusoff *et al.*, 1980) and also has a rich concentration of beta carotene and xanthophylls (Journey *et al.*, 1993). Dried duckweed contains crude protein up to 40% (Vandyke *et al.*, 1977) and can be compared with soybean meal as a source of plant protein. Duckweed was used as feed for poultry because of its high nutritional value (Musaffarov *et al.*, 1968). It has great potential as feed for poultry, and has high moisture content (Leng, 1995). In present study large leaves duckweed has high dry matter (DM), crude protein (CP), ether extract (EE), crude fiber (CF), N and aflatoxin compared to small leaves and mixed leaves duckweed. The ash content was higher in small leaves whereas nitrogen free extract was higher in mixed leaves of duckweed (Table 3).

**Table 3. Chemical composition of duckweed leaves.**


Composition	Small leaves	Large leaves	Mixture of small & large leaves
DM	4.62	6.53	3.77
CP	25.30	29.67	24.01
EE	6.91	7.30	1.37
CF	10.73	11.14	10.21
Ca	3.99	4	4.75
P	1.17	0.94	1.16
N	4.04	4.74	3.84
Na	0.67	0.426	1.8
K	1.205	1.225	4.75
Ash	28.03	19.00	23.69
Nitrogen free extract	32.89	32.89	40.72
Aflatoxin (ppb)	55	76	-
Gross energy (MJ/kg)	12.31	12.89	12.7

Duckweed was mixed with the commercial feed categorized into group A, B and C according to composition (Table 3). The dry matter contents of all groups were almost


similar. Crude protein was higher (22.67 %) in group B while in group A it was 21.0% but in group C it was very lower (20.65%).

**Table 4. Composition of duckweed and commercial feed after mixing.**


Composition	Group A	Group B	Group C
DM	92.41	92.05	92.46
CP	21.0	22.67	20.65
EE	4.54	3.65	5.97
CF	5.57	5.89	4.17
Ca	0.9	1.6	0.4
P	0.82	0.73	0.72
Na	0.12	0.17	0.07
K	0.53	1.23	4.75
Ash	13.18	19.55	7.22
NFE	55.71	48.24	61.99
Gross energy (MJ/kg)	10.2	12.5	16.175



**Fig. 3a Average weight per bird in different treatments.**



**Fig. 3b Average weight gain per bird in different treatments.**



**Fig. 3c Average weight gain per bird per day in different treatments.**

The gross energy was higher in group C (16.175 MJ/kg) and in group A and B it was lower (10.21, 2.5MJ/kg). Crude fiber (5.89%), calcium (1.60%), sodium (17%) and ash (19.55%) were higher in group B whereas nitrogen free extract (61.99%) and ether extract (5.97%) was higher in group C. Gross energy was higher (16.175 MJ/kg) in group C (commercial feed) and low in group B (12.5 MJ/kg) and A (10.2 MJ/kg).

In the beginning of study, the weight of chicks in all treatments was same. During third and 4<sup>th</sup> week, weight of chicks was higher in group C followed by group A and B (Fig 3a). In case weight gain per bird there was no significant difference in weight during 1<sup>st</sup> and 2<sup>nd</sup> week but during week 3<sup>rd</sup> and 4<sup>th</sup> week, weight was higher in group C (Fig 3c). In case of average weight gain, the weight was higher in group C (Table 5). Similar studies was conducted by Ngamsaeng *et al.* (2004) on 24 Muscovy ducks in order to evaluate the protein quality in water spinach (WS) or duckweed (DW) alone or mixed (WS+DW) and they reported average daily gain was highest for ducks fed diet DW (22.4 g/bird/day) and lowest when fed diet WS (6.2 g/bird/day).

Khang, (2003) found Chicks fed DW had somewhat higher weight gains (8.3 g/day) compared with chicks fed the diets without DW (7.8 g/day) but no differences were seen in between diets with different CP levels. Khanum *et al.* (2005) reported the final live weight and average daily body weight gain were significantly lower in the control diet but were not different between duckweed diets. Samnang (1999) found in the station, soyabean supplemented chickens grew faster than duckweed supplemented.

The growth of very young broiler chickens may be retarded with increasing levels of *Lemna gibba* dehydrated meal in the diet whereas layer hens produced effectively (Haustein *et al.*, 1990a). The growth of chickens fed duckweed, group A and group B was less than the group C which consist of totally commercial feed (P=0.001). It was due to the low intake of chickens fed duckweed. Duckweed was eaten but the amount was less because the birds digestive tract adapts to the type and quantity of food available (Klasing, 1998).




Fig. 4a Feed intake in different treatments.




Fig. 4b Feed intake per bird in different treatments.




Fig. 4c Feed intake per bird per day in different treatments.

During present investigation it was observed that the crop size of chicken fed duckweed was enlarged. This was due to the fluffy nature of duckweed as it absorbs more moisture as a result of which the crop size become enlarge. Duckweed was eaten but the amount was less than reported by Nguyen and Brian, (2004). The birds digestive tract adapts to the type and quantity of food available (Klasing, 1998). The greater size of crop in birds fed duckweed may be because of higher intake of fiber. As a result of which the group A and group B the chickens has low feed intake and due to this low intake the weight also reduce when compared to the group C which was offered no duckweed. Klasing, (1998) also reported that crop capacity increased on diets high in grass or leaves compared with a diet based on ground grains. The chickens with access to duckweed tended to have a lower carcass yield than control chickens, probably

because of higher contents in the digestive tract as indicated by large size of crop. Mortality rate was very low through out the experiment only two mortality was seen in the group C d and no mortality was observed in group A and group B. When postmortem of chicks were performed it was observed that these mortalities were not due to any disease but due to a litter which block the respiratory passage and becomes the cause of death (Figure 4abc, Table 5).

Feed Conversion Ratio (FCR) is a measure of a bird's efficiency in converting feed mass into increased body mass. The smaller the FCR the more efficient birds are at converting feed to meat. FCR of group B was maximum and FCR of group C was minimum. So group C with smaller FCR, the birds were more efficient in converting feed to meat (Fig. 5, Table 5).




Fig. 5 Feed conversion ratio in different treatments and weeks.

Table 5. Average weight/bird, weight gain per bird, weight gain per bird per day, feed intake, intake per bird per day and feed conversion ratio.

	wt/bird	wt gain/bird	wt gain/bird/day	Feed intake	Intake/bird	Intake/bird/day	FCR
A	396	163	23	3637	295	42	1.72
B	356	139	19	3729	311	44	2.2
C	473	216	30	3861	366	52	1.61

At the end of the trail one chicken from each of the replicate was slaughter and record the weight of different organs. The chickens with access to duckweed (group A and group B) tended to had lower carcass yield than control chickens (group

C), probably because of higher contents in the digestive tract as indicated by large size of crop. As shown in the table 6 the live weight of group C was very high (1013g) as compared to group A (811g) and group B (777g). Meat and bone weight was also

high in group C (473g) compared to group A (383g) and group B (318g). Skin weight of group C was increased (141g) as compared to group A (97g) and group B (86g). The weight of other organs such as head, liver, intestine, gizzard with feed,

empty gizzard, shank was high in the chicks belong to group C. After group C the weight of organ was high in the chicks belong to group A and weight of organs was less in the chicken belong to group B.

**Table 5. Mean values for carcass of chicks given duckweed as replacement for commercial feed.**

Organs	Organ weight (g)		
	Group A	Group B	Group C
Live weight	811	777	1013
Meat and Bone	383	318	473
Skin	97	86	141
Head	26	24	32
Liver	23	25	35
Heart	4	6	6
Intestine	74	79	88
Gizzard with feed	32	29	32
Empty gizzard	18	14	17
Shank	33	31	42

Kabir *et al.* (2005) found that body weight, feed intake, feed efficiency, protein efficiency, energy efficiency and profitability linearly declined as the proportion of duckweed increased in the diet. Duckweed did not affect on livability ( $P>0.05$ ). The growth of very young broiler chickens may be retarded with increasing levels of *Lemna gibba* dehydrated meal in the diet whereas layer hens produced effectively (Haustein *et al.*, 1990a). Khanum *et al.* (2005) found that the final live weight and average daily body weight gain was significantly lower in the control diet but were not different between duckweed diets. Samnang, (1999) found that Soyabean supplemented chickens grew faster than duckweed supplemented. Khang (2003) conducted experiment in order to evaluate the effects of duckweed on the laying performance of local (Tau Vang) hens and concluded that egg production, egg quality, feed conversion and net profit are highest when fresh duckweed replaces 75% of the protein from roasted soya beans in a diet based on broken rice. Even at 100% of roasted soya beans replacement by duckweed, the egg production and margin of income over feed costs were better than on the control diet in which the supplementary protein came only from roasted soya beans.

#### IV. CONCLUSION

Sewage grown duckweed can be successfully utilized as poultry feed. The intake and growth rate of chicks offered duckweed was less due to the fluffy nature of duckweed but they survived. Preparation of duckweed does not require extra labor cost for growing and harvesting. It can be produced at house hold level and save farmers money but at commercial level where weight gain is preferred, duckweed does not perform well as chicks do not gain much weight. So the duckweed plant from the wastewater material can be converted into useful product.

#### ACKNOWLEDGEMENT

The authors would like to thanks the Institute of Animal Sciences, National Agricultural Research Center, Islamabad for allowing the use of their facilities.

#### REFERENCES

- [1] Abdullaev, D. A. 1969. The use of common duckweed as green feed for chickens (In Russian). *J. Uzbekskii. Bio. Cheskii. (USSR)*.13: 42-43.
- [2] AOAC. 1994. Official methods of analysis of association of official analytical chemists. Washington. D.C.
- [3] Boyt, F. L., S. E., Bayley and J. Zoltet. 1977. Removal of nutrients from treated municipal wastewaters by wetland vegetation. *J. Water. Poll. Con. Feder.*, (49).
- [4] Haustein, A.T., R. H. Gilman, P. W. Skillicorn, V. Vergara, V. Guevara and A. Gastanaduy. 1990a. Duckweed. A useful strategy for feeding chickens: performance of layers fed with sewage-grown Lemnaceae species. *J.Poul. Sci.* 69: 1835-1844.
- [5] Hillman, W. S. and D. D. Culley. 1978. The uses of duckweed. *Ani. Sci.* 66: 442-451.
- [6] Journey, W. K., P. Skillicorn and W. Spira. 1993. Duckweed Aquaculture A new aquatic farming system for developing countries. World Bank, Washington.D.C.
- [7] Kabir, J., M. A. Islam, M. U. Ahamad and M. A. Howlider. 2005. Use of duckweed (*Lemna minor*) in the diet of broiler. *J. Indian.Animal.Research.* 39(1). Print ISSN : 0367-6722.
- [8] Khang, N. T. K. 2003. Use of Duckweed (*Lemna Minor*) as a protein supplement for local (Tau Vang) chicks, and growing and laying hens, MSc Thesis. Swedish University of Agricultural Sciences, Uppsala, Sweden.
- [9] Khanum, J., A. Chwalibog and K. S. Huque. 2005. Study on digestibility and feeding systems of duckweed in growing ducks. *J. Livestock Research for Rural Development.* 17 (5).
- [10] Klasing, K. C. 1998. Comparative Avian Nutrition. CAB International. British Library, London, UK.
- [11] Leng, R.A., J. H. Stambolie and R. Bell. 1995. Duckweed - a potential high-protein feed resource for domestic animals and fish. *Live. Stock. Res. Rural. Dev.*7(1).
- [12] Muzaffarov, A. M., T. Taubayev, M. Abdiyev. 1968. The use of Lemna minor L. for poultry feeding (In Russian). *J.Usbekskii Biologia Cheskii Journal (USSR)*. 12: 44-46.

- [13] Ngamsaeng, A., S. Thy, T. R. Preston. 2004. Duckweed (Lemna minor) and Water spinach (Ipomoea aquatica) as protein supplements for ducks fed broken rice as basal diet. *Live stock Res. Rural. Dev.* 16: 18-24.
- [14] Nguyen, T. and O. Brian. 2004. The effect of supplementing different green feeds (water spinach, sweet potato leaves and duckweed) to broken rice based diets on performance, meat and egg yolk colour of Luong Phuong chickens. Retrieved February 23, 112, from MEKARN Research Reports.
- [15] Rodriguez, L. and T. R. Preston. 1996. Comparative parameters of digestion and N metabolism in Mong Cai and Mong Cai Large White cross piglets having free access to sugar cane juice and duckweed. *Live stock Res. Rural. Dev.* 8: 72-81.
- [16] Rusoff, L. L., W. W. Blakenc and D. D. Culley. 1980. Duckweeds (Lemnaceae family): A potential source of protein and amino acids. *J. Agri. Food. Chem.* 28: 848-850.
- [17] Samnang, H. 1999. Duckweed versus ground soya bean as supplements for scavenging native chickens in an integrated farming system. *Live stock Res. Rural. Dev.* 11, 1. <http://www.cipav.org.co/lrrd/lrr11/1/sam111.htm>
- [18] Susarla, S., V. F. Medina and S. C. McCutcheon. 2002. Phytoremediation: an ecological solution to organic chemical contamination. *J. Ecol. Eng.* 18: 647-658.
- [19] Skillicorn, P., W. Spira and W. K. Journey. 1993. Duckweed Aquaculture - A new aquatic farming system for developing countries. Washington. D. C.
- [20] Vandyke, J. M. and D. L. Sutton. 1977. A feeding trial for grass carp using fresh duckweed (Lemna sp.) as food. *J. Fish. Bio.* 2: 273-278.

#### AUTHORS

**First Author** – Shazia Iram, Department of Environmental Sciences, Fatima Jinnah Women University, Rawalpindi, Pakistan.

**Second Author** – Sumera Abrar, Department of Environmental Sciences, Fatima Jinnah Women University, Rawalpindi, Pakistan., Email: [sumeraabrar75@gmail.com](mailto:sumeraabrar75@gmail.com)

**Third Author** – Iftikhar Ahmad, Institute of Animal Sciences at National Agricultural Research Centre (NARC), Islamabad, Pakistan.

**Fourth Author** – Tassawar Khanam, Institute of Animal Sciences at National Agricultural Research Centre (NARC), Islamabad, Pakistan.

**Fifth Author** – Atiya Azim, Institute of Animal Sciences at National Agricultural Research Centre (NARC), Islamabad, Pakistan.

**Sixth Author** – Mukhtar Ahmad Nadeem, Institute of Animal Sciences at National Agricultural Research Centre (NARC), Islamabad, Pakistan.