

Allelopathic Potential of *Papaver Somniferum* L. on Seed Germination Against Three Different Varieties of *Zea Mays*

Sidra-Tul-Muntaha, Sayeda Sarah Mubarak, Fatima Khan Sherwani, Nazma Azim

Department of Botany, Govt. Frontier College for Women Peshawar, Pakistan

DOI: 10.29322/IJSRP.11.04.2021.p11241

<http://dx.doi.org/10.29322/IJSRP.11.04.2021.p11241>

Abstract- To evaluate the allelopathic effect of *Papaver somniferum* against the 3 varieties of *Zea mays* seeds cv. Azam, Pahari and Iqbal. Aqueous extracts were made by soaking the powder of dried leaves of *Papaver somniferum*. For the aqueous extract treatment, the extract was soaked in distilled water for 24 and 48 hours separately. For mulching treatment, the seeds were sowed in soil and sprinkled powder of leaves. For litter, the petri dishes were grounded with randomly cut pieces of filter paper wetted with extract. For hot water treatment the water was boiled for 5 minutes. The powder was then mixed in boil water. Azam got effected in 24h extract in aqueous extract treatment while Pahari in 48h treatment. In mulching all the three varieties got inhibited by extract. Azam showed resistance toward the inhibitory effect of extract in litter experiments. Iqbal was affected in hot water extract. It is concluded that the extracts obtained from *Papaver somniferum* showed inhibitory effects on all the three varieties except few exceptions in treatments. The plumule and radical growth of seeds got affected the most among other aspects.

Index Terms- allelopathy; maize; mulching; litter; hot water extract.

I. INTRODUCTION

Allelopathy is the favorable or unfavorable effects of a plant on another plant due to the release of biochemicals, which are known as allelochemicals. This could be from plant parts by extracts, root, evaporation, disintegration of litter, and other processes in both natural and agricultural ecosystems (7). Allelochemicals are released from plants and crops, may have reverse effects in agricultural fields and controlled forest ecosystems that cause reduction in crop yield. These metabolites get stored and preserved for some time and put an influence on the growth and development of surrounding plants (19). Weeds are unwanted plants, which penetrate different crops and impose negative effect on their production. There are uncountable reports on the destructive effects of weeds on crops (11 and 3). Some type of allelochemicals, micro-organisms and soil conditions each play a significant role in determining the preservation of allelochemicals in soils. After extraction and verification of allelochemicals, it becomes necessary to examine their behavior in soil (13). In the past, many methods were applied for assessment of allelopathic potential. For example, aqueous extract method

applied on medicinal plants exposed high allelopathic potential to some extent (9 and 8). To our knowledge, little is known about the allelopathic potential of *Papaver somniferum* tissues or extracts. Therefore, the current study aims to explore the allelopathic potential of *P. somniferum* with an emphasis on the inhibition of seed germination of maize varieties i.e., Azam, Pahari and Iqbal by different methods. We tested the following hypothesis: the root and shoot fresh aqueous extracts of *P. somniferum* would reduce seed germination of each of the 3 target varieties, and that this reduction will increase with increasing soaking time.

II. MATERIAL AND METHOD

Leaves of *P. somniferum* were collected from Govt. Frontier College Peshawar and were air dried at room temperature. Powder was made from the dried leaves. Two aqueous extracts were made by soaking powder for 24 hours and 48 hours, respectively. Two Petri dishes were set for 3 varieties of maize seeds i.e., Azam, Pahari and Iqbal. Each Petri dish had 5 seeds randomly arranged.

Procedure of aqueous extracts:

1.25 gm of dried leaves powder was separately soaked in 100ml distilled water at 25°C for 24 and 48 hours and filtered to get aqueous extract. These extracts were tested against Azam, Pahari and Iqbal maize varieties. 3 filter papers were layered in Petri dishes. The filter papers were moistened with the extract. The control series were moistened with distilled water. After 72 hours germination, growth of plumule and radical were observed. Five Seedlings were taken out randomly for fresh and dry weight determination. Seedlings were dried for 72 hours.

Procedure of litter:

5 grams of randomly cut filter papers were set on the base Petri dishes. Test series were wet with aqueous extract and control series with distilled water. 5 seeds of each variety were set on litter. After 72 hours of germination, the length of plumule and radical were measured.

Procedure of mulching:

Five g of crushed dried leaf powder was layered on surface of plastic glasses containing sterilized moist sand. For each treatment, three replicates, each with 5 seeds were made. The control consisted of fine pieces of filter paper. After 72hours of

germination, growth of plumule and radical were measured. 5 seedlings were randomly taken out for fresh and dry weight and moisture contents.

Procedure of hot water extract:

1.25 gram of dried plant part boiled in water for 5 mints. The extract was cooled at room temperature. The seeds were arranged in Petri dishes and set for 72 hours to germinate. Each treatment of this experiment was carried out with 3 replicates with 5 seeds in it. The data obtained were subjected to statistical analysis of variance (ANOVA) by statistix 8.1 software.

III. RESULTS

The allelopathic activity of *Papaver somniferum* was determined on different varieties of *Zea mays* and observed its plumule and radical growth. In all activities the germination of plumule length and radical length of *Zea mays* was noted as compared to control.

Effect of aqueous extracts:

As shown in figure 1, the germination in Azam and Iqbal variety 100% growth while Pahari showed lesser growth than them. In test series of 24h test, Azam's germination rate decreased while Pahari increased, and Iqbal showed no effect. 48h series showed growth inhibition in Iqbal while increased in Pahari and showed no effect in Azam variety. Plumule growth of Azam is inhibited in 24h and 48h extracts as showed in table no 1. In Pahari, the growth was inhibited less by 24h extract than inhibition by 48h. Iqbal variety was inhibited by 24h extract while it was stimulated by 48h extract. Radical growth of Azam variety was inhibited by both the extracts but 24h showed more inhibitory effect than 48h. In Pahari, the growth increased by 24h while in 48h test it showed inhibition. In Iqbal, the radical was inhibited by both the extracts. Aqueous extracts from leaves soaked for 24 and 48 h significantly reduced plumule growth and radical length growth in all three varieties of maize: Azam, Pahari and Iqbal. The plumule growth in Pahari and radical growth of Iqbal was stimulated in 24h and reduced in 48h treatment. (Table no. 1)

Effect of Mulching:

This aspect when tested by using *P. somniferum* mulch in experiments it was seen that germination was inhibited in Azam and showed no effect in Pahari while in Iqbal the germination percentage increased as compared to control as shown in the figure 2. The plumule length was inhibited in all the three varieties but in Iqbal, the plumule length was reduced up to 50%. The radical length was very much reduced by the extract as compared with control series. (Table no. 2)

Effect of litter:

It is observed that before decay many species release phytotoxic substances. In current study it was observed that litter of *P. somniferum* when used as growth medium the germination percentage was not affected in Azam and Pahari while increased in Iqbal. Extracts significantly reduced the plumule growth and radical length growth in Pahari and Iqbal varieties of maize. However, the plumule growth and radical growth was stimulated in Azam. (Table no. 3)

Effect of hot water extract:

The percentage of germination was not affected by the extracts in Azam and Pahari while in Iqbal it showed stimulatory effect. The plumule length was reduced by the extract in all three varieties but Pahari was more affected than other two varieties. The radical emergence was reduced in Pahari and Iqbal while stimulated in Azam. (Table no. 4).

The growth of plumule and radical was reduced by the extracts in mulching experiment in all the three varieties. Pahari was affected the most by the phytotoxic effect of extract. Whereas Iqbal was affected in mulching and hot water extract.

DISCUSSIONS

The findings obtained from the aqueous extract treatment agree with (2 and 21) in which it was proven that enhancing soaking duration increased phytotoxicity of aqueous extracts. The extract inhibited the growth of both plumule and radical of the varieties.

(14) Confirmed that plants release chemicals also from dead tissues, and their absorption in the soil could be accelerated by leaching thus facilitating their harmful effects in the field. Allelopathic substances which are released by the plants accumulate in the soil to physiological activity level. This aspect was proven in mulching treatment where the radical of all the three varieties were reduced to high extents.

In litter treatment the extract caused not significant inhibition in Pahari and Iqbal. Azam resisted the repressing effect and increased in length compared to control. These results were agreed with the allelopathic effects of *Ageratina adenphora* plant litter on growth of *Lantana camara* (15 and 20).

(5 and 17) reported hot water extracts to be allelopathic against test species. The use of hot water extract is unnatural, but it reduces the time for extraction of allelochemicals. Similarly, it caused significant repression in plumule length of all three varieties and radical of Pahari and Iqbal.

Above mentioned results were like the inhibitory effect of the aqueous extracts from other species like *Dodonaea viscosa* (1), *Cenchrus ciliaris* and *Bothriochloa pertusa* (10), *Anagallis arvensis* (20), *Azadirachta indica* A Juss (20), *Tamarindus indica* L. (18), *Broussonetia papyrifera* (11) and *Lactuca sativa* (4), which support our findings. Other researchers like (16, 12 and 6) investigated similar phytotoxicity for other plant species, which further strengthened the results.

IV. CONCLUSION

Current work led us to the conclusion that the extracts obtained from powder of leaves of *Papaver somniferum* caused inhibitory effect on *Zea mays* varieties Azam, Pahari and Iqbal. Increasing soaking duration can be more inhibitory. Since the extract reduced the plumule and radical in aqueous extract treatment (24h and 48h). Also, in mulching, litter, and hot water treatments.

V. CONFLICT OF INTEREST STATEMENT

On behalf of all authors, the corresponding author states that there is no conflict of interest.

REFERENCES

- [1] Barkatullah FH, and Ibrar M. 2010. Allelopathic potential of *Dodonaea viscosa* (L.) JACQ. Pak. J. Bot. 42(4), 2383-2390.
- [2] Basharat A. Dar, Saud L. Al-Rowaily, Abdulaziz M. Assaeed, Magdy I. El-Bana, Ahmed K. Hegazy and Jahangir A. Malik, 2017. Allelopathic potential of *Argemone ochroleuca* from different habitats on seed germination of native species and cultivated crops. Pak. J. Bot., 49(5): 1841-1848
- [3] Bhowmik PC, Doll JD. 1992. Corn and soybean response to allelopathic effects of weed and crop residues. Agron. J. 74, 601–606.
- [4] Chon SU, Jang HG, Kim DK, Kim YM, Boo HO, Kim. YJ 2005. Allelopathic potential in lettuce (*Lectuca sativa* L.) plant. Scientia Horticultorea 206(3), 309-317.
- [5] Chung IM, Kim JT, Kim S. 2007. Evaluation of allelopathic potential and quantification of Momilactone, b from rice hull extracts and assessment of inhibitory bioactivity on paddy field weeds. J. Agric. Food Chem. 54(7), 2527-2536.
- [6] Elizabeth GB, Pivello VR, Meirelles ST. 2008. Allelopathic evidence in *Brachiaria decumbens* and its potential to invade the Brazilian Cerrados. Braz. Arch. Biol. Technol. 51(4), 825-831.
- [7] Ferguson JJ, Rathinasabapathi. 2009. Allelopathy: How Plants Suppress Other Plants. University of Florida, IFAS Extension. Horticultural Science 1 – 4.
- [8] Fujii, Y., M. Furukawa, Y. Hayakawa, K. Sugawara and T. Shibuya. 1991. Survey of Japanese medicinal plants for the detection of allelopathic properties Weed Res. Japan., 36: 36-42
- [9] Fujii, Y., T. Shibuya and T. Yasuda. 1990. Survey of Japanese weed and crops for the detection of water extractable allelopathic chemicals using Richard’s function fitted to lettuce germination test. Weed Res. Japan., 35: 362-370.
- [10] Hussain F, Ahmad B, Ilahi I. 2010. Allelopathic effects of *Cenchrus ciliaris* L. and *Bothriochloa pertusa* (L.) a. campus. Pak. J. Bot. 42(5), 3587-3604.
- [11] Hussain F, Niaz F, Jabeen M, Burni T. 2004. Allelopathic potential of *Broussonetia papyrifera* Vent. Pak. J. Pl. Sci. 10(2), 69-77.
- [12] Hussain S, Siddiqui SU, Khalid S, Jamal A, Qayyum A, Ahmad Z. 2007. Allelopathic potential of *Senna* (*Cassia angustifolia* Vahl.) on germination and seedling characters of some major cereal crops and their associated grassy weeds. Pak. J. Bot. 39(4), 1145-1153.
- [13] Inderjit K, Keating L. 1999. Allelopathy: principal and practice. John promises for biological control in advance in agronomy, (Eds). Sparks D.L., Academic Press, 67, 141-231.
- [14] Inderjit, Duke SO. 2003. Eco physiological aspects of allelopathy. Planta 217(4), 529-539.
- [15] Kaul S, Bansal GL. 2002. Allelopathic effect of *Ageratina adenophora* on growth and development of *Lantana camara*. Ind. J. Pl. Physiology 7(2), 195-197.
- [16] Marwat KB, Khan MA. 2006. Allelopathic proclivities of tree leaf extracts on seed germination and growth of wheat and wild oats. Pak. J. Weed Sci. Res. 12(4), 265-269.
- [17] Peneva A. 2007. Allelopathic effect of seed extracts and powder of coffee (*Coffea arabica* L.) on common cocklebur (*Xanthium strumarium* L). Bul. Jour. Agr. Sci. 13, 205-211.
- [18] Pervez SS, Pervez MM, Nishihara E, German H, Fujii Y. 2003. *Tamarindus indica* L., leaf is a source of allelopathic substance. Plant Growth Regulator 40(2), 107-115.
- [19] Putnam, A.R. and W.B. Duke. 1974. Biological suppression of weeds evidence for allelopathy in accessions of cucumber Sci., 185:370-71
- [20] Rebaz Z, Shaukat SS, A.Siddiqui I. 2001. Allelopathic potential of *Anagalis arvensis* L.: A Cosmopolitan weed,.Pak. J. Biol. Sci. 4(4), 446-450.
- [21] Samreen U, Hussain F, Sher Z. 2009. Allelopathic potential of *Calotropis procera* (Ait.) Ait. Pak. J. Pl. Sci. 15(1), 7-14.
- [22] Xuan TD, Tsuzuki E, Hiruki T, Mitsuhiro M, Khanh TD, Chung IM. 2004. Evaluation of phytotoxicity of neem (*Azadirachta indica* A. Juss) to crops and weeds. Crop Protection 23(4), 335- 345.

AUTHORS

First Author – Sidra-Tul-Muntaha, Department of Botany, Govt. Frontier College for Women Peshawar, Pakistan
Second Author – Sayeda Sarah Mubarak, Department of Botany, Govt. Frontier College for Women Peshawar, Pakistan
Third Author – Fatima Khan Sherwani, Department of Botany, Govt. Frontier College for Women Peshawar, Pakistan
Fourth Author – Nazma Azim, Department of Botany, Govt. Frontier College for Women Peshawar, Pakistan

Corresponding author’s email: sidtmun99@gmail.com

TABLES

Treatment 24h	Azam	Pahari	Iqbal	Treatment 48h	Azam	Pahari	Iqbal
Germination							
Control	100	93.3	100	Control	100	93.3	100
Test	93.3	100	100	Test	100	100	93.3
Radical							
Control	33.66	26.73	34.83	Control	33.66	26.73	34.83
Test	20.77	30.2	17.33	Test	24.13	17.27	16.30
Plumule							
Control	5.40	2.57	0.40	Control	5.40	2.57	0.40

Test	0.00	2.46	0.00	Test	0.50	1.46	1.00
Fresh weight							
Control	3.13	2.33	1.73	Control	3.13	2.33	1.73
Test	3.40	1.73	2.33	Test	2.73	1.66	2.00
Dry weight							
Control	1.66	1.60	1.13	Control	1.66	1.60	1.13
Test	1.73	0.73	1.66	Test	1.66	1.06	1.66
Moisture content							
Control	88.40	70.88	67.40	Control	88.40	70.88	67.40
Test	99.10	141.67	39.33	Test	64.80	66.67	19.90

Table no. 1: Effect of Aqueous extract of leaves of *Papaver somniferum* on percent germination, plumule and radical growth (mm), Fresh and dry weight (mg), and percent moisture contents of *Zea mays* varieties Azam, Pahari and Iqbal. Each value is a mean of three replicates, each with 5 seeds

Treatment	Variety	Variety		
		Azam	Pahari	Iqbal
Control	Germination	100	100	86.67
	Test	73.3	100	100
Control	Plumule	55.76	66.93	62.96
	Test	52.66	64.03	31.1
Control	Radical	124.20	117.33	94.87
	Test	98.87	93.97	51.80
Control	Fresh weight	2.53	2.66	3.53
	Test	2.40	3.06	3.53
Control	Dry weight	1.40	1.40	1.51
	Test	1.53	1.60	1.57
Control	Moisture content	84.10	90.27	103.70
	Test	57.13	94.80	123.63

Table no. 2: Effect of mulch of leaves of *Papaver somniferum* on percent germination, plumule and radical growth (mm), Fresh and dry weight (mg), and percent moisture contents of *Zea mays* varieties Azam, Pahari and Iqbal. Each value is a mean of three replicates, each with 5 seeds.

Treatment	Variety	Variety		
		Azam	Pahari	Iqbal
Control	Germination	100	100	93.3
	Test	100	100	100

	Plumule			
Control		3.00	5.46	2.66
Test		4.26	3.33	1.06
	Radical			
Control		26.93	32.00	29.60
Test		30.60	29.00	28.86
	Fresh weight			
Control		1.26	1.53	1.73
Test		2.46	2.20	1.40
	Dry weight			
Control		0.73	1.06	1.06
Test		1.40	1.60	1.06
	Moisture content			
Control		77.76	43.33	46.16
Test		77.76	44.50	51.12

Table no. 3: Effect of litter of leaves of *Papaver somniferum* on percent germination, plumule and radical growth (mm), Fresh and dry weight (mg), and percent moisture contents of *Zea mays* varieties Azam, Pahari and Iqbal. Each value is a mean of three replicates, each with 5 seeds.

Treatment	Variety	Germination		
		Azam	Pahari	Iqbal
Control		100	100	93.3
Test		100	100	100
	Plumule			
Control		3.26	5.20	1.06
Test		1.26	1.00	0.60
	Radical			
Control		27.26	31.13	29.40
Test		33.40	26.73	24.46
	Fresh weight			

Control	1.20	1.40	1.69
Test	2.63	2.46	2.67
	Dry weight		
Control	0.73	0.66	1.13
Test	1.7	1.86	2.33
	Moisture content		
Control	65.27	113.87	43.00
Test	52.05	20.10	25.90

Table no. 4: Effect of hot water extract of leaves of *Papaver somniferum* on percent germination, plumule and radical growth (mm), fresh and dry weight (mg), and percent moisture contents of *Zea mays* varieties Azam, Pahari and Iqbal. Each value is a mean of three replicates, each with 5 seeds.

FIGURES



Fig 1: showing germinated varieties in aqueous extract treatment.




Fig 2: Showing the germinated varieties in mulch treatment. The emerging plumule can be seen clearly.