

PHYTOCHEMICAL STUDIES OF *ACMELLA CAULIRHIZA* AND *SPERMACOCE PRINCEAE* USED BY POSTPARTUM MOTHERS IN NYAMIRA COUNTY, KENYA

¹Jespher Nyaboke Onyango²Jared Misonge Onyancha, ³Dr. Jacksn Odhiambo Onyuka⁴Prof. John Memba Ochora, PhD.⁵Patrick Ogembo Getonto, ⁶ Charles Wambugu Maina

¹ Medical Laboratory Department, Mount Kenya University

² Pharmacognosy Department, Mount Kenya University

³ Medical Laboratory Department, Mount Kenya University

⁴ Botany Department, Jomo Kenyatta University of Agriculture and Technology

⁵ Department of Community Health Nursing, Mount Kenya University

⁶ Muranga County Referral Hospital

Abstract- Introduction: Traditional medicine have been used in health maintenance, disease prevention and treatment for example *Acmella caulirhiza* used to treat a child's mouth sores and *Spermacoce princeae* used to accelerate healing of umbilical cord and to clean the system after birth.

Objective: The main objective of the present study was to determine phytochemical compounds of *A. caulirhiza* and *S. princeae* used by postpartum mothers in Nyamira County, Kenya.

Methodology: The study area was Nyamira County where the two plant specimens were collected. Plant materials were identified at East Africa Herbarium. Plant specimens were transported to M.K.U. Pharmacognosy laboratory where processing was done. Phytochemical analysis methods were employed to determine phytochemicals compounds in the crude plant extracts. Data was stored in Excel spread sheet in a personal computer protected with a password. Data was presented using tables and photographs.

Results: Phytochemical examinations revealed that *Acmella caulirhiza* contains flavonoids, terpenoids, coumarins and sterols compounds. On the other hand, *Spermacoce princeae* contains flavonoids, terpenoids, tannins, saponin alkaloid and glycoside compounds.

Conclusion and recommendation: The plants may be used in treating puerperal sepsis although commercially available drugs are recommended as they are highly effective. The two plants can be a potent source of complementary and modern medicine. Further research is recommended to isolate and identify pure compounds of the two plants

Index Terms- Traditional medicine, phytochemical investigation, Asteraceae, Rubiaceae.

I. INTRODUCTION

Traditional medicine has been used in health maintenance, disease prevention and treatment (WHO, 2014). The use of medicinal plants has been known to mankind as the oldest practice of healthcare (Yogayata and Vijay, 2012). Nowadays, isolation and characterization of biologically active compounds from medicinal plants continues and drug discovery techniques have been applied to the standardization of herbal medicines, to elucidate analytical marker compounds (Marcy and Douglas, 2005). Medicinal plants used to manage postpartum complications include; *Basella alba* plant which belongs to Basallaceae family, is used to manage stomachache, stimulate milk production and is used to remove the placenta after birth (Jeruto *et al.*, 2015). *Toddalia asiatica* and *Pentas longiflora* species which belongs to Rutaceae and Rubiaceae family respectively. The leaves of these herbs are used to manage Urinary tract infections (Jeruto *et al.*, 2015).

Acmella caulirhiza is similarly known as *Spilanthes acmella*. It is a flowering herbal plant, which belongs to Compositae/Asteraceae family (Berhane *et al.*, 2014). It is an annual or perennial herb. Locally it is known as Ekenyunyuntamonwa (Ekegusii) and Ajuok-olwa Salamatwe (Dholuo) (Kokwaro, 2009). It is used by different communities in Kenya and the rest of Africa to treat various medical conditions. Example in Kenya, its flowers and leaves are used to treat venereal diseases (Jeruto *et al.*, 2015). It is used to relieve painful sores of the mouth, gums and throat, as well as stomach ache (Kokwaro 2009). Also it is used to treat decayed teeth, gingivitis or wounds in the mouth, toothache and sore throat (Kipruto *et al.*, 2013). The Zulu people of South Africa use *A. caulirhiza* as a local analgesic for toothache and to ease sensitivity of gums during dental extractions (Crouch *et al.*, 2005).

Spermacoce princeae is a flowering herbal plant which belongs to the family Rubiaceae (Augustin *et al.*, 2015). Locally it is known as Omoutakiebo (Ekegusii), Gakungathe (kikuyu), Murkugwet (kipsigis) and Nyamwoch (Dholuo) (Kokwaro, 2009). It grows in tropical regions and it is used extensively. Normally it is used by different communities in Kenya and the rest of Africa to treat

several diseases. Just to mention a few, leaves and roots are used to treat chronic asthma, cancer, mastitis in cows and venereal diseases by the Nandi people in Kenya (Jeruto *et al.*, 2011). Another study carried out in Vihiga County, Kenya found that, cold infusion is made from leaves and drunk in the treatment of diarrhea (Antony *et al.*, 2016). Leaves and stem are used to treat female infertility in Baham, Cameroon (Telefo *et al.*, 2011). In Cameroon, leaves of *Spermacoce princeae* are warmed on fire, ground and mixed with red oil and salt, then is taken orally in treatment of kidney disease (Focho *et al.*, 2009).

II. MATERIAL AND METHODS

2.1 Study Area

The study site was North Mugirango and West Mugirango constituencies of Nyamira County. The study points in West Mugirango constituency were; Sirona (0° 33'14.8536 S and 34° 58' 2.4996 E), Bonyunyu (0° 31'36.2532 S and 34° 53'20.4108 E) and miruka (0° 29'13.902 S and 34° 53'20.3208 E) whereas the study point in North Mugirango constituency was; Magong'a (0° 28' 46.7724 S and 34° 57'6.4836 E) in Nyamira County. In this County, local inhabitants regularly use medicinal plants for personal and domestic animal health. Local inhabitants in this County, follow traditional beliefs and customs. Further, most inhabitants living in this area have a tendency of harvesting the medicinal plants from undisturbed vegetation. This is due to the fact that many plant species grow in the study region (Omwenga *et al.*, 2015). Postnatal mothers use *Acmella caulirhiza* and *Spermacoce princeae* to treat child sores and to clean reproductive system respectively in women after birth. Nyamira County is one of highly populated area with approximately 912.5 Km² with a population of 598,252 and a population density of 656 persons per Km² according to (KNBS, 2009).

2.2 Plant materials collection

Acmella caulirhiza and *Spermacoce princeae* medicinal plant specimen were collected from West Mugirango and North mugirango constituencies in Nyamira County with acceptable bio-conservation methods (WHO, 2003a). Harvesting was done in a dry weather morning after the dew had evaporated (Prajapati *et al.*, 2010). The two specimens were carried separately in gunny bags and transported to Pharmacognosy Laboratory of Mount Kenya University within 72 hours of collection (WHO, 2003a).

Fig. 1: The two medicinal plant materials collected



2.3 Processing of plant materials

Processing was done within 72 hours after collection. Herbarium preparations were established and the voucher specimens were processed in duplicate. They were mounted on herbarium sheets, pressed to flatten, to dry and were labeled. Voucher specimen (Number JN001 and JN002) were identified at East African Herbarium in the National Museums of Kenya on basis of morphological characteristics and compared with the voucher specimens recorded in East Africa Herbarium. Voucher specimen (Number JN001 and JN002) were deposited at Mount Kenya University Botanical Herbarium Laboratory in the school of Pharmacy. The collected materials were washed thoroughly with tap water and then air dried under a shade at room temperature for one week. When dried, the plant materials (*A. caulirhiza* and *S. princeae*) were ground into coarse powder using a porcelain mortar and pestle (Hena, *et al.*, 2010). The coarse powder materials were labeled and stored in brown paper bags under a dry condition, away from light at room temperature till the time of extraction and phytochemical screening (Prajapati *et al.*, 2010).

2.4 Plant extraction using organic solvents

Using a top loading Weighing Electronic Balance (Models TP-B 2000), 50 grams of the Kenyan *Acmella caulirhiza* and 50 grams *Spermacoce princeae* each powder was weighed separately and transferred into separate conical flasks, labeled with the

constituency of collection, plant species and date. Then 500mls of 100% Ethyl Acetate (Loba Chemie Company Lot#L157601502) was added to cover each plant materials and covered with a stopper, then macerated in the solvent at room temperature for 48 hours with intermittent agitation. Using a funnel and Whatman filter paper No. 1 the crude extracts from each of the plant materials were strained separately into glass reagent bottles then covered with stoppers. The process was repeated with 500mls of 100% Ethanol Analar Normapur (VWR Prolabo Company Batch 12D250511) and Methanol (Loba Chemie Company Lot #B193331604). The filtrates were labeled and concentrated in a rotary evaporator at 40 degree Celsius for Ethyl acetate, 60 degree Celsius for Ethanol and Methanol respectively. Using analytical balance, empty beakers were weighed, the extracts from the distillation flask were transferred into them, labeled appropriately and the solvents were evaporated in an Oven set at appropriate temperature. Quantity of each crude plant extract paste was calculated by the formula: Plant crude residue = (weight of beaker + extract) - (weight of empty beaker). The extracted paste of each plant species examined was kept in beakers covered in a refrigerator a waiting for bioactivity assay (Afolayan *et al.*, 20008).

2.5 Aqueous extraction of crude plant material

Aqueous extracts of *Acmella caulirhiza* and *Spermacoce princeae* was made from crude plant material according to Bibi *et al.*, (2012) by weighing 20 grams of *Acmella caulirhiza* and 20 grams of *Spermacoce princeae*. They were boiled separately in 400mls distilled water in beakers of 400ml capacity on Hot Plate set at 100⁰ C for 5 minutes. The extracts were cooled, using a funnel and Whatman filter paper (No. 1) they were filtered and freeze dried according to Pikal *et al.*, (2010), to extract dry powders from the aqueous solutions of the two plants. Freeze-drying was done in the following steps; freezing, primary drying and secondary drying. Primary drying involves; evacuating the system, increasing shelf temperature resulting to product temperature 2–3⁰ C below collapse temperature. Secondary drying involves; removing unfrozen water from the solute phase by desorption through raising temperatures. The dry and lyophilized extracts were weighed and stored in a freezer for bioactivity testing (Bibi *et al.*, 2012).

2.6 Phytochemical investigation

Phytochemical analysis for various secondary metabolites of the plant extract was done using methods described below. The groups of phytochemical compounds investigated were alkaloids, flavonoids, tannins, coumarins, steroids, sterols, saponins, terpenoids and glycosides.

Test for tannins

Ferric Chloride Test: *Acmella caulirhiza* 0.5 grams and 0.5gms *Spermacoceprinceae* dry powdered samples were boiled in 10ml distilled water on a hot plate separately and the extracts were filtered. 2ml portion of each filtrate was measured and 3drops of 0.1% ferric chloride solution was added. Formation of a green colored solution shows tannins presence (Kiran and Prasad 2015).

Test for coumarins

Onto separate filter papers, a few drops of ammonia were added and then a drop of *Acmella caulirhiza* and *Spermacoceprinceae* extracts were added separately. Fluorescence on the paper shows coumarins presence (Sangeetha *et al.*, 2014).

Test for flavonoids

Ferric chloride test; 2ml *Acmella caulirhiza* and 2ml *Spermacoceprinceae* crude extracts were separately treated with 5 drops of Ferric chloride solution each. Formation of a blackish red colour shows flavonoids presence (Kiran and Prasad 2015).

Tests for steroids

Liebermann test: To 1ml *Acmella caulirhiza* and 1ml *Spermacoceprinceae* extract solution in 10ml chloroform solution separately, 3drops of acetic anhydride and Sulphuric acid solutions were added slowly from the side of test tube separately. Observation of a brown ring at the junction of the two layers and the upper layer turns green separating the liquids shows steroids presence (Kiran and Prasad, 2015).

Test for sterols

Liebermann test: Few grams of *Acmella caulirhiza* and *Spermacoceprinceae* in separate test tubes were dissolved in 0.5 ml hot acetic anhydride and 0.5 ml of chloroform was added. Observations of blue-green ring at interphase a positive reaction (Sabri *et al.*, 2012).

Testing for saponins

Froth test: Using a top loading weigh balance, 0.5 grams *Acmella caulirhiza* and 0.5gms *Spermacoceprinceae* powder were weight, placed in separate test tubes and 10ml distilled water added to each. Then themixture was shaken and left to stand. Persistent froth shows saponins presence (Kiran and Prasad 2015).

Test for terpenoids

Five ml *Acmella caulirhiza* and 5ml *Spermacoceprinceae* extracts were mixed separately in 2ml chloroform and then 3ml concentrated sulphuric acid was added. At the interface a reddish brown color shows terpenoids presence (Ablude, 2001).

Test for glycosides

Keller-kiliani test; *Acmella caulirhiza* (50mg), and 50mg *Spermaceocephalae* extracts in separate test tubes, 2ml and 1ml glacial acetic acid and ferric chloride solution were added respectfully. The contents were heated and cooled then transferred to a test tube containing 2ml concentrated sulphuric acid. Formation of a brown ring at the interface (presence of deoxy sugar characteristic of cardenolides) and observation of pale green colour in the upper layer (steroidal nucleus) shows glycoside presence (Kiran and Prasad 2015).

Test for alkaloids

Using a top loading weigh balance, 0.5gms of *Acmella caulirhiza* and 0.5gms *Spermaceocephalae* dried powdered sample free extract were measured, agitated with 3mls diluted hydrochloric acid and filtered separately. Then the filtrates were tested for alkaloids using dragendorff's reagent: To 2mls of *Acmella caulirhiza* and *Spermaceocephalae* filtrate in separate test tubes, 2 mL Dragendorff's reagent was added to each. A prominent yellow precipitate shows presence of alkaloids (Harborne, 2005).

2.7 Data Analysis and Presentation

Data was stored in Excel spread sheet in a personal computer protected with a password. Also a flash disk secured with a password was used as a backup. Data was presented using tables and photographs.

2.8 Ethical Considerations

Ethical clearance was obtained from Mount Kenya University Ethical Review Committee and NACOSTI before commencement of the study. Phytochemical analysis was conducted in Mount Kenya University Pharmacognosy laboratory.

III. FINDINGS

Quantity of crude plant extracts

Table 1: % yields of extract obtained by different extraction solvent

Percentage yield					
Plant species	Part used	Ethyl Acetate	Ethanol	Methanol	Aqueous
<i>Acmella caulirhiza</i>	Whole plant	1.68 (2.1%)	3g (3.75%)	4g (5%)	2.45g (12.25%)
<i>Spermaceocephalae</i>	Whole plant	2g (2.5%)	4g (5%)	5(6.25%)	3g (15%)

Fig. 2: Preliminary phytochemical screening of *Acmella caulirhiza* whole plant



Fig. 3: Preliminary phytochemical screening of *Spermacoce princeae* whole plant



Determination of Phytochemical Compounds in Crude Plant Extracts of *A. caulirhiza* and *S. princeae*

Phytochemical examinations revealed that *Acmella caulirhiza* contains flavonoids, terpenoids, coumarins and sterols compounds. However, tannins, saponin and glycoside were absent. On the other hand, *Spermacoce princeae* contains flavonoids, terpenoids, tannins, saponin alkaloid and glycoside compounds. However, sterols and coumarins were not found.

Table 2: Phytochemical Analysis of *Acmella caulirhiza* and *Spermacoce princeae* from Nyamira County

Phytochemical compounds	Flavonoids	Terpenoids	Coumarins	Sterols	Tannins	Saponins	Alkaloids	Glycoside	Steroid s
<i>Acmella caulirhiza</i>	+	+	+	+	-	-	-	-	-
<i>Spermacoce princeae</i>	+	+	-	-	+	+	+	+	-

IV. DISCUSSION

Medicinal plants usage is increasingly popular among the Gusii community of Nyamira County. Many medicinal plants grow around the homestead and have been used naturally for many years by traditional healers to control common health problems.

Determination of Phytochemical Compounds in Crude Plant Extracts of *Acmella caulirhiza* and *Spermacoce princeae*

Phytochemical examinations revealed that, *Acmella caulirhiza* contains flavonoids, terpenoids, coumarins and sterols phytochemical compounds. However, tannins, saponin and glycoside were absent. The presence of coumarin compound in this plant is consistent with other study Chhabra *et al.*, (1989). Other findings indicates that, *A. caulirhiza* leaves and flowers contain spilanthol compounds believed to prevent bacterial pathogens in addition to numbing the pain when used in toothache therapy (Neil and Jerald, 2005). Besides, *A. caulirhiza* plant is believed to treat sore throat and stomachache traditionally (Kipruto *et al.*, 2013; Giday *et al.*, 2010; Njoroge and Bussmann, 2006; Chhabra *et al.*, 1989).

On the other side, *Spermacoce princeae* contained flavonoids, terpenoids, tannins, saponin alkaloid and glycoside phytochemical compounds. However, sterols and coumarins were not found. This result concurs with the phytochemical results reported by Jeruto *et al.*, (2011) on phytochemical compounds present in this plant. *S. princeae* is used traditionally to clean reproductive system after birth. Other findings have shown that *S. princeae* treats venereal diseases, pneumonia, typhoid, Chronic asthma, cancer, wounds, eye

problems, venereal diseases, diarrhoea skin and kidney diseases (Jeruto *et al.*, 2011). This validates therapeutic value of this plant, in traditional management of typhoid, pneumonia and eye infections (Jeruto *et al.*, 2011).

Therapeutic value of medicinal plants is determined by the presence of phytochemical compounds having certain functional and pharmacological activity (Geeta *et al.*, 2012). Flavonoids compounds have been reported to have antimicrobial activity (Pandey *et al.*, 2010; Cowan, 1999). Example flavonoids such as flavone and flavonol glycosides, apigenin, isoflavones, galangin flavanones, and chalcones have been revealed to possess effective antibacterial activity (Cushnie and Lamb, 2005). A new flavanone, 7-hydroxy-6,8-dimethoxyflavanone, showed anti-mycobacterial activity against *M. tuberculosis* H37Ra at 50 µg/mL MIC value (Prawat *et al.*, 2013). Mode of action results from their ability to inactivate microbial cell envelope transport proteins, enzymes and adhesins (Kumar and Pandey 2013; Mishra *et al.*, 2009; Cowan, 1999). Lipophilic flavonoids may disrupt microbial membranes (Mishra *et al.*, 2009; Cowan, 1999). Flavonoids inhibit cytoplasmic membrane function and they inhibit DNA gyrase and hydroxyacyl-acyl carrier protein dehydratase activities (Zhang *et al.*, 2008).

Terpenoids have antimicrobial activity against bacterial and fungi (Omojate *et al.*, 2014; Ghoshal *et al.*, 1996). Example petalostemumol terpenoid exhibited excellent effects against *Bacillus subtilis* and *Staphylococcus aureus* (Omojate *et al.*, 2014). They also control *Listeriamonocytogenes* Cowan, (1999). Terpenoids derived from essential oil inhibitory 60% fungi while they inhibit 30% bacteria (Omojate *et al.*, 2014; Mohd *et al.*, 2014). Terpenoids act by disrupting cellular membrane by lipophilic compounds (Cowan, 1999; Hatice and Ayse 2014). Coumarin, has been reported to contain antimicrobial activity against *Candida albicans* and chronic infections (Thornes *et al.*, 1982). Example coumarin compound isolated from *Angelica lucida* L. is effective against oral pathogens such as *Streptococcus viridians* and *Streptococcus mutans* (Widelski *et al.*, 2009). They act by activating other cells of immune system stimulating macrophages, indirectly affecting the disease causing agents (Casley *et al.*, 1997).

Sterols have been reported to contain antimicrobial activity. Examples stem sterols of *Withania somnifera* is reported to have antimicrobial effects against *P. aeruginosa*. As well stem sterol of *Euphorbia hirta* and *Terminalia chebula* is active against *S. aureus* (Geeta *et al.*, 2012). There is no information on antimicrobial mode of action of plant steroids and sterols. Tannins have been reported to contain bactericidal activity against gram positive organisms such as *S. aureus* and used for the treatment of diarrhea an example of enteric diseases (Omojate *et al.*, 2014; Chung *et al.*, 1993). Tannins have been used as antiseptic whose activity is due to manifestation of the phenolic group. Tannin-rich plants have been shown to be toxic to filamentous fungi, yeasts, and bacteria (Omojate *et al.*, 2014). Example tannins extracted from *Solanum trilobatum* plant have been reported to possess high antibacterial activity against *Staphylococcus aureus* and *Proteus vulgaris* at 2.5mg/ml concentration (Doss *et al.*, 2009). Tannin compounds act by interfering with bacterial cell wall causing disintegration of bacterial colonies thus inhibiting microbial growth (Erasto *et al.*, 2004). They also act by forming complexes with proteins and polysaccharide through hydrogen bonding, hydrophobic bond and covalent bond resulting to inactivation of microbial adhesion, enzymes and cell envelope transport proteins (Haslam, 1996; Haslam *et al.*, 1988). Also they have the ability to inactivate microbial adhesins, enzymes, cell envelope and transport proteins (Omojate *et al.*, 2014). Condensed tannins have been confirmed to bind to cell walls of ruminal bacteria, thus inducing bacterial stasis and protease activity (Jones *et al.*, 1994).

Saponins have been reported to have antimicrobial activity (Moyo *et al.*, 2012). Example Yucca saponin have antibacterial activity against *Staphylococcus aureus* and *Lactobacillus* (Tanaka *et al.*, 1996). Mode of action is by altering with the permeability of cell walls hence they exert toxicity on all tissues. Also they form complexes with cell membranes hence elicit changes in cell morphology leading to cell lysis (Moyo *et al.*, 2012).

Alkaloids have antibacterial properties against gram-positive and gram-negative bacteria (Donald *et al.*, 2016). Example alkaloids Chloroform and ethanol extracts from *Callistemon citrinus* leaves have been reported to have antibacterial activity against *Staphylococcus aureus*, *E. coli*, *S. typhi* and *P. aeruginosa* via disc diffusion methods (Krishna *et al.*, 2012; Donald *et al.*, 2016). They act by disrupting cell wall membrane (Cowan, 1999). Glycosides: have been reported to contain antibacterial activity. Example glycosides compounds extracted from *Caesalpinia coriaria* (Jacq) Willd exhibited antibacterial activity against *E.coli*, *Staphylococcus aureus* and *Klebsiella pneumonia* (Anandhi *et al.*, 2014). Glycosidic compounds (G1) extracted from *Citrus laurantiifolia* L. fruits exhibited broad spectrum antibiotic effects against *Staphylococcus aureus*, *Streptococcus*, *Pseudomonas aeruginosa* and *Escherichia coli* (Sameerah *et al.*, 2013). Mode of action glycoside act by causing leakage of cellular materials by breaking the outer membrane. Also glycosides inhibit the respiration and growth of pathogenic micro-organisms by entering the inner membrane and inactivating the enzyme system dehydrogenase (Anandhi *et al.*, 2014).

V. CONCLUSION AND RECOMMENDATION

Demonstration of phytochemical compounds in the two plants such as flavonoids, terpenoids, coumarin, sterols, tannins, saponin, alkaloids and glycosides validates their use in management of puerperal sepsis. Mode of action of compounds present in the two plants indicates that the plants have therapeutic potential. Phytochemical compounds analysis lays a foundation for ethnobotanical and pharmacological investigations for new drug discovery.

A. caulirhiza and *S. princeae* plants may be used in treating puerperal sepsis. The two plants may be used as medicine and can be a potent source of complementary and modern medicine. Further research is recommended to isolate and identify pure compounds of the two plants

REFERENCES

- [1] **Ablude, F. O. (2001)**. Mineral and phytate content of ten vegetable grown in Nigeria and calculation of their phytate; Zn and Ca: phytate molar ratios, "Adv. Food Science"23:36-39
- [2] **Afolayan, A. J., Ncube, N. S. and Okoh, A. I.(2008)**. Assessment techniques of antimicrobial properties of natural compounds of plant origin: Current methods and future trends. *African Journal of Biotechnology*7 (12): 1797-1806.
- [3] **Augustin, N. Donatien, G., Simeon, P. C. F. and Huguette, N. M. (2015)**. Toxicological Evaluation of the Aqueous Leaf Extract of *Spermacoce princeae* (Rubiaceae): A Traditional Antibacterial Preparation. *International Journal of Toxicological and Pharmacological Research* 7(3); 123-129.
- [4] **Anandhi, D., Srinivasan, P.T., Praveen, Kumar G. and Jagatheesh, S. (2014)**. Influence of flavonoids and glycosides from *Caesalpinia coriaria* (Jacq) wild as bactericidal compound. *Int.J.Curr.Microbiol.App.Sci* 3(4): 1043-1051
- [5] **Antony, O. R., Michael, K., Makokha, A. O. and Festus, M. T. (2016)**. Types of Herbal Medicine Used for HIV Conditions in Vihiga County, Kenya. *European Journal of Medicinal Plants* 13(2): 1-23
- [6] **Berhane, K., Tinde, A., Laurentius, J. M. and Zemedu, A. (2014)**. Use and management of traditional medicinal plants by Maale and Ari ethnic communities in southern Ethiopia. *Journal of Ethnobiology and Ethnomedicine*, 10:46
- [7] **Bibi, Y., Nisa, S., Zia M., Waheed, A., Ahmed, S. and Chaudhary, F. M.(2012)**. *In vitro* cytotoxicity activity of *Aesculus indica* against breast carcinoma cell lines (MCF-7) and phytochemical analysis. *Pak. J. Pharm. Sci.*, 25 (1): 183-187
- [8] **Casley-Smith, J. R. and Casley-Smith, J. R. (1997)**. Coumarin in the treatment of lymphoedema and other high-protein oedemas, p. 348. *In* R. O'Kennedy and R. D. Thornes (ed.), *Coumarins: biology, applications and mode of action*. John Wiley & Sons, Inc., New York, N.Y.
- [9] **Chhabra, S. C., Mahunnah, R. L. A. and Mshiu, E. N. (1989)**. Plants used in traditional medicine in eastern Tanzania: II. Angiosperms (Capparidaceae to Ebenaceae) *Journal of Ethnopharmacology*, 25 (1989), pp. 339-359
- [10] **Chung, K. T., Stevens, S. E., Lin, W. F. and Wei, C. I. (1993)**. Growth inhibition of selected food-borne bacteria by tannic acid, propyl gallate and related compounds. *Letters in Applied Microbiology* 17, 29-32.
- [11] **Crouch, L. A., Mulholland, D. A. and Nair, J. J. (2005)** A novel alkylamide from the leaves of *Acmella caulirhiza* (Asteraceae), a traditional surface analgesic. *South African journal* 71(2): 228-230
- [12] **Cushnie, T. P. T. and Lamb, A. J. (2005)**. "Antimicrobial activity of flavonoids," *International Journal of Antimicrobial Agents*, vol. 26, no. 5, pp. 343-356.
- [13] **Cowan, M. M. (1999)**. Plant products as antimicrobial agents. *Clin Microbiol Rev.* 12(4):564-82.
- [14] **Donald, M., Tariro, C., and Stanley, M. (2016)**. Antibacterial Properties of Alkaloid Extracts from *Callistemon citrinus* and *Vernonia adoensis* against *Staphylococcus aureus* and *Pseudomonas aeruginosa*. *International Journal of Medicinal Chemistry* Volume 2016 Article ID 6304163, page 7
- [15] **Doss, A., Mohammed, M. H. and Dhanabalan, R. (2009)**. Antibacterial activity of tannins from the leaves of *Solanum trilobatum* Linn. *Indian Journal of Science and Technology* Vol.2 No 2.
- [16] **Erasto, P., Bojase-Moleta, G. and Majinda, R. R. T. (2004)**. Antimicrobial and antioxidant flavonoids from the roots wood of *Bolusathus spesiosus*. *Phytochem.* 65, 875-880.
- [17] **Focho, D. A., Ndam W. T. and Fonge, B. A. (2009)**. Medicinal plants of Aguambu – Bamumbu in the Lebiale highlands, southwest province of Cameroon. *African Journal of Pharmacy and Pharmacology* Vol. 3 (1). pp. 001-013.
- [18] **Geeta, S., Padma, K. and Alka, J. (2012)**. Antibacterial Potential of Sterols of some Medicinal Plants. *International Journal of Pharmacy and Pharmaceutical Sciences*. Vol 4, Issue 3,
- [19] **Ghoshal, S., Krishna, B. N., Prasad, and Lakshmi, V. (1996)**. Antiamoebic activity of *Piper longum* fruits against *Entamoeba histolytica* in vitro and in vivo. *J. Ethnopharmacol.* 50:167-170.
- [20] **Giday, M., Asfaw, Z. and Woldu, Z. (2010)**. Ethnomedicinal study of plants used by Sheko ethnic group of Ethiopia. *J. Ethnopharmacol.* 132 (1):75-85.
- [21] **Harborne, J. B. (2005)**. *Phytochemical Methods*, Springer (India) Pvt. Ltd, New Delhi, 17
- [22] **Haslam, E. (1996)**. Natural polyphenols (vegetable tannins) as drugs: possible modes of action. *J. Nat. Prod.* 59:205-215.
- [23] **Haslam, E. Ya, C., Gaffney, S. H. and Lilley, T. H. (1988)**. Carbohydrate polyphenol complexation, p. 553. *In* R. W. Hemingway and J. J. Karchesy (ed.), *Chemistry and significance of condensed tannins*. Plenum Press, New York, N.Y.
- [24] **Hatice, Z. and Ayse, H. B. (2014)**. Antibacterial and Antioxidant Activity of Essential Oil Terpenes against Pathogenic and Spoilage-Forming Bacteria and Cell Structure-Activity Relationships Evaluated by SEM Microscopy. *Molecules Journal* 19(11), 17773-17798
- [25] **Hena, J. S. I., Adamu, A. K. L., Iortsuun, D. N. and Olonitola, O. S. (2010)**. Phytochemical Screening and Antimicrobial Effect of the Aqueous and Methanolic Extracts of Roots of *Balanites aegyptiaca* (Del.) On Some Bacteria Species. *Science World Journal* Vol 5 (No 2)
- [26] **Jeruto, P., Mutai, C., Catherine, L. and Ouma, G. (2011)**. Phytochemical constituents of some medicinal plants used by the Nandis of South Nandi district, Kenya. *Journal of Animal and Plant Sciences*; Vol. 9, Issue 3: 1201- 1210.
- [27] **Jeruto, P. Too, E. Mwamburia, L. A. and Amuka, O. (2015)**. An Inventory of Medicinal Plants used to Treat Gynaecological-Obstetric-Urino-Genital Disorders in South Nandi Sub County in Kenya. *Journal of Natural Sciences Research* ISSN 2224-3186 (Paper) Vol.5, No.18, 2015
- [28] **Jones, G. A., McAllister, T. A., Muir, A. D. and Cheng, K. J. (1994)**. Effects of sainfoin (*Onobrychis viciifolia* scop.) condensed tannins on growth and proteolysis by four strains of ruminal bacteria. *Applied Environmental Microbiology* 60:1374-1375. Kenya national bureau of statistics census (2009). Kipruto, A. Sinei, F. A., Okalebo, A., Hannington, N., Mugo, B. and Josephat, M. M. (2013). An Investigation of the Antimicrobial Activity of *Acmella caulirhiza*. *African Journal of Pharmacology and Therapeutics* Vol. 2 No. 4 Pages 130-133.
- [29] **Kiran, C. D. N. and Prasad, B. S. S. (2015)**. Preliminary pharmacognostic and phytochemical Studies on *Nerium oleander* Linn. (White cultivar). *Journal of Pharmacognosy and Phytochemistry*; 4(1):185-188
- [30] **Kokwaro, J. O. (2009)**. *Medicinal plants of East Africa*: East Africa Literature Bureaus Nairobi-Kenya. Page 11 production. *J Lab Clin Med*; 53:807.
- [31] **Krishna, K. V. V. S., Surendra, G., Anjana, M. and Nagini, K. S. K. (2012)**. Phytochemical screening and antimicrobial activity of *Callistemon citrinus* (L.) leaves extracts. *International Journal of Pharmaceutical and Technology Research*.; 4 (2):700-704.)
- [32] **Kumar, S. and Pandey, A. K. (2013)**. Chemistry and Biological Activities of Flavonoids: An Overview. *The Scientific World Journal* Volume 2013 Article ID 162750, 16 pages
- [33] **Marcy, J. B. Douglas, K. A. (2005)**. Drug discovery from medicinal plants *Life Sciences* 78; 431 – 441

- [34] Mishra, A. K. Mishra, A. Kehri, H. K. Sharma, B. and Pandey, A. K. (2009). "Inhibitory activity of Indian spice plant *Cinnamomum zeylanicum* extracts against *Alternaria solani* and *Curvularia lunata*, the pathogenic dematiaceous moulds," *Annals of Clinical Microbiology and Antimicrobials*, vol. 8, article 9
- [35] Mohd, I., Shakeel, A. and Manik, S. (2014). Antimicrobial activity of terpenoids from *Sphaeranthus indicus* L. *Asian Journal of Plant Science and Research*, 4(1):1-6
- [36] Moyo, B., Masika, P. J. and Muchenje, V. (2012). Antimicrobial activities of *Moringa oleifera* extracts. *African Journal of Biotechnology*; 12:34-42.
- [37] Neil, C. and Jerald, J. (2005). A novel alkylamide from the leaves of *Acmella caulirhiza* (Asteraceae), a traditional surface analgesic. *South African Journal of Botany*.
- [38] Njoroge, G. N. and Bussmann, R. W. (2006). Traditional management of ear, nose and throat. *J. Ethnobiol. and Ethnomed.* 2: 1-9.
- [39] Omojate, G. C., Enwa, F. O, Jewo, A. O. and Eze, C. O. (2014). Mechanisms of Antimicrobial Actions of Phytochemicals against Enteric Pathogens. *Journal of Pharmaceutical, Chemical and Biological Sciences* ISSN: 2348-7658; 2(2):77-85
- [40] Omwenga, E. O., Ogot, C., Paul, K. M. and Paul, O. O. (2012). Ethnobotanical Identification and Anti-microbial Evaluation of Some Anti-Diarrhoeal Plants Used by the Samburu Community, Kenya. *Malaysian Journal of Microbiology*, Vol 8(2) pp. 68-74
- [41] Pandey, A. K. Mishra, A. K. Mishra, A. Kumar, S. and Chandra, A. (2010). "Therapeutic potential of *C. zeylanicum* extracts: An antifungal and antioxidant perspective," *International Journal of Biological and Medical Research*, vol. 1, pp. 228-233.
- [42] Pikal, J. M., Takayuki, D. and Patel S. (2010). Determination of End Point of Primary Drying in Freeze-Drying Process Control. *AAPS PharmSciTech.* 11(1): 73-84.
- [43] Prajapati, N. D., Purohit, S. S., Sharma and Tarun, K. (2010). *A handbook of Medicinal plants*. A complete source book. Agrobios publishers India. Page 140,150. Prawat, U., Chairerk, O., Lenthas, R., Salae, A. W. and Tuntiwachwuttikul, P. (2013). Two new cycloartane-type triterpenoids and one new flavanone from the leaves of *Dasymaschalondasymaschalum* and their biological activity. *Phytochem Lett* 6: 286-290.
- [44] Sabri, F. Z., Belarbi, M., Sabri, S., Alsayadi, M. M. S. (2012). Phytochemical Screening and identification of some compounds from Mallow. *J. Nat. Prod. Plant Resource* 2 (4):512-516
- [45] Sameerah, A. Z., Adnan, J. M. A. and Ghosoon, F. A. (2013). Antibacterial activity of the glycosidic extract From *Citrus laurantiifolia* L. fruits, *Scholars Research Library* 5(6):73-78
- [46] Sangeetha, V. S., Michael, B. and Beena, L. (2014). Phytochemical Analysis of *Annona Reticulata* L. Leaf Extracts. *Int. Res J Pharm. App Sci.*, 4 (5):4-8
- [47] Tanaka, O., Tamura, Y., Masuda, H. and Mizutani K. (1996). Application of saponins in foods and cosmetics: Saponins of Mohave yucca and *Sapindus mukurossi*. Pages 1-11 in Saponins Used Food and Agriculture. G. R. Waller, and K. Yamasaki, eds. Plenum Press, NY. Telefo, P. B., Lienou, L. L., Yemele, M. D., Lemfack, M. C, Mouokeu C, Goka CS, Tagne, S. R. and Moundipa, F. P. (2011). Ethnopharmacological survey of plants used for the treatment of female infertility in Baham, Cameroon. *Journal of Ethnopharmacology* 136: 178-187.
- [48] Thornes, R. D., Lynch, G. and Sheehan, M. W. (1982). Relationship between structure and Anti-Coagulant activity of coumarin derivatives. *Lancet.* 2: 328-339.
- [49] Vichith, L., Hugo J. B. and Lars, B. (2011). Traditions and plant use during pregnancy, childbirth and postpartum recovery by the Kry ethnic group in Lao PDR. *J Ethnobiol*
- [50] Widelski, J., Popova, M., Graikou, K., Glowniak, K. and Chinou, I. (2009). Coumarins from *Angelica lucida* L. Antibacterial Activities. *Molecules* 14, 2729-2734
- [51] World Health Organization (2003a). Guidelines on good agricultural and collection practices for medicinal plants. Page 11
- [52] World Health Organization (2014). Traditional Medicine Strategy 2014-2023 Page 1, 2
- [53] Yogayata, S. P and Vijay, D. W. (2012). Herbal medicines and nutritional supplements used in the treatment of glaucoma. *Research Journal of Pharmaceutical, Biological and Chemical Sciences* 3 (1): 331
- [54] Zhang, L., Kong, Y., Dalei, W., Haitao, Z., Jian, W., Jing, C., Ding, J., Lihong, H., Jiang, H. and Shen, X. (2008). Three flavonoids targeting the β -hydroxyacyl-acyl carrier protein dehydratase from *Helicobacter pylori*: Crystal structure characterization with enzymatic inhibition assay *Protein Sci.* 17(11): 1971-1978.

AUTHORS

First Author – Jespher Nyaboke Onyango, BSc, Mount Kenya University and email address rebecca.nyaboke059@gmail.com.

Second Author – Jared Misonge Onyancha, MSc, Kenyatta University and email address onyancha.jared@mku.ac.ke

Third Author - Dr. Jackson Odhiambo Onyuka, PhD. Mount Kenya University and email address jonnyuka@mku.ac.ke

Fourth Author – Prof. John Memba Ochora, PhD. Jomo Kenyatta University of Agriculture and Technology, and email address jochora@jkuat.ac.ke

Fifty Author – Patrick Ogembo Getonto, BSc, Mount Kenya University and email address ppgetonto059@gmail.com

Sixty Author - Charles Wambugu Maina, BSc, Mount Kenya University and email address cwamaina2015@gmail.com

Correspondence Author – Jespher Nyaboke Onyango, jnyaboke@mku.ac.ke, rebecca.nyaboke059@gmail.com ; 0726734760

ACKNOWLEDGMENT

I am glad to almighty God for being with me through this far.

I am importantly grateful to my supervisors Prof. John Memba Ochora Department of Botany Jomo Kenyatta University of Agriculture and Technology, Dr. Jackson Odhiambo Onyuka from the school of Medicine, Department of Medical Laboratory Science, Mount Kenya University and Mr. Jared Misonge Onyancha from the School of Pharmacy, Department of Pharmacognosy, Mount Kenya University who strongly and patiently supervised this research project, giving me valuable guidance throughout this research thesis period. My appreciation goes to my immediate supervisor Mr. Masasi of Mount Kenya University health centre and

my colleagues at work for encouraging me to work hard. Lastly my appreciation goes to Mount Kenya University and Department of Medical Laboratory, school of Medicine for providing me with the opportunity to take my MSc studies. May God bless them.