

# A Field Level Studies on Selected Serum Biochemical Constituents in Cyclic, Non-Cyclic and Endometritic Crossbred Cows

Vijaya Bhaskar M. Reddy<sup>1\*</sup>, A. Karthik<sup>1</sup>, P. Sasikala<sup>2</sup>

<sup>1\*</sup> Vijaya Bhaskar M. Reddy<sup>1\*</sup>, Ph. D, Department of Livestock Production Management, C.V.Sc, SVVU, Tirupati, Andhra Pradesh, India

<sup>1</sup> A. Karthik: M. Sc, Department of Microbiology, C.V.Sc, Tirupati, Andhra Pradesh, India

<sup>2</sup> P. Sasikala, M. Sc, Department of Livestock Production Management, C.V.Sc, Tirupati, Andhra Pradesh, India

**Abstract-** Forty-five Jersey Sahiwal cows selected from nearby villages from tirupati. Jersey Sahiwal were divided into three equal groups i.e. cyclic, non cyclic and endometritic. Blood samples were collected from all the experimental animals, serum was separated and stored at -20°C until assayed for glucose (g/dl), total protein (g/dl), urea (mg/dl) and cholesterol (mg/dl). The results revealed significantly ( $P < 0.05$ ) higher values of glucose ( $58.08 \pm 2.59$ ) and cholesterol ( $290.72 \pm 15.95$ ) in endometritic cows as compared to cyclic ( $50.72 \pm 1.12$ ,  $199.12 \pm 9.38$ ) and non-cyclic cows ( $50.56 \pm 1.12$ ,  $202.96 \pm 14.84$ ). Total protein level differed significantly ( $P < 0.05$ ) among cows of all the three groups, being highest in endometritic ( $19.16 \pm 1.00$ ), followed by non cyclic ( $15.23 \pm 0.89$ ) and lowest in cyclic ( $9.19 \pm 0.45$ ) cows. However, serum level of urea did not differ in cyclic ( $30.88 \pm 2.42$ ), non cyclic ( $33.80 \pm 3.43$ ) and endometritic ( $37.12 \pm 3.45$ ) animals, although highest value was recorded in endometritic animals.

**Index Terms-** Blood, biochemical constituents, cows, cyclic, non cyclic, endometritic.

## I. INTRODUCTION

The breeding efficiency of dairy cows is lowered by a number of reproductive disorders like endometritis, an-oestrus and repeat breeding, affecting adversely the productive and reproductive performance of cows, and resulting in great economic losses to dairy farmers (Dutta et al., 1988). The basic causes of the reproductive problems in a herd are multiple and include managemental, nutritional and pathological factors. The findings of many authors (El-Azab et al., 1993; Balakrishnan and Balagopal, 1994; Qureshi, 1998) suggest that normal blood levels of various biochemical constituent are indispensable for normal function of various systems of body including reproductive system. Changes in various biochemical constituents have been blamed for reproductive failures. Thus, serum biochemical profile might be a potential aid in characterizing these problems. Blood glucose appears to be one of the key nutrients affecting cyclicity in farm animals and a minimum level of 60-40mg/ml is required to maintain the physiological processes of the body (Duke, 1970). According to Dowine and Gelman (1976), low blood glucose may be

associated with infertility. Nadiu and Rao (1982) and Dutta et al. (1988) reported significantly lower serum glucose level in an-oestrus than normally cycling animals. El-Azab et al. (1993) reported significantly higher serum protein in cyclic cows than the non cycling one's. However, Tegegne et al. (1993) found inconsistent trend. Qureshi (1998) reported higher blood urea level in an-oestrus animals than those resuming cyclicity within 45 days postpartum. Burle et al. (1995) reported lowest serum concentration of cholesterol in an-oestrus than in cycling cows. Greatly variable reports were available on the level of these biochemical constituents, hence the present study was planned to assess the levels of selected biochemical constituents in cyclic, non-cyclic and endometritic crossbred cows.

## II. MATERIALS AND METHODS

The study was conducted at field level. Fifteen animals each for three groups i.e., cyclic, non-cyclic and endometritic were selected. The animals with a corpus luteum (CL) on one of the ovaries were considered as cyclic animals, while cows having no functional CL were included in non cyclic group. The animals having thick uterus and containing pus in the oestral mucus were placed in endometritic group. Blood samples were collected by jugular veinipuncture; serum was separated and stored at -20°C until analyzed. Serum glucose, urea, cholesterol and total proteins were determined spectrophotometrically according to the methods described by Braham and Trinder, (1972), Merck (1974), Allain et al. (1974) and Doumas et al. (1981), respectively.

## III. STATICAL ANALYSIS

The data were analyzed statistically using one way analysis of variance technique (Steel and Torrie, 1980). Significant means were compared by Duncan's Multiple Range Test (Duncan, 1955).

## IV. RESULTS AND DISCUSSION

Serum biochemical constituents (mean  $\pm$  SE) in cyclic, non cyclic and endometritic crossbred cows are given in Table 1. Mean concentrations of serum glucose in cyclic, non-cyclic and

endometritic cows were  $50.72 \pm 1.12$ ,  $50.56 \pm 1.13$  and  $58.08 \pm 2.59$  g/dL, respectively. The results clearly indicate that serum glucose level of endometritic cows was significantly ( $P < 0.01$ ) higher than cyclic and non-cyclic animals. Majeed et al. (1990) recorded higher serum glucose level in endometritic animals than healthy ones. Increased blood glucose level in endometritic animals may result from either an imbalance between hepatic output and peripheral uptake of the sugar or disturbances in the endocrine regulatory mechanisms which influence these processes. Abnormal functioning of hormone producing organs may influence glucose levels (Coles, 1986). Increased activity of the anterior pituitary, adrenal cortex and thyroid may result in increased blood glucose level. Mechanism by which hyperpituitarism resulted is incompletely understood; however it may be related to increased production of adrenocorticotrophic hormones (Coles, 1986). Total protein level was significantly higher ( $P < 0.05$ ) in endometritic ( $19.16 \pm 1.00$  g/dL) cows as compared to cyclic ( $9.19 \pm 0.45$  g/dL) and non-cyclic ( $15.23 \pm 0.89$  g/dL) cows. The level in non cyclic cows was significantly ( $P < 0.05$ ) higher than that of cyclic animals. Majeed et al. (1990) and Lyubetsky (1997) reported higher values of total serum protein in endometritic buffaloes and cows compared to cyclic buffaloes and cows, respectively. However, Burle et al. (1995) reported significantly higher value of total serum protein in cyclic cows. The variation could be due to differences in breeds, environment and level of nutrition. In agreement to the findings of the present study,

higher level of total serum protein was associated with low fertility, as reported by Hewett (1974). Excessive intake of protein in the feed can reduce fertility and increase the number of service per conception. Fertility is impaired more by feeding excessive protein to older cows. The mechanism by which high level of protein adversely affects reproduction in dairy cows is unknown (Randel, 1990). No significant difference in concentration of urea in cyclic ( $30.88 \pm 2.42$  mg/dL), non-cyclic ( $33.80 \pm 3.45$  mg/dL) and endometritic ( $37.12 \pm 3.45$  mg/dL) cows was observed in the present study. The values observed in endometritic cows were apparently higher but not significantly different. Zaman et al. (1985) and Majeed et al. (1990) also observed non significant ( $P < 0.05$ ) difference in blood urea among cyclic, non cyclic and endometritic animals. Serum cholesterol level was found to be significantly ( $P < 0.05$ ) higher in endometritic ( $290.72 \pm 15.95$  mg/dL) as compared to cyclic ( $199.12 \pm 9.38$  mg/dL) and non-cyclic ( $202.96 \pm 14.84$  mg/dL) cows. However, the difference between cyclic and non cyclic animals was non significant. Similarly, Zaman et al. (1985) reported a non significant difference in levels of plasma cholesterol of cyclic and non-cyclic buffaloes. Majeed et al. (1990) found a non significant difference in serum cholesterol level between endometritic and healthy buffaloes.

Identify the constructs of a Journal – Essentially a journal consists of five major sections. The number of pages may vary depending upon the topic of research work but generally comprises up to 5 to 7 pages.

**Table1: Serum biochemical constituents (mean  $\pm$  SE) in cyclic, non-cyclic and endometritic crossbred cows**

Parameters	Cyclic cows	Non-cyclic cows
Glucose (g/dL)	$50.72 \pm 1.12^b$	$50.56 \pm 1.13^b$
Total protein (g/dL)	$9.19 \pm 0.45^c$	$15.23 \pm 0.89^b$
Urea (mg/dL)	$30.88 \pm 2.42^a$	$33.80 \pm 3.45^a$
Cholesterol (mg/dL)	$199.12 \pm 9.38^b$	$202.96 \pm 14.84^b$

Values with different superscripts in a row differ significantly ( $P < 0.05$ ).

REFERENCES

[1] Allain, C.C., L.S. Poon and C.S.G. Chan, 1974. Enzymatic determination of total serum cholesterol. Clin. Chem., 20: 470. 2<sup>nd</sup> Ed. Edited by W.R. Faulkner and S.M. Mietes. Washington, D.C.

[2] Balakrishnan, V. and R. Balagopal, 1994. Serumcalcium, phosphorus, magnesium, copper and zinc levels in regular breeding buffaloes. Indian Vet. J., 71: 23-25.

[3] Braham, D. and P. Trinder, 1972. Methods for determination of blood glucose level by spectrophotometer. Anstlyst, 97: 142.

[4] Burle, P.M., N.S. Mangle, M.D. Kothekhar and D.R. Kalorey, 1995. Blood biochemical profiles during various reproductive states of Sahiwal and Jersey X Sahiwal cattle. Livestock Adv., 20(7): 13-20.

[5] Coles, E.H., 1986. Veterinary Clinical Pathology, 4<sup>th</sup> Ed. W.B. Saunders Company, London.

[6] Doumas, B., D.D. Bayse, R.J. Carter, T. Peters and R. Schaffer, 1981. A candidate reference method for determination of total protein in serum. Clin. hem., 27: 1642. 8<sup>th</sup> Ed. Edited by W.R. Faulkner and S.M. Mietes. Washington, D.C.

[7] Dowine, J.G. and A.L. Gelman, 1976. The relationship between changes in body weight, plasma glucose and fertility in beef cows. Vet. Rec., 11: 210-212.

[8] Duke, H.H., 1970. Physiology of Domestic Animals. 8<sup>th</sup> Ed., Comstock Publishing Associates. Ithaca and London.

[9] Duncan, D.B., 1955. Multiple ranges and multiple F test. Biometrics, 11: 1-42.

[10] Dutta, J.C., R.N. Baruah, L. Dutta and S.C. Talukdar, 1988. Blood biochemical studies in anoestrous and normal cyclic cattle. Indian Vet. J., 65: 239-241.

- [11] El-Azab, M.A., A. Badr, El-Sadawy, G. Shawki and T.M. Borkat, 1993. Some biochemical changes in relation to postpartum ovarian activity in dairy cows. *Indian J. Anim. Sci.*, 63(12): 1244-1247.
- [12] Hewett, C., 1974. Studies on the causes and effect of variation in the blood profile of Swedish dairy cattle. *Acta Vet. Scand.*, 50: 152-156.
- [13] Lyubestsky, V.I., 1997. Plasma protein fractions in cows with post parturient edometritis. *Visnik- Agrarnoi-Nauli.*, 6: 28-33.
- [14] Majeed, M.A., J. Iqbal and M.N. Chaudhry, 1990. Blood chemistry of clinical merits in Nili-Ravi buffaloes of two age groups and at two stages of lactation. *Pakistan Vet. J.*, 10(2): 55-59.
- [15] Merck, E., 1974. *Clinical Laboratory Techniques*. 11<sup>th</sup> Ed., 379 Dermasted, Federal Republic of Germany. Nadiu, K.V. and A.R. Rao, 1982. A study on the etiology of an-oestrus in crossbred cows. *Indian Vet. J.*, 59(10):781-782.
- [16] Qureshi, M.S., 1998. Relationship of pre and post partum nutritional status with reproductive performance in Nili-Ravi buffaloes under the conventional farming system in NWFP, Pakistan.
- [17] Ph.D. Thesis, Univ. Agric. Faisalabad, Pakistan. Randel, R. D., 1990. Nutrition and post partum rebreeding in cattle. *J. Anim. Sci.*, 68: 853 - 862. Steel, R.G.D. and J.H. Torrie, 1980. *Principles and Procedures of Statistics. A Biometrical approach*. 2<sup>nd</sup> Ed. McGraw Hill Book Co. Inc. New York.
- [18] Tegegne, A., K.W. Entwistle and E.M. Mugerwa, 1993. Plasma progesterone and blood metabolite profiles in post partum small East African Zebu cows. *Trop. Anim. Hlth. Prod.*, 25: 101- 110.
- [19] Zaman, M.S., C.S. Ali and K.M. Ahmad, 1985. Comparative study of blood glucose, cholesterol, protein and urea contents in cyclic, non-cyclic and s`sub-oestous lactating buffaloes. *Pakistan Vet. J.*, 5 (2): 72-75.
- [20] Ijaz Ahmad, L.A. Lodhi, Z.I. Qureshi and M. Younis Pakistan 2004. Studies on blood glucose, total proteins, urea and cholesterol levels in cyclic, non-cyclic and endometritic crossbred cows *Vet. J.*, 24(2):92-94

#### AUTHORS

**First Author** – Vijaya Bhaskar M. Reddy, Ph. D, Department of Livestock Production Management, C.V.Sc, SVVU, Tirupati, Andhra Pradesh, India.

**Second Author** – A. Karthik: M. Sc, Department of Microbiology, C.V.Sc, Tirupati, Andhra Pradesh, India., Tel: +91-9704582512, E-mail: vijaymutlam@gmail.com

**Third Author** – P. Sasikala, M. Sc, Department of Livestock Production Management, C.V.Sc, Tirupati, Andhra Pradesh, India., Tel: +91-9492150844, E-mail: vbrsasikala@gmail.com

**Correspondence Author** – M. Vijaya Bhaskar Reddy, Tel: +91-9705497317, E-mail: vijaymutlam@gmail.com, vijaybhaskar24@gmail.com