

# New Antigenic Approach to Diagnose Visceral Leishmaniasis in India

Kumar Rajesh\*, Kumar Sanjay\*\*

\* Associate Professor, Faculty of Advance Studies in Life Sciences, Chhatrapati Shahu ji Maharaj University, Kanpur, 208024, Uttar Pradesh, India

\*\* PhD Scholar, Faculty of Advance Studies in Life Sciences, Chhatrapati Shahu ji Maharaj University, Kanpur, 208024, Uttar Pradesh, India

**Abstract-** Visceral Leishmaniasis (VL) is a vector borne anthrozoontic disease caused by a protozoan, *Leishmania donovani*, of *Trypanosomatidae* family. The most precise methods used for its diagnosis includes the analysis of spleen & liver smears (90%), Bone Marrow smear (80%), sternal or iliac crest puncture, and referred as gold standard for diagnosis. Various serological tests like Aldehyde test, antimony test, Leishmania skin test, Direct agglutination Test(DAT), Indirect fluorescent antibody test (IFAT), ELISA etc. are also used to diagnose VL, but in the case of immunocomprised patient, diagnosis of VL is difficult due to low titer of antibody. Several molecular diagnostic techniques are also available but they need from more sophisticated laboratory and expertise person. The current article is devoted to the detection of specific antigen which are useful in diagnosis of VL in VL patients as well as in immunocomprised patients.

**Index Terms-** Visceral Leishmaniasis, LD bodies, Antibody detection, Antigens

## I. INTRODUCTION

Visceral Leishmaniasis (VL) also known as Kala-azar (Hindi: kala means black, azar means sickness, also known as Assam fever, Dumdum fever, Sikari disease, Burdwan fever, Shahib's disease and tropical splenomegaly) was first described in 1824, in Jessore District of Bengal, what is now Bangladesh. Leishmaniasis, because of its importance, was selected by WHO out of six most precarious diseases under its special programme for research and training in tropical diseases. It ranks next to malaria among the human diseases caused by protozoan. The disease is endemic, covering almost the five continents, in 88 countries (16 developed and 72 developing or 66 countries are in the Old World while 22 countries located in the New World) with a total of 350 million people at risk and 12 million cases of infection. Out of 500, 000 cases of visceral Leishmaniasis, more than 90 percent are reported from India, Bangladesh, southern Sudan, Nepal and northeast Brazil At the turn of the 20<sup>th</sup> Century VL was observed in many parts of India. In India, kala-azar (Visceral leishmaniasis; VL) affects millions of people, and the state of Bihar accounts for nearly 90% cases, followed by West Bengal and Eastern Uttar Pradesh. In Uttar Pradesh recently few active cases have been seen from eastern districts e.g. Varanasi, Gorakhpur, Gonda, Faizabad etc. where there is under reporting of kala-azar and post kala-azar dermal leishmaniasis in women and children 0-9 years of age. Untreated cases of kala-azar are associated with up to 90% mortality. Although disease is to

widely prevalent, the accurate data on the incidence and prevalence of Leishmaniasis is lacking especially from this part of the country.

## II. MATERIALS AND METHOD

The standard strain of *Leishmania donovani* (Strain *HOM/IN/80/Dd8*: WHO reference strain originally isolated from human Kala-azar patients from Muzaffarpur, Bihar, India), is taken for the preparation of various antigens. NNN<sup>o</sup> Media is used to culture promastigotes. The polyacrylamide gel electrophoresis is carried out by Davis, B. J., 1964.

Pathological methods are mainly based on demonstrate of parasites in biopsies or aspirated from infected spleen, bone marrow and lymphnodes (Zijlstra et al., 1992). Their sensitivities are in the range of 90-95% (Manson Bahr 1987; Zijlstra et al., 1992). Parasitological diagnosis remains the gold standard in Leishmaniasis diagnosis, because of its high specificity.

Detection of LD bodies in smear of Spleen, bone marrow Painful, cumbersome and some times. Leishmanian test or Intradermal skin reaction or Montenegro reaction (Singh S, 2003) was also used. This is a test for Delayed type hypersensitivity. In this test an injection of 0.5 ml ( $5 \times 10^7$  phenol killed promastigotes) in volar aspect of the forearm of the patient leads to the formation of an induration measureable after 48 / 72 hours. Comparing the size of the induration with a control injection with phenol-saline in the other forearm it is possible to determine either an individual is infected or not. But this test not gives satisfactory result, because of large number of false positive and negative cases

**Serological Diagnosis:** Various serological tests are being used for the diagnosis of the disease but their demerits are that they cross react with other chronic diseases (Chatterjee, 1957; Evans, 1973). Other serological tests which have been used for the diagnosis are as:-

**Napier Test.** In Napier Test, About 1-2 ml of serum from a case of kala azar will be taken in a small glass tube and to it will add a drop or two of 40 % formalin. The aldehyde test procedure is simple. Five cubic millimeters venous blood is drawn from the patient and left for clotting. Serum is separated by removing supernatant after centrifugation. To one ml of serum, 2-3 drops of commercial formalin (40% formaldehyde) is added. Opaque jellification like white of a boiled egg indicates a positive reaction. Jellification time indicates intensify of infection as indicated in the table below.

**Antimony Test (Chopra et al):** This also depends upon a rise of serum gamma globulin. In Antimony test, 4% of urea stibamine solution in distilled water will add to whole serum from kala-azar patients in a small tube. A positive test will be indicated by the formation of a profuse flocculent precipitate. A negative result is indicated when the two fluids mix without precipitation. As a diagnostic aid antimony test is less reliable than the aldehyde test.

**Complement Fixation Test with W.K.K. Antigen:** This test depends upon the presence of certain immune bodies in blood sera of kala-azar patients, the antigen used in the test is prepared from either human tubercle bacilli (Witebsky, Kligenstein & Kuhn: WKK antigen) or from Kedrowsky's acid-fast bacillus. The test is essentially an antigen-antibody reaction in which the antibodies present in the serum are bound to an antigen.

**Direct Agglutination Test:** The Direct agglutination test (DAT) is a highly specific and sensitive test. It is inexpensive and simple to perform making it ideal for both field and laboratory use. The test is based on antigen-antibody reaction.

**Indirect Immuno fluorescent Assay Test (IFAT):** This is a commonly used technique for Kala azar. This is a specific and sensitive test requiring fewer amounts of antigens and detects the low titer of antibodies present in patient serum, which are demonstrated in the very early stage of infection and are undetectable six to ninth months after cure. If the antibodies persist in low titers, it is good indication of a probable relapse. The sensitivity and specificity of this test is 96 % and 98 % respectively.

**Enzyme linked Immunosorbant Assay:** ELISA has been used as a potential serodiagnostic tool for almost all infectious disease. Enzyme linked immunosorbent assay (ELISA) is also equally sensitive and specific but specificity depends upon the antigen used, requiring less amount of antigen. This is rapid inexpensive and simple to perform and can be used for diagnosis of Kala-azar for epidemiological work.

**Rapid Immunochromatographic assay (dipstick or strip test):** A promising ready-to-use immunochromatographic strip test based on rK39 antigen has been developed as a rapid test for use in difficult field conditions. The recombinant antigen is immobilized on a small rectangular piece of nitrocellulose membrane in a band form, and goat anti-protein A is attached to the membrane above the antigen band. After the finger is pricked, half a drop of blood is smeared at the tip of the strip, and the lower end of the strip is allowed to soak in 4 to 5 drops of phosphate-buffered saline, placed on a clean glass slide or tube. If the antibody is present, it will react with the conjugate (protein A colloidal gold) that is pre dried on the assay strip. The mixture moves along the strip by capillary action and reacts with rK39 antigen on the strip, yielding a pink band. In the strip of patients who are infected, two pinkish lines appear in the middle of the nitrocellulose membrane (the upper pinkish band serves as a procedural control). Several studies from the Indian subcontinent reported the test to be 100% sensitive).

**Western Blot:** A new method, the Western blot for detection of anti-leishmania antibodies in the sera is developed which allows specific serodiagnosis of Visceral Leishmaniasis in patients living in non-endemic areas (Salotra P, 1999). In non-endemic areas,

**Molecular Diagnosis:** A variety of nucleic acid detection methods have also been developed. PCR assay can detect presence of presence of parasite DNA or RNA in blood sample of any clinical sign or symptoms. Different DNA-based methods have been used for characterization of isolates of Leishmania at (Bernhard Ronacher 2002)

- (a) **Genus level:** by performing PCR using Leishmania common primers depending on direct identification of the parasite in clinical
- (b) **Species level:** different methods could be applied, such as PCR which depends on species specific primers (Ibrahim et al., 1994), analysis of restriction fragment length polymorphism (RFLP), where the substrate can be whole genomic DNA, kDNA or PCR amplification products, analysis of kinetoplast and nuclear DNA including Southern blot hybridisation with specific DNA probes (Jackson et al., 1984; Beverley et al., 1987; Barker, 1989; Van Eys et al., 1989&1991)
- (c) **Strain level:** These methods are capable to detect variation among isolates of Leishmania belonging to the same species such as sequencing.

**Antigen Detection:** Antigen detection is more specific than antibody-based immunodiagnostic tests (De Colmenares et al., 1995). This method is also useful in the diagnosis of disease in cases where there is deficient antibody production (as in AIDS patients) (Riera et al., 2004). Because of the conditions prevailing in endemic area, any sophisticated method cannot be employed on a wider scale. There is a need for a simple rapid and accurate test with good sensitivity and specificity, which can be used without any specific expertise. A promising ready to use Latex test (KAtex) based on detection of antigen in urine has been developed as a rapid test for use in difficult field conditions (Attar et al., 2001).

### III. RESULT & DISCUSSION

All above parasitological as well as serological techniques which are used to diagnose VL in India have some limitations and not is useful in field conditions. As we know that parasitological diagnosis is gold standard due to its specificity. Detection of LD bodies in smear of Spleen, bone marrow are invasive Painful, cumbersome and are not conveniently applicable under field conditions (Osman, 1998), but according to Neva & Sacks, 1990; Weiss, 1995, there is no possibility to distinguish between leishmania amastigotes belonging to the different species. *François Chappuis et al, 2007 demonstrated that*, The accuracy of microscopic examination is influenced by the ability of the laboratory technician and the quality of the reagents used. Leishmanian Skin Test not gives satisfactory result, because of large number of false positive and negative cases. LST requires culture facility to produce LST antigens. The different antigen preparation impact test sensitivity and the test

did not recognize present or past infections. In the case of gammaglobulin detection (serological diagnosis) in serum of the patients, the serum globulins also increase in a variety of infections and thus this test is considered to be rather non-specific. A positive reaction may also be seen in diseases like tuberculosis, cirrhosis of liver, malaria, etc. Further, in kala-azar, the test becomes positive only when infection is at least three months old and may remain so even after six months of cure. (Napier & Antimony) are non-specific, and show cross reaction with other pathological conditions, hence are unsuitable for diagnosis. Compliment fixation tests by W.K.K. antigen gives better results However, cross reactions are observed in cases of pulmonary tuberculosis, leprosy and Mycobacterium infections. Most serological tests are non-specific, permitting detection of antigen shared by related organisms. They do not distinguish antigens confined to a single species as group reactions mask specific ones. The major disadvantage of DAT is the relative long incubation time of 18 h and the need for serial dilutions of serum. Also, DAT has no prognostic value. DAT remains positive for a longtime after the disease is cured, and thus cannot be used as a test of cure or for diagnosis of relapses. Immunochromatographic dip stick test rK 39 has a drawback that an individual with a positive rK39 strip test result may suffer from an illness (malaria, typhoid fever, or tuberculosis) with clinical features similar to those of VL yet be misdiagnosed as suffering from VL. Using western blotting, one can find even minor antigenic differences among various organisms and thus detect cross reactive antigens. This process is time consuming, technically cumbersome, and expensive. Two urinary antigens of 72-75 and 123 kDa have been reported to be very useful in diagnosis and prognosis of Kala-azar with sensitivity of 96% and specificity of 100%.

*Cruz et al. 2006* demonstrated that the detection of antigens in serum is complicated by the presence of high levels of antibodies, circulating immune complex, serum amyloid, rheumatoid factors, and autoantibodies,

Total 7 bands were obtained with Dd8 strain which were 35Kda, 27Kda, 19 Kda, 15Kda, 10Kda, 9.5Kda and 8 Kda. Of these 7 bands, 8.0 Kda protein fractions were shown to react with all kala-azar sera.

However, sera from cured cases of kala-azar and other disease sera did not show any reaction with this 8.0 Kda protein fraction. This 8.0Kda protein fraction was observed in all 14 isolates tested including DD8. Antigenic fractions of 8 of these isolates were electro-transferred on to the surface of NCP and subsequently immunoblotted.

#### IV. CONCLUSION

Many of the diagnostic methods / procedures are reported with due course of time but none of them was found completely reliable for the diagnosis of the VL. Review of several diagnostic approaches revealed that antigen detection is most sensitive and better technique to diagnose VL patients including immunocomprised ones. This review article will provide definitely the insight to the readers about the various methods and be helpful for the development of new specific antigens for the diagnosis of VL

#### REFERENCES

- [1] Adriano Gomes Silva et al. (2008). Serological reactivity of different antigenic preparations of *Leishmania amazonensis* and *Leishmania braziliensis* complex. *Revista da sociedade Brasileira de Medicina tropical* 41 (2): 135-141.
- [2] B. Sarkari et al. (2005). A capture ELISA for the Diagnosis of Visceral Leishmaniasis using a Monoclonal Antibody against a Leishmanial Urinary Antigen. *Iran. Biomed. J.*, 9(3), 117-122,
- [3] B. Sherchand Jeevan et al. (2003). KATex Test for the Detection of Urinary Antigens in Visceral Leishmaniasis Patients of Nepal. *J. Nepal Med. Assoc.*, 42, 191-195,
- [4] Boelaert M et al. (1999). Operational validation of the direct agglutination test for diagnosis of visceral leishmaniasis. *American J Trop Med Hyg.* 60(1):129-134.
- [5] C. Riera et al. (2004). Evaluation of a latex agglutination test (KATex) for detection of *Leishmania* coinfection: value in diagnosis and post-treatment follow-up. *Eur. J. Clin. Microbiol. Inf. Dis.*, 23(12), 899-904.
- [6] C. Vilaplana et al. (2004) Noninvasive Method for Diagnosis of Visceral Leishmaniasis by a Latex Agglutination Test for Detection of Antigens in Urine Samples. *J. Clin. Microbio.*, 42, 1853-1854
- [7] C.A.P. Tavares et al. (2003) Molecular diagnosis of leishmaniasis. *Expert Rev. Mol. Diagn.*, 3(5), 657-667
- [8] DeKorte et al. (1990). Introduction of an improved direct agglutination test for the detection of *Leishmania infantum* infection in southern France. *Parasitol Res.* 6: 526-30.
- [9] Edrissian GH and Darabian P. (1979). A comparison of enzyme linked immunosorbent assay and indirect fluorescent antibody test in the serodiagnosis of cutaneous and visceral leishmaniasis in Iran. *Trans R Soc Trop Med Hyg* 73:289-292.
- [10] F. Chappuis, et al. (2006) Field validity, reproducibility and feasibility of diagnostic tests for visceral leishmaniasis in rural Nepal. *Trop. Med. Int. Health*, 11(1), 31-40.
- [11] Harith AEL et al. (1988). Improvement of a DAT for field studies of visceral leishmaniasis. *J Clin Microbiol* 26:1321-1325.
- [12] Héctor Dardo Romero et al. (2009). Comparative study of serologic test for the diagnosis of asymptomatic VL in an endemic area. *Am J Trop. Med. Hug.* 81 (11). 27-33
- [13] Hommel M et al. (1978). The micro-ELISA technique in the serodiagnosis of visceral leishmaniasis. *Ann Trop Med Parsitol* 72:213-218.
- [14] Israel cruz et al. (2006). Comparison of New diagnostic tools for management of pediatric Mediterranean visceral Leishmaniasis. *Journal of Clinical Microbiology*, 44, 2343-2347.
- [15] K. Agrawal (2006). Field Adaptable Tests for Kala-Azar. *J. Armed Forces India*, 62, 178-179,
- [16] M. Hommel et al. (2001). Katex for the diagnosis of human visceral leishmaniasis. *Med. Trop. (Mars)*, 61(6), 503-505
- [17] Marleen Boelaert et al. (2007). Evaluation of rapid diagnostic test: visceral Leishmaniasis. *Nature reviews Microbiology*, 530-539
- [18] ML Das et al. (2007). Use of rk39 for diagnosis of post kala-azar Dermal leishmaniasis in Nepal southeast asian j trop med public health Vol 38 No. 4 919-625.
- [19] P. Salotra and R. Singh (2005). Rapid & reliable diagnostic tests for visceral leishmaniasis. *Indian J. Med. Res.*, 122, 464-467,
- [20] Raj et al. (1999). Serodiagnosis of leishmaniasis with recombinant ORFF antigen. *Am J Trop Med Hyg* 61 : 482-7.
- [21] RK Goyal and TM Mohapatra (2004). Superiority of DAT over ELISA as a diagnostic and Seroepidemiological tool for the diagnosis of Indian Kala-azar *Indian Journal of Medical Microbiology*, 22 (1):57-60
- [22] S Rijal et al. (2004). Evaluation of a urinary antigen-based latex agglutination test in the diagnosis of kala-azar in eastern Nepal. *Trop. Med. & Int. Health*, 9, 724-729
- [23] S. Singh et al. (2005). Applications of molecular methods for *Leishmania* control *Exp. Rev. Mol. Diagn.*, 5(2), 251-265
- [24] S. Sunda and M. Rai (2002). Laboratory Diagnosis of visceral Leishmaniasis *Clin. Diagn. Lab. Immunol.*, 951-958,

- [25] S.H. El-Safi et al. (2003). Field evaluation of latex agglutination test for detecting urinary antigens in visceral leishmaniasis in Sudan. *Eastern Mediterr. Health J.*, 9(4), 844-855,
- [26] S.Sundar, et al. (2005). Detection of Leishmanial Antigen in the Urine of Patients with Visceral Leishmaniasis by a Latex Agglutination test. *Am. J. Trop. Med. Hyg.*, 73(2), 269-271.
- [27] *Schoone et al.* (2001). A fast agglutination test (FAST) for the detection of anti-Leishmania antibodies. *Trans, Royal Soc, Trop Med, Hyg.* Jul-Aug; 95(4):400-1.
- [28] Selcuk Kilic et al. (2008). Evaluation of serological tests for the Diagnosis of VL. *Turk J Med. Sci.* 38(1) 13:19.
- [29] Singla N et al. (1993). Evaluation of the direct Kala-azar in India. *Trans R Soc Trop Med Hyg.* 87:276-278.
- [30] Sinha R and Sehgal S. (1994) Comparative evaluation of serological test in Indian Kala-azar. *J Trop Med Hyg* 97:333-340.
- [31] Srivastava L and Singh VK.(1988). Diagnosis of Indian Kalaazar by dot enzyme – linked immunosorbent assay (dot-ELISA). *Ann Trop Med Parasitol* 82:331- 334.
- [32] Srivastava L. (1989). Comparative evaluation of serological test in diagnosis of Kala-azar. *Indian J Med Res.* 89:265-270.
- [33] Z.A. Attar et al. (2001). Latex agglutination test for the detection of urinary antigens in visceral Leishmaniasis. *Acta Tropica* 78, 11-16.

#### AUTHORS

**First Author** – Kumar Rajesh, Associate Professor, Faculty of Advance Studies in Life Sciences, Chhatrapati Shahu ji Maharaj University, Kanpur, 208024, Uttar Pradesh, India

**Second Author** – Kumar Sanjay, PhD Scholar, Faculty of Advance Studies in Life Sciences, Chhatrapati Shahu ji Maharaj University, Kanpur, 208024, Uttar Pradesh, India

**Correspondence Author** –Sanjay Kumar C/O Dr. Rajesh Kumar, Faculty of Advance Studies in Life Sciences, Email: [Skumarbt88@gmail.com](mailto:Skumarbt88@gmail.com), Phone: +91-945-385-2292