

Value Chain and Evaluation of Milk Quality in Smallholder Dairy Farms Around Addis Ababa, Ethiopia.

Melesse Etifu Meranga*¹ Mustefa Abu Kufa²

¹. Wolkite University, Department of Animal Production and Technology

². Federal Ministry of Agriculture, Addis Ababa

DOI: 10.29322/IJSRP.9.01.2019.p8510

<http://dx.doi.org/10.29322/IJSRP.9.01.2019.p8510>

Abstract- Evaluation of milk value chain and quality of milk were assessed in selected districts of surrounding Addis Ababa from September 2016 to April 2017. A total of 180 randomly selected market-oriented smallholder dairy farmers were involved in a cross-sectional study that was carried out by way of questionnaire survey, rapid market appraisal, farm inspection and group discussion. The overall mean family size of respondents in this study was 5.63 ± 1.926 persons with average livestock holding per household of 23.93 ± 11.755 animals. Cattle were the predominant species representing 84.3% of the total TLU. The average number of lactating cows owned by the respondent farmers was 1.76 ± 0.920 local and 2.79 ± 3.445 cross bred animals. Average daily milk yield of crossbred and local milking cows were 9.11 ± 2.902 and 1.889 ± 0.6707 liters respectively. Overall mean lactation lengths of crossbred and local milking cows were 9.7 ± 0.46 and 6.26 ± 0.6624 months. Sixty milk samples were collected and the analysis of microbiological and physicochemical were carried out. The overall mean chemical compositions of milk for fat (%), protein (%) and solids not fat (%) contents were 3.5693 ± 0.10892, 2.9646 ± 0.04621 and 6.9632 ± 0.12175 in bulk Tank milk samples. The overall mean microbiological count of log (TBC cfu/ml), log (CC cfu/ml) and log (SCC/ml) of raw milk was 8.2285 ± 0.10041, 3.3363 ± 0.10010 and 5.1622 ± 0.07382, respectively. The proportion of raw milk used for household consumption was relatively small (5%). The major part (86%) of milk produced by smallholders is destined to market. The main outlets for raw milk identified were cooperatives (55.6%), processors (20.0%), vendor (20.0%), directly to consumer (2.8%) and hotels/restaurants (1.7%). Price variations (cited by 87% of the respondents), lack of fair market (72.2%), lack of demand during fasting (49.4%), lack of preserving facilities, and absence of quality based payment and no/less say in deciding milk price by producers were the major problems of raw milk marketing.

Index Terms- Coliform, Milk, Physicochemical, Total bacteria, Value-chain.

I. INTRODUCTION

Ethiopia has a huge potential to be one of the key countries in dairy production for various reasons [1]. These include a large

population of milk cows in the country estimated at 9.9 million [2] a conducive and relatively disease free agro-ecology, particularly the mixed crop–livestock systems in the highlands that can support crossbred and pure dairy breeds of cows [3], a huge potential for production of high quality feeds under rain fed and irrigated conditions, existence of a relatively large human population with a long tradition of consumption of milk and milk products and hence a potentially large domestic market [4].

A number of fundamental constraints underlie these outcomes, including traditional technologies, limited supply of inputs (feed, breeding stock, artificial insemination and water), poor or non-existent extension service, high disease prevalence, poor marketing infrastructure, lack of marketing support services and market information, limited credit services, absence of effective producers' organizations at the grass roots levels, and natural resources degradation [5]. In addition, policy decision on milk and milk product marketing are taken in the absence of vital information on how they affect dairy producers, traders, exporters, and consumers. Similarly, current knowledge on dairy product market structure, performance and prices is poor for designing policies and institutions to overcome the perceived problems in the marketing system [6].

Traditional farmers sell their raw milk informally due to absence of organized marketing network that has made the produced milk unable to reach the consumer. Further losses incurred are quality losses by storing in unclean storage utensil, which is prone to high microbial contamination. Losses in spillage and contamination occur where handling during and after milking are traditional and care is not satisfactory. Additionally the trade in the sub-sector is constrained by various structural, production, information exchange, and promotional problems, as well as financial constraints.

Therefore, this study is conducted to evaluate value chain and quality of milk around Addis Ababa.

II. MATERIALS AND METHODS

This study is conducted in purposively selected districts around capital city of Addis Ababa namely Welmera and Sululta which are known for smallholder dairies are practiced for distribution of milk and milk products to the nearby city Addis Ababa. Welmera district is located 28 km west of Addis Ababa at

09°02 North latitude and 38°34 East longitudes with its altitude ranging from 2060-3380 m.a.s.l. whereas Sululta district lies between 39°30' N Latitude and 38° 30' and 39° 00 E longitude. It is located 40 km north west of Addis Ababa.

Study population

Smallholder farmers in Sululta and Welmera districts owning crossbred and indigenous cattle for milk production constituted the study population.

Study design

A cross-sectional study by way of questionnaire survey, rapid market appraisal, farm inspection, group discussion, interviewing key respondents and laboratory analysis of raw milk samples was carried out from September 2016 to April 2017. Marketing actors and smallholder dairy farmers in the selected study area were study participants.

Sample size determinations

The sample size was determined by using mathematical model of [7]. The sample size, N, can then be expressed as largest integer less than or equal to $0.25/SE^2$.

$$N=0.25/SE^2$$

Where, confidence level of 95% and confidence interval of 5%, were considered.

Based on the above formula the computed sample size was 180.

Sampling procedure

To select a representative sample, the potential of the two districts were identified. Sululta district has 23 peasant associations (PAs), of which eight have potential in dairy production. From the list of these eight PAs, three were selected randomly. These included Moye-Gajo, Chanco-Buba, and Warrarso-Malima PAs. Then ninety households owning dairy cattle were selected randomly from three PA's (thirty from each PA). Welmera District has also 24 PAs, of which six have potential in dairy production. From the list of these six, three of them namely Geggelikuyu, Bekeka na kore-oddo and Gebarobi PAs were selected. Then ninety households owning dairy cattle were selected randomly from the three PA's.

Data collection

Questionnaire survey

The questionnaire that was structured and closed type for its major part was pre-tested before its full administration. The questionnaire was focusing on demographic characteristics of the study participants, husbandry practices, milk production, processing, and marketing and utilization situations. Furthermore, marketing constraints of raw milk was investigated.

Rapid Market Appraisal (RMA)

Rapid Market Appraisal (RMA) using checklists and observation was implemented to understand how a product or commodity flows to reach the end users.

Farm inspection

Farms were inspected once at the same time with the questionnaire survey. Activities observed during the farm visit

encompassed kinds of utensils used, milking practices, milk handling and storage conditions.

Group discussions

Group discussions at three different PA'S of Sululta and three PA'S of Welmera were undertaken, in order to understand the overall community situations and get insight about milk marketing, milk handling, limitations and strength milk marketing. Groups were composed of 10 to 12 members constituted by different age and social groups. Discussion participants were identified in consultation with the wereda development agents. A sample checklist, which served as a guide and consisting of the main points for the group discussion was prepared.

Interviewing key respondents

Chairmen of PA'S, representatives of the sub PA'S and extension workers were interviewed. The agricultural office workers at PA'S levels were also participants in the process.

Collection of raw milk samples

Raw milk samples were collected at farm and milk collections centers by following strict aseptic procedures. Physicochemical test of raw milk was performed and the presence of bacteriological agents was assessed; standard plate count, coliform and somatic cell count tests were done. Before sampling, milk was thoroughly mixed after which 25 ml of milk was transferred into sterile sampling bottles. The milk sample bottles were capped, labeled with a permanent marker and stored in an ice packed cool box and transported to the Ethiopian Meat and Dairy Technology Institute, Debre-zeit where the different analysis were conducted.

Bacteriological quality tests

Tests employed to determine the quality of milk were Standard plate count, Coliform count, and Somatic cell count. Detailed description of the steps followed in each of the methodologies is presented in the following sections.

Standard plate count (SPC)

The standard plate count of raw milk samples was performed by putting one ml of milk sample into a sterile test tube having 9 ml peptone water. After mixing, the sample was serially diluted up to 1: 10⁻⁷ and duplicate samples of 1 ml of diluted milk samples were streaked on 15-20 ml standard plate count agar media and then incubated for 48 hours at 37°C to encourage bacterial growth. Finally, colony counts were made using colony counter. Single bacteria species or clusters grow to become visible colonies that were then counted. All plate counts were expressed as the number of colony forming units (cfu) per milliliter. Results from plates, which contained 10 to 300 colonies per plate were recorded. If plates from two consultative decimal dilutions yield colony counts of 10 to 300, the counts for each dilution were computed by the following formula [8]

$$N = \frac{\Sigma \text{ colonies}}{[(1 \times n_1) + (0.1 \times n_2)] \times d}$$

Where: N = number of colonies per milliliter of milk,

$\sum C$ = sum of colonies on plates counted,
 n_1 = number of plates on lower dilution counted,
 n_2 = number of plates in next higher dilution counted and
 d = dilution from which the first counts are obtained.

Coliform count (CC)

One ml of milk sample was added into sterile test tube having 9 ml peptone water. After mixing, the sample was serially diluted up to 1: 10⁻⁴ and duplicate samples (1 ml) were pour plated using 15-20 ml Violet Red Bile Agar solution (VRBA). After thoroughly mixing, the plated sample was allowed to solidify and laying over by Violet Red bile Agar solution (VRBA) then incubated at 37°C for 24 hours. Finally, colony counts were made using colony counter. Typical dark red colonies were considered as coliform colonies.

Somatic cell count (SCC)

For counting somatic cells, the microscopic method was used. Milk film preparation, staining and counting were done according to the standards set by International Dairy Federation [9]. To obtain a uniform distribution of cells, milk samples were mixed by moving upside down gently 25 times and letting it to stand for 2 minutes to permit air bubbles and foam disappear. Microscopic slides were degreased with alcohol before milk film preparation. A 0.01ml of milk was taken with a 50µl micropipette calibrated at 10 and spread evenly over one cm² area on a microscopic slide and allowed to dry at room temperature on a leveled table. One cm² area was delineated by a template prepared from a cap board. Dried films were fixed with ethanol for 15 minutes. Stained with toluidine blue for 5 minutes and washed with tap water gently and allowed to dry in a dust free area. Stained slides were stored in slide box until counted. Using oil immersion objective those cell nuclei clearly recognizable and those at the periphery with more than 50% of the cell body in view were counted. Twenty fields were counted from given sampled milk. The number of cells per ml of milk was calculated by multiplying the average number of cells per field with Magnifications filed (Laboratory manual).

$$\text{Somatic cell per ml of milk} = \frac{\sum SCOperfield * 10,000}{0.0346 * 20}$$

Where $\sum SCOperfield$ = the summations somatic cell counted per each field

0.0346 = oil immersion calibrated
 20 = Total number of field counted

Physicochemical test

The chemical compositions of milk (fat, protein, and solid not fat) and physical characteristics (density and freezing point), of the milk samples were determined by Ekomilk analyzer (Bulgaria), according to manufacturer’s instructions. Milk samples were mixed gently 4-5 times to avoid any air enclosure in the milk. Then 25 ml samples were taken in the sample-tube and put in the sample- holder one at a time with the analyzer in the recess position. Then when the starting button activated, the analyzer sucks the milk, makes the measurements, and returns the milk in the sample-tube and the digital indicator (IED display) shows the specified results.

Data analysis

The data collected from the study area were entered into Micro-soft-Excel spreadsheet for managing the data and analyzed using SPSS version 20. Descriptive statistics like means, standard deviation and frequency distribution were used to describe the farming system characteristics in the study area. One-way ANOVA statistical analysis was used for comparison of the performance variation. The correlation statistical analysis was used to study the interaction between the farming system characteristics and the interaction between physicochemical and microbiology of raw milk sample.

III. RESULTS

The overall mean family size for all respondents was 5.63 ± 1.926 persons. The family size ranged from 2 to 12 people. Fifty two percent of the family members were male and the rest (48%) were female. The overall average livestock holding per household was 23.93± 11.755. The average family and herd size of the two districts namely Sululta and Wolmera pointed out by the respondents (Table 4).

Table 1. Average family size and herd in smallholder dairy farms in the study district.

Variable s	Sululta(N=90)	Wolmera(N=90)	Overall(N=180)
	Mean ±S. D	Mean ±S. D	Mean ±S. D
Family size	5.49±1.819	5.77±2.028	5.63±1.926
Male	2.88±1.211	3.04±1.469	2.96±1.346
Female	2.68±1.198	2.76±1.248	2.72±1.220
Livestoc k	25.22±12.382	22.63±11.009	23.93±11.755
Cattle	14.69±11.619	11.37±3.905	13.03±8.802
Lactating cows	4.57±4.316	3.38±1.427	3.97±3.260
Local cows	2.18±0.384	2.08±.278	2.13±0.336
Cross bred cow	4.28±0.450	4.32±.470	4.30±0.459
Sheep	5.92±4.238	6.66±4.490	6.26±4.356
Equines	1.80±0.924	2.16±1.256	1.97±1.105

S.D=standard deviations N=number of respondents

Cattle composition

Table 2 shows the size and composition of cattle owned by the smallholders in the study areas. All the surveyed smallholders owned on average 13.03 ± 8.802 (12.29 TLU) cattle. The average number of Lactating cows owned by the respondent farmers was 1.76±0.920 local or 1.76 TLU and 2.79±3.445 crossbred animals or 4.185 TLU. Cattle were the predominant species representing 84.3% of the total TLU. The smallholders prefer to have crossbred cows because of their greater milk production, even though they require high management and susceptible to disease than local breeds.

Table 2. Cattle herd size and composition in TLU in smallholder farms.

Variables	Sululta(N=90)		Wolmera(N=90)		Overall(N=180)	
	Mean ±S. D	TL U	Mean ±S. D	TL U	Mean ±S. D	TL U
Cattle	14.69±1.619	13.006	11.37±3.903	11.391	13.03±8.802	12.29
L.M. cows	1.68±0.837	1.68	1.83±0.993	1.83	1.76±0.920	1.76
C.M. cows	3.30±4.511	4.95	2.21±1.252	3.315	2.79±3.445	4.185
Calves	3.23±3.083	0.646	2.76±1.126	0.552	3.01±2.357	0.602
Heifers	2.78±3.131	1.668	2.16±1.094	1.296	2.47±2.359	1.482
Bulls	1.56±0.940	1.872	1.39±0.549	1.668	1.48±0.777	1.776
Oxen	2.19±0.518	2.19	2.73±1.166	2.73	2.48±0.956	2.48

S.D =standard deviation TLU= tropical livestock units N= number of respondents 1TLU=250kg of live weight of livestock, L.M. cows=local milking cows C.M. cows= crossbred milking cows

Milking and milk handling practices

Ninety four percent of the respondents of the study area were using plastic pail for milking and milk handling. Nearly 6% were using Stainless steel pail. Difficulties of using these utensils were difficult for cleaning (1.1%), accessibility in local markets (5%) and no problem of using these utensils (93.9%) were indicated by the respondents of the study area. Through group discussions with the participant of the study areas it was pointed out that all the respondents practice washing the utensils used for milking and milk handling. Commonly they were washing the milking utensils with warm water by using soap and finally allow drying till milking. In the study area cows are hand milked and calves are allowed to suckle their dams prior to as well as after milking. About 100% of the respondents in Sululta and Wolemera area pointed out that they milk their cows two times a day at morning and evening. They milked their cows at barn, where the animals are sheltered. As illustrates on Table 3, all respondents were washing their hands and vessels before milking. Seventy two percent of respondents were also washing udder before milking. Nearly 19% of the smallholders were using individual towels for cleaning udder of milking cows in 52.2% of the cases collective towels were used while in the rest (28.9%) no towel use was practiced.

Table 3. Observed milking practices in the study areas (N=180)

Variable	Frequency Yes		Frequency NO	
	Frequency	Percent	Frequency	Percent
Wash milkier hands and vessels	180	100.0		
Wash udder before milking	129	71.7	51	28.3
Wash udder before and after milking			180	100.0
Use of individual towels	34	18.9	145	80.6
Use of collective towels	94	52.2	86	47.8
No towel	52	28.9	128	71.1

Milk production and use aspects

Mean of Lactation length of crossbred and local milking cows were 9.72 ± 0.45 and 6.353 ± 0.7681 in Sululta, 9.68 ± 0.47 and 6.167 ± 0.5567 in Wolmera district respectively and overall mean of lactation length of crossbred and local milking cows were 9.7 ± 0.46 and 6.26 ± 0.6624 months respectively. Average daily milk yield of cross bred and local cows in Sululta were 9.56 ± 3.010 and 1.809 ± 0.4574 Liter/day respectively. Moreover, crossbred and local cows in Wolmera areas were 8.60 ± 2.703 and 1.96 ± 0.8193 liters/day respectively. Overall mean summery of daily milk yield at the study areas of crossbred milking cows (9.11 ± 2.902) and local milking cows (1.889 ± 0.6707) liters as shows on table 4.

Table 4. Average lactation length and daily milk yield of local and cross bred milking cow of small holder farmers.

Variables	Sululta(N=90)	Wolmera(N=90)	Overall(N=180)
	Mean ± S.D	Mean ± S.D	Mean ± S. D
Lactation length of local cattle in month	6.353± 0.7681	6.167 ± 0.5567	6.26 ± 0.6624
Lactation length of cross bred cows in month	9.72 ± 0.45	9.68 ± 0.47	9.7 ± 0.46
Average daily milk yield of local cows(/liter/day)	1.809± 0.4574	1.96± 0.8193	1.889 ± 0.6707
Average daily milk yield of	9.56 ± 3.010	8.60 ± 2.703	9.11 ± 2.902

cross bred cows(/lit/day)

relatively small. As figure 1 illustrates, the major part of milk produced by smallholders is destined to market. Smallholders also process milk to butter and cheese. Milk was soured for 2-3 days before processing it in to butter and cheese. The one way of ANOVA analysis showed significance difference at (P<0.01) and (P<0.05) among the District from which the milk sample for milk produced and milk sold per day/liter.

S.D= standard deviations, N=number of respondents

Overall mean of milk producing, Processing, consuming and selling per day per household was 26.88±4.76, 1.23±1.603, 1.29±1.176 and 23.32±5.22 liters respectively (Table 5). The proportion of raw milk used for household consumption was

Table 5. Milk production and partition in to different use categories at smallholder farm level.

Category/area of study	Mean ± S. D	95% CI	Df	F	P
Milk produced at Sululta farm/lit/day	28.01±3.135	27.35-28.67	1	10.762	0.001**
	Wolmera	25.74±5.756	178		
	Overall	26.88±4.76	179		
Milk processed/lit/day	Sululta	1.29±1.493	1	0.215	0.643
	Wolmera	1.18±1.713	178		
	Overall	1.23±1.603	179		
Milk consumed at home/lit/day	Sululta	1.43±1.272	1	2.534	0.113
	Wolmera	1.16±1.059	178		
	Overall	1.29±1.176	179		
Milk sold/lit/day	Sululta	24.11±4.67	1	4.245	0.041*
	Wolmera	22.52±5.631	178		
	Overall	23.32±5.22	179		

**P-value is significant at 0.01 levels S.D= standard deviation and CI = confidence interval df= degree of freedom F=F ratios P= P value

Milk marketing

Table 6 shows distance between production and market place. Nearly 54.9% of the households were nearby to the market center for their raw milk marketing while about 4% of the households travel more than 10 km.

Table 6. Distance of market center for milk in smallholder dairy farmer of study areas (N=175 households).

Distance of marketing place	Frequency	Percent
Less than 1 km (nearby)	96	54.9
Between 1 – 5 km (proximity)	70	39.9
Between 5 – 10 km (intermediate)	2	1.1
More than 10 km(far)	4	2.3

Milk sales outlet

The main outlets for raw milk identified as shows in (Table7) were Cooperatives, Processors, Vendor, Directly to Consumer and Hotels/restaurants 55.6%, 20.0%, 20.0%, 2.8% and 1.7% respectively.

Table 7 .Marketing channel of smallholder farmer of the study area.

Milk out let	Frequency	Percent
Cooperatives	100	55.6
Hotels/restaurants	3	1.7
Vendor	36	20.0
Processors	36	20.0
Directly to Consumer,	5	2.8

Raw Milk marketing constraints

Table 8 illustrates raw milk marketing constraints at specific study area. The respondent farmers indicated that, price variations (87.2%), lack of fair market (72.2%) and lack of demand (49.4%) during fastening were the major problem of raw milk marketing in descending order of importance.

Table 8. Descriptions of marketing problems of small holder at the study area

Constraints	Frequency Yes	Percent	Frequency No	Percent
price variations	157	87.2	23	12.8

Lack of fair market	130	72.2	50	27.8
Lack of demand during fastening	89	49.4	91	50.6

As shows on table 9 milk price decided by producer, processor and collector were 6.1%, 25%, and 68.9% respectively as ascending order. Additionally through group discussion almost the entire group member pointed out they have less /no power to decided milk price at the study area.

Table 9 Power of decisions in milk price

Decisions of milk price	Frequency	Percent
Producer	11	6.1
Processor	45	25
Collector	124	68.9

Quality based payment was also another raw milk marketing constraints of the study area. They indicated quality based payment was enhanced quality of milk supplied to processors at the same time as encouraging them to produce more and quality milk. Through group discussions of respondents in the sturdy areas pointed out they possessed less preserving facilities for surplus milk produced and demand especially during fasting were great influence on raw milk marketing. Additionally, they showed that less adopted technologies for enhancing shelf life of raw milk in the study areas.

Factors influencing milk production, consumption and marketing

Milk production was positively and significantly correlated with experience of raising cattle for milk productions, raw milk sold ($P < 0.01$) and significantly correlated with distance of milk marketing ($P < 0.05$). Milk sold was positively and significantly correlated with experience of raising cattle, milk productions and distance of milk sold ($P < 0.01$). Milk consumption was negatively and significantly correlated with cattle herd size ($P < 0.01$) (Table 10).

Table 10. Correlations among different characteristics of small holder dairy farmers

Variable	Family size	ERCMP	Cattle	MPF/day	MS/day	MC/day	DMP
Family size	1						
ERCMP	.149*	1					
Cattle	.124	.012	1				
MPF/day	.121	.375**	-.119	1			
MS/day	.085	.342**	-.146	.908**	1		
MC/day	-.006	-.160	-0.613**	.103	.091	1	
DMP	-.155*	.232**	.235	.175*	.284**	-.080	1

*correlation is significant at the 0.05 level and **highly significant at the 0.01 level ERCMP=experience of raising cattle for milk productions, MPF= milk produced at farm, MS= milk sold, MC= milk consumed and DMP= distance of marketing place.

Physicochemical and microbiological quality of milk

The average chemical compositions of milk for fat (%), protein (%) and solids not fat (%) content were 3.6043 ± 0.12200 , 2.9749 ± 0.05147 and 6.9992 ± 0.13452 in raw milk samples mixture from producer respectively. Additionally, the mean of milk chemical compositions for fat (%), protein (%) and solids not fat (%) content were 3.3243 ± 0.15814 , 2.8929 ± 0.08510 and

6.7114 ± 0.24844 in raw milk samples mixture from collector respectively. The average physical properties of milk sample indicates on (Table 11) with density, freezing point 1.02721 ± 0.000477 and -0.47143 ± 0.00774 in raw milk sample from producer; 1.02623 ± 0.000874 and -0.45788 ± 0.016510 in raw milk sample from collector respectively.

Table 11. Physicochemical properties of milk at farm and collection points in study area.

Variables and category		N	Mean ± Std. Error	95% CI
Fat (%)	Producer	49	3.6043 ± 0.12200	3.3590 - 3.8496
	Collector	7	3.3243 ± 0.15814	2.9373 - 3.7112
	Overall	56	3.5693 ± 0.10892	3.3510 - 3.7876
Protein (%)	Producer	49	2.9749 ± 0.05147	2.8714 - 3.0784
	Collector	7	2.8929 ± 0.08510	2.6846 - 3.1011
	Overall	56	2.9646 ± 0.04621	2.8720 - 3.0572
SNF (%)	Producer	49	6.9992 ± 0.13452	6.7287 - 7.2696
	Collector	7	6.7114 ± 0.24844	6.1035 - 7.3193
	Overall	56	6.9632 ± 0.12175	6.7192 - 7.2072
Density	Producer	49	1.02721 ± 0.000477	1.02625 - 1.02817
	Collector	7	1.02623 ± 0.000874	1.02410 - 1.02837
	Overall	56	1.02709 ± 0.000432	1.02622 - 1.02795
Added water (%)	Producer	40	14.4087 ± 1.45661	11.4625 - 17.3550
	Collector	7	14.2700 ± 3.05519	6.7942 - 21.7458
	Overall	47	14.3881 ± 1.30856	11.7541 - 17.0221
Freezing point	Producer	49	-0.47143 ± 0.00774	-0.4870 - (-0.4559)
	Collector	7	-0.45788 ± 0.016510	-0.4983 - (-0.4175)
	Overall	56	-0.46974 ± 0.007066	-0.4839 - (-0.4556)

S.E = Standard error, C.I = confidence interval N= number of sample

The overall Average of microbiological count of log (TBC cfu/ml), log (CC cfu/ml) and log (SCC/ml) of raw milk was 8.2577 ± 0.10499, 3.3210 ± 0.11295 and 5.0806 ± 0.08484 for milk sample from the producer; 8.2577 ± 0.10499, 3.3400 ± 0.10352 and 5.1205 ± 0.07533 for milk sample from collector respectively (Table 12).

Table 12. Microbiological quality of milk at farm and collection points in study area.

Variables and category		N	Mean ± Std. Error	95% CI
log(CC/cfu/ml)	Producer	49	3.3210 ± 0.11295	3.0939 - 3.5480
	Collector	8	3.4569 ± 0.26971	2.8191 - 4.0946
	Overall	57	3.3400 ± 0.10352	3.1327 - 3.5474
log(TBC/cfu/ml)	Producer	44	8.2572 ± 0.11195	8.0314 - 8.4830
	Collector	8	8.2601 ± 0.31391	7.5179 - 9.0024
	Overall	52	8.2577 ± 0.10499	8.0469 - 8.4684
log(SCC)	Producer	47	5.0806 ± 0.08484	4.9098 - 5.2513
	Collector	8	5.3548 ± 0.11766	5.0766 - 5.6331

Overall	55	5.1205 ± 0.07533	4.9694 - 5.2715
---------	----	------------------	-----------------

Log=logarithm in base ten (normal logarithm), CC=coli form count, SCC=somatic cell count, TBC= total bacterial count, cfu=colony forming unit per ml of milk sample, S.E = standard error, C.I = confidence interval

The overall mean of chemical compositions of milk for fat (%), protein (%) and solids not fat (%) contents were 3.5693 ± 0.10892, 2.9646 ± 0.04621 and 6.9632 ± 0.12175 in raw milk sample mixture from the two districts respectively. Additionally the overall mean of physical properties of milk sample indicates on (Table 13) with density, freezing point 1.02665 ± 0.00061 and -0.47088 ± 0.0070 in raw milk sample from the two districts. The ANOVA showed significance difference (P< 0.05) due to the source area of raw milk samples for fat, protein and freezing point. Moreover, ANOVA showed highly significance difference at (P<0.01) due to the source area of raw milk samples for solid not fat.

Table 13. Physicochemical properties of milk for the two districts.

Variable		N	Mean ±Std. Error	95% CI	Df	F	P
Fat (%)	Sululta	27	3.3185 ±0.13027	3.0507 - 3.5863	1	5.32	0.025*
	Wolmera	29	3.8028 ±0.16215	3.4706 - 4.1349	54		
	Overall	56	3.5693 ±0.10892	3.3510 - 3.7876	55		
Protein (%)	Sululta	27	2.8485 ±0.05218	2.7413 - 2.9558	1	6.46	0.014*
	Wolmera	29	3.0728 ±0.06985	2.9297 - 3.2158	54		
	Overall	56	2.9646 ±0.04621	2.8720 - 3.0572	55		
SNF (%)	Sululta	27	6.6307 ±0.14321	6.3364 - 6.9251	1	7.80	0.007* *
	Wolmera	29	7.2728 ±0.17720	6.9098 - 7.6357	54		
	Overall	56	6.9632 ±0.12175	6.7192 - 7.2072	55		
Density	Sululta	27	1.02593 ±0.00051	1.02488 - 1.02698	1	1.27	0.266
	Wolmera	30	1.02730 ±0.00105	1.02514 - 1.02945	55		
	Overall	57	1.02665 ±0.00061	1.02544 - 1.02787	56		
Added water	Sululta	26	15.4562 ±1.80236	11.7441 - 19.1682	1	0.82	0.37
	Wolmera	21	13.0657 ±1.90496	9.0920 - 17.0394	45		
	Overall	47	14.3881 ±1.30856	11.7541 -17.0221	46		
Freezing point	Sululta	27	-0.45456±0.00987	-0.47485-(-0.43428)	1	5.21	0.026*
	Wolmera	30	-0.48557±0.00934	-0.50468 -(-0.46647)	55		
	Overall	57	-0.47088 ±0.0070	-0.48498- (-0.45679)	56		

** highly significant at the (P< 0.01) and* is significant at the (P< 0.05) level.

S.E = standard error, C.I = confidence interval, Df= degree of freedom p= p value

The overall mean of microbiological count of log (TBC cfu/ml), log (CC cfu/ml) and log (SCC/ml) of raw milk was 8.2285 ±0.10041, 3.3363 ±0.10010 and 5.1622 ±0.07382 for milk sample from the two districts respectively (Table 14). The ANOVA

showed significance difference at (P<0.01) due to the source area for log (TBC cfu/ml)

Table 14. Microbiological quality of milk for the two districts of study area

Variable	N	Mean ±Std. Error	95% CI	Df	F	P
log(CC/cfu/ml)						
Sululta	30	3.3925 ±0.14411	3.0978 - 3.6873	1	.302	0.585
Wolmera	31	3.2819 ±0.14074	2.9944 - 3.5693	59		
Overall	61	3.3363 ±0.10010	3.1361 - 3.5365	60		
log(TBC/cfu/ml)						
Sululta	27	7.9548 ±0.12902	7.6896 - 8.2200	1	7.774	0.007**
Wolmera	29	8.4834 ±0.13799	8.2008 - 8.7661	54		
Overall	56	8.2285 ±0.10041	8.0273- 8.4298	55		
log(SCC)						
Sululta	28	5.2643 ±0.10299	5.0530 - 5.4757	1	1.753	0.191
Wolmera	31	5.0699 ±0.10406	4.8573 - 5.2824	57		
Overall	59	5.1622 ±0.07382	5.0144 - 5.3099	58		

** Highly significant at the (P< 0.01).

Log=logarithm in base ten (normal logarithm), CC=coli form count, SCC=somatic cell count, TBC= total bacterial count, cfu=colony forming unit per ml of milk sample, S.E = standard error, C.I = confidence interval

Relationship among and between physicochemical and microbiological test of Milk

Milk protein was positively and significantly correlated with fat, solid not fat and density of milk (P<0.01) and significantly correlated with each other (P<0.01). Milk protein was

negatively and significantly correlated with added water and freezing point. Fat, solid not fat (SNF) and density of milk were also negatively and significantly correlated with added water and freezing point (P<0.01) (Table 15).

Table15. Correlations among different characteristics of physicochemical and microbiological test of milk from selected area of small holder dairy farmers

Milk	Fat (%)	Prot. (%)	SNF	Density	AW	Fp	log(CC)	log(TBc)	log(SCC)
Fat (%)	1								
Prot. (%)	.671**	1							
SNF	.624**	.977**	1						
Density	.565**	.969**	.997**	1					
AW	-.562**	-.951**	-.984**	-.976**	1				
Fp	-.634**	-.918**	-.942**	-.529**	1.000**	1			
log(CC)	-.148	-.062	-.068	-.141	.089	.053	1		
log(TBC)	.138	.075	.070	.030	-.066	-.081	-.075	1	
log(SCC)	.068	.075	.074	.001	-.228	-.127	-.097	.005	1

** . Correlation is significant at the (P< 0.01) level.

SNF= solid not fat, AW= added water, Fp= freezing point, log (CC) = logarithms of Coli form count, log (TBC) =logarithms of total bacterial count and log (SCC) = logarithms of somatic cell count.

IV. DISCUSSION

The overall mean family size obtained in the present study for all respondents was 5.69 ± 1.87 person's less than those reported by [10] for Girar Jarso (5.77 persons) and by [11], 6.12 persons per house hold at kuyu wored. The family size ranged from 2 to 12 persons at the study area which is comparable with the report of [12] that family size ranged from 1 to 13 persons in Addis Ababa. About 78.9% of the households were basic educations and above in this study. This value is by far higher than the report of [13] for Gondar area (38.5%). This is mainly indicate that the education coverage between the study areas were different.

The cattle herd size of the study area was 12.27 TLU. The work of [14] indicated that the cattle herd size at Bilalo and Lemmu areas are 8.57 TLU and 10.38 TLU respectively .In the present study area the cattle herd was dominated by crossbreds that results in larger TLU cattle herd size as compared with Bilalo and Lemmu of Arsi area.

The present study also showed that milk production was positively and significantly correlated with experience of raising cattle for milk productions, raw milk sold ($p < 0.01$) and significantly correlated with distance of milk marketing ($p < 0.05$). Whereas the family sizes were not correlated with cattle herd size. On contrast finding reported by [12] and [14] indicated that family size and cattle herd size were positively and significantly correlated. This variations may be due to hired labor was means of overcoming family labor resource.

The average milk yield of cross bred cows in the study area was 9.11 ± 2.902 litres per day, which was comparable with average milk yield of 10 liters reported by [15- 17] Moreover, the average milk yield of local cows was 1.889 ± 0.6707 which was comparable with reported by [18] indicated that the overall average daily milk yield of local cows in the first and second lactations in North Gonder Zone was 1.69 and 1.86 liters, respectively.

The overall average lactation length of local and crossbred cows was 6.26 ± 0.6624 and 9.7 ± 0.46 months, respectively in the study area. The lactation length of the indigenous cows observed

in this study is comparable with the national average of 7 months [19]. The lactation length in crossbred cows observed in this study is shorter than the lactation length of 11.7 months reported for crossbred cows in the central highlands of Ethiopia [20]. The variation in lactation length in the present study may be credited to feed shortage and poor genetic potential of the sample population.

Overall mean of milk producing, Processing, consuming and selling per day per household was 26.88 ± 4.76 , 1.23 ± 1.603 , 1.29 ± 1.176 and 23.32 ± 5.22 liters respectively. Eighty six point seven percent (86.77%) of the milk produced in the area was sold by the producer through different channels. Amount of milk processed, consumed and used for calves was 4.6%, 4.8% and 3.84% respectively. This study is inconsistent with study conducted around Addis Ababa indicated that from total milk production 73% is sold, 10% is left for household consumption, 9.4% goes to calves and 7.6% is processed into butter [21].

Marketing channels are routes through which products pass as they are moved from the farm to the consumer. From this study the main outlets for raw milk identified were cooperatives, processors, vendor, directly to consumer and hotels/restaurants. These are consistent with the result in any marketing system various actors participate in marketing of commodities and process of transactions made. These include itinerate /mobile traders, semi-whole sellers, retailers, cooperatives and consumers as reported by [22]. Collectors collect the milk from the small holder and commercial dairy producers, they sale it to retailers, hotels, restaurants and processors. There exist two types of collectors in the milk value chain. Cooperative collection centers are a formal collectors organized by the bureau of agriculture in their respective districts. They have members of small holder dairy producers which supply daily produce of milk in order to supply to the larger processors in Addis Ababa markets. In addition to collecting from cooperative and individual collectors, larger processors are also collect milk from smallholder farmers giving them additional cents over a liter of milk than other collectors. This condition had negative effect on cooperative collection centers and mutual agreement and win-win approach should be followed among all the actors involving milk supply chain.

Among constraints of milk marketing, price variations, Lack of fair market and Lack of demand during fastening were the most indicated ones. The current study agreed with the report by [23] for Kenyan highlands inaccessibility of fresh milk marketing. Through group discussion almost the entire group member pointed out they have less /no power to decided milk price at the study area. Quality based payment was also another raw milk marketing constraints of the study area. They indicated quality based payment was enhanced quality of milk supplied to processors at the same time as encouraging them to produce more and quality milk. Finally, milk marketing constraints were possessing less preserving facilities for surplus milk produced and demand especially during fasting were great influence on raw milk marketing.

Nearly 19% of the smallholders were using individual towels for cleaning udder of milking cows in 52.2% collective towels were used while in the rest (28.9%) no towel use practiced. It was reported by [24] that pre-milking udder preparations play an important part in the contamination of milk during milking. Most of the dairy owners did not use towel and a few dairy owners used a single towel for all cows commonly to dry the udders. The reuse of towel for cleaning and sanitizing may result in recontamination of the udder. Since drying was not or in sufficiently practiced, contamination level of milk was becoming higher.

The overall mean fat percentage (3.5693 ± 0.10892) of whole milk collected from the smallholder farmers in the current study is less than the fat content of whole milk collected from smallholder farmers reported by [25] for eastern Wollega (6.05%) and also slightly less than reported by [23] for Bahir Dar Zuria (4.14%). The variation in fat percentage observed in the present study may probably due to variation in stage of lactation, feeding regime and parity. The overall mean protein (2.9646 ± 0.04621) content from bulk milk obtained in the current study is lower than those reported by O'Connor (1994) for local cows' milk and also lower than [26] for whole milk in the central highlands of Ethiopia (3.1%). The average SNF (6.9632 ± 0.12175) content of milk obtained in the current study is slightly lower than reported by [22] for eastern Wollega (8.22%).

The overall mean total bacterial count of cows' milk produced in the study area was $8.2285 \log_{10} \text{cfu/ml}$. The total bacterial count obtained in this study is generally high as compared to the acceptable level of 1×10^5 bacteria per ml of raw milk [27] The current study is consistent with [28] reported that the minimum and maximum total bacterial count of raw cows' milk produced in southern region to be 6 to $8.8 \log_{10} \text{cfu/ml}$. Commonly, lack of knowledge about clean milk production and use of unclean milking equipment would be some of the factors which contributed to the poor hygienic quality of milk produced in the study area.

The overall mean coliform count of milk produced in the area was $3.3363 \log_{10} \text{cfu/ml}$. The coliform count of cows' milk obtained in the current study is smaller than with reported by [26] for districts of southern region ($3.8 \log_{10} \text{cfu/ml}$). The current result is also inconsistent with the reported by [29] for cows' milk collected from different producers in the central highland of Ethiopia ($6.57 \log_{10} \text{cfu/ml}$). The higher coliform count obtained in this study may be due to the initial contamination of the milk samples either from the cows, the milkers, milk containers and the

milking environment. The overall mean of somatic cell count in log (SCC/ml) of raw milk was 5.1622 ± 0.07382 for milk sample from the two districts.

V. CONCLUSIONS

Dairy production became a crucial element of the farming activities and income generating for household in "Sululta" and "Wolmera" district of Oromia special zone surrounding Addis Ababa.

The proportion of raw milk used for household consumption was relatively small and the major part of milk produced by smallholders is destined to market. Smallholders also process milk to butter and cheese. Milk was soured for 2-3 days before processing it into butter and cheese. The main outlets for raw milk identified were cooperatives, processors, vendor, directly to consumer and Hotels/restaurants. Price variations, lack of fair market, lack of demand during fastening, lack of quality based payment and lack of preserving facilities were the major problem of raw milk marketing in the study areas.

Hygienic conditions of milking and storage processes, transferring of milk into different containers and sieves, unclean milk equipment were basic determinants of milk quality. Majority of raw milk samples from producer and collector bulk milk sample had higher TAPC and coliform counts, which was higher than the international acceptable limits.

VI. CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

VII. ACKNOWLEDGMENT

The Authors would like to thank Ministry of Education and Agriculture bureau, Oromia special zone for their co-operation and offering facilities. The Authors also thank dairy cooperatives, farm attendants, farm managers, AI technicians and veterinary professionals working in dairy farms who agreed to participate in the data collection process.

REFERENCES

- [1] Pratt, A., Staal, S & Jabbar M. (2008). Dairy Development for the Resources Poor Policy Initiative of Kenya and Ethiopia, Dairy Development Case Studies. Rome, Italy: Pro-Poor Livestock, Part 2.
- [2] CSA (Central Statistical Authority) (2008). Livestock and Livestock Characteristics. Agricultural Sample Survey, Statistical Bulletin, Volume II, Addis Ababa, Ethiopia.
- [3] Ahmed, M. Ehui, S. and Yemesrach, A. (2003). Milk development in Ethiopia. Socio- economics and policy research International Livestock Research Institute, Nairobi, Kenya, Working paper, 58:47.
- [4] Holloway, G., Nicholson, C., Delgado, C., Staal, C.S. and Ehui, S. (2000). How to make milk market: A case study from the Ethiopian highlands. Socio-economic and policy research, ILRI, Nairobi, Kenya, working paper, 28: 28.
- [5] Berhanu G., D., Hoekstra, and Azage Tegegne, (2006). Commercialization of Ethiopian Agriculture: extension service from input supplier to knowledge broker and facilitator, Working Paper no. 1, Nairobi, Kenya: International Livestock Research Institute
- [6] Ayele, S., Assegid, W., Jabbar, M., Ahmed, M. and Belachew, H. (2003). Livestock marketing in Ethiopia: A review of structure, performance and development initiatives, Socio-economic and Policy Research, ILRI, Nairobi, Kenya, Working Paper 52:35.

- [7] Arsham H., (2007). Arsham Mathematical and Computer Modelling Oxford University Press, London.
- [8] American Public Health Association (APHA). 1993. Standard Methods for the Examination of Dairy Products, 16th ed. APHA, Washington, DC.
- [9] IDF (1995) Methods for Assessing the Bacteriological Quality of Raw Milk from the Farm. Brussels, Belgium. Bulletin No 256. PP 4-8.
- [10] Tolera, D. (2007). Smallholder dairy production technology transfer and adoption constraints in mixed farming system in Girar Jarso District of North Shoa Zone Oromia Regional State. MSc. Thesis, Addis Ababa University, Faculty of Veterinary Medicine, Debre-Zeit, Ethiopia.
- [11] Abera K. (2008). Milk production and marketing in smallholder dairy in Kuy District of North Shoa zone, Oromia regional state. MSc. Thesis, Addis Ababa University, Faculty of Veterinary Medicine, Debre-Zeit, Ethiopia.
- [12] Kelay, B. (2002). Analysis of dairy cattle breeding practices in selected areas of Ethiopia. PhD. Thesis, Humboldt University of Berlin, Department of Animal Breeding in the Tropics and Sub Tropics. Berlin, Germany.
- [13] Sisay A. (2006). Qualitative and Quantitative Aspect of Animal Feed in Different Agroecological Areas of North Gonder. MSc Thesis, Alemaya University, Alemaya, Ethiopia.
- [14] Abdnasir, I.B. (2000). Smallholder dairy production and dairy technology adoption in the mixed farming in Arsi highlands, Ethiopia. PhD. Thesis, Humboldt University of Berlin, Department of Animal Breeding in Tropics and Sub tropics, Germany.
- [15] Yoseph, M. (1999). Impacts feed resources on productive and reproductive performance of dairy cow in the peri-urban dairy production systems in the Addis Ababa milk shed and evaluation of non-conventional feed resources using Sheep, MSc. Thesis, School of Graduate Studies, Alemaya University of Agriculture, Alemaya, Ethiopia.
- [16] Azage, T. and Alemu, G. (1998). Prospects for peri-urban milk development in Ethiopia. In: Proceedings of 5th National Conference of Ethiopian Society of Animal Production, May15-17, 1997, Addis Ababa. Ethiopia, Pp 248.
- [17] Mekonnen, H.M., Asmamaw, K., Courreau, J.F. (2006). Husbandry practices and health in smallholder dairy farms near Addis Ababa, Ethiopia. *Preventive Veterinary Medicine*, **74**:99-107.
- [18] Zewdu W. (2004). Indigenous cattle genetic resources, their husbandry practices and breeding objectives in North-western Ethiopia. M.Sc. Thesis. Alemaya University of Agriculture, DireDawa, Ethiopia. 127p.
- [19] CSA (Central Statistical Authority) (2005). Livestock and livestock characteristics. Agricultural Sample Survey, Statistical Bulletin, Volume II, Addis Ababa, Ethiopia.
- [20] Zelalem, Y. and Ledin, I. (2001a). Milk production, processing, marketing and the role of milk and milk products on smallholder farmers' income in the central highlands of Ethiopia. In: Proceedings of the 8th Annual Conference of the Ethiopian Society of Animal Production, August 24-26, 2000. Addis Ababa, Ethiopia, Pp 139-154.
- [21] Baltenweck, I. A. and Staal, S. J. (2000). Determinants of adoption of dairy cattle technology in the Kenya highlands: a Spatial and dynamic Approach. International Livestock Institute. Nairobi, Kenya, Pp 1-7.
- [22] Alganesh, T. (2002). Traditional milk and milk products handling practices and raw milk quality in Eastern Wollega. M.Sc. Thesis, Alemaya University, Alemaya, Ethiopia.
- [23] Asaminew T. (2007). Production, handling, traditional processing practices and quality of milk in Bahir Dar milk shed area, Ethiopia. MSc. Thesis, Haramaya University, Haramaya.
- [24] Zelalem Y. and Ledin I. (2001b). Efficiency of Smallholder butter-making in the central highlands of. In: Proceedings of the 8th Annual Conference of the Ethiopian Society of Animal Production (ESAP)
- [25] O'Connor, C.B. (1994). ILRI Rural Dairy Technology training manual No. 1. International Livestock Research Institute (ILRI), Addis Ababa, Ethiopia. 133p.
- [26] Fekadu, B. (1994). Present situation and future aspects of milk production, milk handling and processing of dairy products in Southern Ethiopia. Food production strategies and limitations: The case of Aneno, Bulbula and Dongora in Southern Ethiopia. Ph.D. Thesis, Department of Food Science. Agricultural University of Norway, Norway.
- [27] Zelalem Y. and Bernard F., (2006). Handling and microbial load of cow's milk and irgo- fermented milk collected from different shops and producers in central highlands of Ethiopia. *Ethiopian Journal of Animal Production*. 6(2)-2006:67-82.

AUTHORS

First Author – Melesse Etifu MSc. (Tropical Animal Production and Health), Wolkite University Department of Animal Production and Technology.

E-mail: etifumelese16@gmail.com

Second Author – Mustefa Abu MSc. (Tropical Animal Production and Health), Federal Ministry of Agriculture, Addis Ababa, E- mail: salimlmak@gmail.com

Correspondence Author – Melesse Etifu

E-mail: etifumelese16@gmail.com

Cell phone: +251-911376067