

Equations for Planetary Ellipses

Eric Sullivan*

*Pittsford Mendon High School, Student, Class of 2016

Abstract- Planetary orbits are ellipses with the sun at one of the foci. The semi major axis of each planetary orbital was used in part with each planets eccentricity to calculate the semi minor axis and the location of the foci. Equations in standard ellipse form were created for each of the planets. In the first model, the sun is placed at (0,0). With this set-up, the equations can be completely derived. Once the equations have been derived, the location of the sun was shifted to the positive (c,0) value. The distances for perihelion and aphelion were calculated by an adapted version of the standard distance equation. After this the accepted value for perihelion and aphelion for each planet was compared to the value measured. The percent error was then calculated.

Index Terms- Aphelion, eccentricity, ellipse equations, perihelion, planetary orbits

I. INTRODUCTION

Planetary orbits and how the planets move is useful information to predict certain astronomical events. The orbits of the eight major planets are ellipses where the sun is located at one of the foci. Each planetary orbit is slightly greater than the preceding orbit, making it so they can never intersect or collide. During the course of this research, the data was converted from kilometers into gigameters. Light-years, parsecs or astronomical units were not used since they were either too large or too small. A gigameter is equivalent a million kilometers thus making the data much easier to work with.

II. IDENTIFY, RESEARCH AND COLLECT IDEA

The idea of creating planetary orbital equations came about during a math lecture on ellipses and conical equations. The information about eccentricity and the semi major axis brought the idea about. A google search revealed no examples of equations related to planetary orbitals. But upon a quick conversation with a math professor and a quick glance through my math textbook made me realize that deriving such equations was possible.

All of the values present are measured in gigameters. The use of gigameters was out of convenience because the astronomical unit, light-year, parsec and kilometers were rather large.

III. WRITE DOWN YOUR STUDIES AND FINDINGS

All of the values present are measured in gigameters. The use of gigameters was out of convenience because the astronomical unit, light-year, parsec and kilometers were rather large.

The standard form of an ellipse with the major axis horizontal can be defined as,

$$\frac{(x - h)^2}{a^2} + \frac{(y - k)^2}{b^2} = 1 \quad (1)$$

All of the eight planets have orbits with a horizontal major axis, therefore their equations will be written in the form of equation 1. The values for the semi major axis are located in table 1.

The semi major axis extends from the center of the ellipse to the farthest point on one side. Both of these sides, the (-a,0) and (a,0), together make up the major axis. Thus, the major axis can be recorded as the distance between (-a,0) and (a,0). The value for the semi major axis is always greater than the value for the semi minor axis. Therefore, the location of the semi major axis value (denoted as a) corresponds with the direction of the major axis. For all planetary ellipses the a value is located under the x term, meaning that the major axis is horizontal. The values for the eccentricity of a planet's ellipse are recorded in table 1 below. The eccentricity and the semi major axis values allows the value for the location of the foci to be calculated.

Any ellipse has an eccentricity value less than one. If the value for eccentricity is equal to one, the result is a parabola. Lastly, if the eccentricity is greater than one, the outcome is a hyperbola.

Eccentricity is calculated by dividing the distance from the center to a foci by the semi major axis. With both the eccentricity and semi major axis already recorded, the eccentricity equation can be re-arranged to solve for the c value. The c value is the distance from the center of the ellipse to both of the foci. The re-arranged equation allows for the c value for each planetary ellipse to be calculated.

These values are located in table 1.

Now with the a and c values calculated, a simple calculation will produce the b value. The b value is the semi minor axis. The calculated values for b can be found in table 1. The equations for each planet's orbit can now be written because both the a and b terms have values.

It must also be noted that all the values in table 1 have been calculated with respect to the sun being located at (0,0). Once the equations have been written, the sun can be shifted to its respective point at the positive foci with coordinates (c,0). This does not change the center of the ellipse however.

Therefore, the model states the center of all planetary ellipses is not the sun but another point in space that is located at the true (0,0) location.

Respectively, both the a and b values can be filled into their appropriate spots in the general equation for an ellipse as noted by equation 1. The values for h and k in this case are both 0. In table 2, all the equations are recorded in order of the planet's distance from the sun.

By using these equations and moving the sun back to its corresponding c value for each planet, the values for aphelion and perihelion were measured. These measured values from the graphs of these equations were then compared to the accepted values.

Mercury's equation is 2.1, Venus' equation is 2.2, Earth's equation is 2.3 and Mars' equation is 2.4. For the outer planets; equation 2.5 is Jupiter, equation 2.6 is Saturn, equation 2.7 is Uranus and equation 2.8 is Neptune.

From these equations, and shifting the position of the sun back to (c,0), the distances for aphelion and perihelion were computed.

Tables 3 and 4 indicate the percentage error in each calculation. Equations 3.2 and 3.3 are the equations used to calculate the values for perihelion and aphelion. These equations extend from the well-known distance formula that is indicated by equation 3.1.

The notation used by equation 3.2 and 3.3 refers to the points along the major axis of an ellipse. d_p and d_A correspond to perihelion and aphelion respectively.

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \quad (3.1)$$

Perihelion is the closest distance that the planet is to the sun. The values for perihelion based on the equations in table 2 were calculated using equation 3.2.

$$d_p = \sqrt{(a - c)^2} \quad (3.2)$$

Aphelion is the farthest distance from the planet to the sun. These values were obtained by using equation 3.3.

$$d_A = \sqrt{(c + a)^2} \quad (3.3)$$

The accepted values in the accepted columns in table 3 and 4 were then compared with the values calculated by equations 3.2 and 3.3. The percentage error was then calculated for each planet for both aphelion and perihelion. The percent error as well as the measured values for aphelion and perihelion are located in table 3 and 4.

The equation used to determine the percentage error is represented by equation 4 below.

$$\% \text{ error} = \frac{|\text{accepted} - \text{measured}|}{\text{accepted}} * 100 \quad (4)$$

Tables:

Planet Name	a	b	c	ϵ
Mercury	57.9	56.6703	11.8695	0.205
Venus	108	107.9974	0.756	0.007
Earth	150	149.9783	2.55	0.017
Mars	228	226.9905	21.432	0.094
Jupiter	779	778.0643	38.171	0.049
Saturn	1430	488.1149	81.51	0.057
Uranus	2870	2866.9619	132.02	0.046
Neptune	4500	4499.7277	49.5	0.011

$\frac{x^2}{57.9^2} + \frac{y^2}{56.6703^2} = 1$	(2.1)
$\frac{x^2}{108^2} + \frac{y^2}{107.9974^2} = 1$	(2.2)
$\frac{x^2}{150^2} + \frac{y^2}{149.9783^2} = 1$	(2.3)
$\frac{x^2}{228^2} + \frac{y^2}{226.9905^2} = 1$	(2.4)
$\frac{x^2}{779^2} + \frac{y^2}{778.0643^2} = 1$	(2.5)
$\frac{x^2}{1430^2} + \frac{y^2}{488.1149^2} = 1$	(2.6)
$\frac{x^2}{2870^2} + \frac{y^2}{2866.9619^2} = 1$	(2.7)
$\frac{x^2}{4500^2} + \frac{y^2}{4499.7277^2} = 1$	(2.8)

Planet Name	Measured	Accepted	% Error
Mercury	46.0305	46	0.0663
Venus	107.244	107.5	0.2381
Earth	147.45	147.1	0.2379
Mars	206.568	206.6	0.01549
Jupiter	740.829	740.5	0.0444
Saturn	1348.49	1352.6	0.3039
Uranus	2737.98	2741.3	0.1211
Neptune	4450.5	4444.5	0.134998

Planet Name	Measured	Accepted	% Error
Mercury	69.7695	69.8	0.0437
Venus	108.756	108.9	0.1322
Earth	152.55	152.1	0.2959
Mars	249.432	249.2	0.0931
Jupiter	817.171	816.6	0.0699
Saturn	1511.51	1514.5	0.1974
Uranus	3002.02	3003.6	0.0526
Neptune	4549.5	4545.7	0.08382

IV. CONCLUSION

The equations that have been derived in this experiment demonstrate similar characteristics and are generally accurate because they relatively match the perihelion and aphelion distances from the sun. A similar equation could be written for Pluto's orbit or other Kuiper Belt objects but due to the rotation of the semi major axis those equations would have to be written in polar form. With the creation of an equation for Pluto, intersection and collision points with Neptune can be determined.

ACKNOWLEDGMENT

I would like to thank my math teacher, Mrs. Kimberly Waterbury, for teaching me about ellipses and sparking my interest into their application into planetary patterns. I would also like to thank Mr. Anthony Martellotta, who helped me with some of the math.

REFERENCES

- [1] Williams, Dr. David R. *NASA Planetary Factsheet*. n.d. 20 March 2016.
- [2] Department, The State Education. "Reference Tables for Physical Setting/Earth Science." *Reference Tables for Physical Setting/Earth Science*. Albany, New York: The University of the State of New York, 2011

AUTHORS

First Author – Eric Sullivan, Pittsford Mendon High School, Student, Class of 2016