

# Supersymmetry: Sonderforschungsbereich: Collaborative Research

Dr. K.N.P. Kumar

Post Doctoral fellow, Department of Mathematics, Kuvempu University, Shankaraghatta, Karnataka, India

**Abstract:** In the following two modules we give a concatenated mathematical statement of the supersymmetry, gravity and warped extra dimension,. It is to be stated in unequivocal terms that the classification is based on the characteristics of the systems under investigation. Perspicuous forbearance, Sophisticated reasoning, constituent structure, concept formulation, concept formulation, beat cadence, interiority of perfect causes, incompatibility of conjunctions', resonant resummations, hypothetical propositions, logical compatibilities, on causal correspondences, predicational inherence, emphasis related phenomenological methodologies, Participational observation, Background actuality, Addressed aestheticism, polished perception, refined proficiency, Existential worldliness, Existential strife, Predicational anteriorities, apophthegm and axiom, brocard- gnome, implantation and inculcation, pronouncement and proposition, declaration dogma, Character consonance, shift and stratagem, Ontological consonance, Primordial exactitude, Phenomenological correlates, Accolytish representation, Atrophied asseveration, supposition, surmise, theorization Anamensial alienisms, Anchorite aperitif, Arcadian Atticism all delineated in the conditionalities and functionalities and orinetatilities of the system which incorporate the rules and regulations, axiomatic predications and postulation alcovishness of the foregoing state. **Characterstics of such systems are taken in to consideration in the complete consolidation and consummation of the systems. It is to be noted that despite the fact the treat is done in two piecemeal for the seemingly inexorable and ineluctable process of typing 24 superscripts and subscripts and attendant and sine qua non confusion. Entire monograph is to be read with one whole and conclusions derived thereof follow suit. Kindly bear with me for it is unassailable task to concatenate all the systems together which however the intention is.**

## INTRODUCTION—VARIABLES USED

- (1) If this symmetry exists, it must be **broken(e&eb)**
- (2) If this symmetry exists the superpartners **would (=) have** the same exact masses as the normal particles, and hence would've been discovered by now.
- (3) If SUSY is to exist at the appropriate scale to solve the hierarchy problem, the LHC — once it reaches its full energy of 14 TeV — **ought (eb)to find** at least one superpartner
- (4) If SUSY is to exist at the appropriate scale to solve the hierarchy problem, the LHC — once it reaches its full energy of 14 TeV **ought (eb) to find** as well as at least a second Higgs particle.
- (5) Otherwise, the existence of very heavy superpartners **would create (eb)** yet another puzzling hierarchy problem, one with no good solution.
- (6) For those of you wondering, the absence of SUSY particles at all energies would be enough **to (e) invalidate** string theory, as supersymmetry is a requirement of string theories that contain the standard model of particles.




Image credit: Matt Strassler.

- (7) Technicolor. No, this isn't a 1950s cartoon; Technicolor is the term for physics theories that **require(e)** new gauge interactions,
- (8) Technicolor is the term for physics theories also **that (e) have** either no Higgs particles or unstable/unobservable (i.e., composite) Higgses.
- (9) There would also be an interesting new slew of observable particles. Although this could have been a plausible solution in principle, the recent discovery of what appears to be a fundamental; spin-0 scalar at the right energy to be the Higgs seems **to (e) invalidate** this possible solution to the hierarchy problem.

**NOTATION**

**Module One**

If this symmetry exists, it must be **broken(e&eb)**

$G_{13}$  : Category one of symmetry exists

$G_{14}$  : Category two of symmetry exists

$G_{15}$  : Category three of symmetry exists

$T_{13}$  : Category one of symmetry must be broken

$T_{14}$  : Category two of symmetry must be broken

$T_{15}$  : Category three of symmetry must be broken

### Module Two

If this symmetry exists the superpartners **would (=) have** the same exact masses as the normal particles, and hence would've been discovered by now

$G_{16}$  : Category one of same exact masses as the normal particles, and hence would've been discovered by now

$G_{17}$  : Category two of same exact masses as the normal particles, and hence would've been discovered by now

$G_{18}$  : Category three of same exact masses as the normal particles, and hence would've been discovered by now

$T_{16}$  : Category one of symmetry exists the superpartners

$T_{17}$  : Category two of symmetry exists the superpartners

$T_{18}$  : Category three of symmetry exists the superpartners

### Module three

If SUSY is to exist at the appropriate scale to solve the hierarchy problem, the LHC — once it reaches its full energy of 14 TeV — **ought (eb)to find** at least one superpartner

$G_{20}$  : Category one of SUSY is to exist at the appropriate scale to solve the hierarchy problem, the LHC — once it reaches its full energy of 14 TeV

$G_{21}$  : Category two of SUSY is to exist at the appropriate scale to solve the hierarchy problem, the LHC — once it reaches its full energy of 14 TeV

$G_{22}$  : Category three of SUSY is to exist at the appropriate scale to solve the hierarchy problem, the LHC — once it reaches its full energy of 14 TeV

$T_{20}$  : Category one of at least one superpartner. Note that we are talking of the investigatable systems that satisfy the definition and statement made in the foregoing of the symmetry. Perspicuous forbearance, Sophisticated seasoning, constituent structure, concept formulation, concept formulation, beat cadence, interiority of perfect causes, incompatibility of conjunctions', resonantial resumptions, hypothetical propositions, logical compatibilities, on causal correspondences, predicational inherence, emphasis related phenomenological methodologies, Participational observation, Background actuality, Addressed aestheticism, polished perception, refined proficiency, Existential worldliness, Existential strife, Predicational anteriorities, apophthegm and axiom, brocard- gnome, implantation and inculcation, pronouncement and proposition, declaration dogma, Character consonance, shift and stratagem, Ontological consonance, Primordial exactitude, Phenomenological correlates, Accolytish representation, Atrophied asseveration, supposition, surmise, theorization Anamensial alienisms, Anchorite aperitif, Arcadian Atticism all delineated in the conditionalities and functionalities and orinetatilities of the system which incorporate the rules and regulations, axiomatic predications and postulation alcovishness of the foregoing state. **Characterstics of such systems are taken in to consideration in the complete consolidation and consummation of the systems**

$T_{21}$  : Category two of at least one superpartner

$T_{22}$  : Category three of at least one superpartner

#### Module four

If SUSY is to exist at the appropriate scale to solve the hierarchy problem, the LHC — once it reaches its full energy of 14 TeV **ought (eb) to find** as well as at least a second Higgs particle

$G_{24}$  : Category one of SUSY is to exist at the appropriate scale to solve the hierarchy problem, the LHC — once it reaches its full energy of 14 TeV

$G_{25}$  : Category two of SUSY is to exist at the appropriate scale to solve the hierarchy problem, the LHC — once it reaches its full energy of 14 TeV

$G_{26}$  : Category three of SUSY is to exist at the appropriate scale to solve the hierarchy problem, the LHC — once it reaches its full energy of 14 TeV

$T_{24}$  : Category one of at least a second Higgs particle. This does not mean there are 3 or 6 Higgs bosons. we reiterate we are talking of the characteristics of the investigatable system which bear the characteristics, namely possession of the mass giving particle. And there are differential masses. Primordial exactitude, Phenomenological correlates, Accolytish representation, Atrophied asseveration, supposition, surmise, theorization Anamensial alienisms, Anchorite aperitif, Arcadian Atticism all delineated in the conditionalities and functionalities and orinetatilities of the system which incorporate the rules and regulations, axiomatic predications and postulation alcovishness of the foregoing state. **Characterstics of such systems are taken in to consideration in the complete consolidation and consummation of the systems**

$T_{25}$  : Category two of at least a second Higgs particle

$T_{26}$  : Category three of at least a second Higgs particle

#### Module five

Otherwise, the existence of very heavy superpartners **would create (eb)** yet another puzzling hierarchy problem, one with no good solution

$G_{28}$  : Category one of existence of very heavy superpartners

$G_{29}$  : Category two of existence of very heavy superpartners

$G_{30}$  : Category three of existence of very heavy superpartners

$T_{28}$  : Category one of another puzzling hierarchy problem, one with no good solution. Please note the classification is based on the characteristics of the investigatable problem, and there are legion. aspectionalities that is taken in to consideration is that they satisfy the conditionalities and functionalities of the main graded system.

$T_{29}$  : Category two of another puzzling hierarchy problem, one with no good solution

$T_{30}$  : Category three of another puzzling hierarchy problem, one with no good solution

#### Module six

For those of you wondering, the absence of SUSY particles at all energies would be enough **to (e) invalidate** string theory, as supersymmetry is a requirement of string theories that contain the standard model of particles

$G_{32}$  : Category one of string theory, as supersymmetry is a requirement of string theories that contain the standard model of particles

$G_{33}$  : Category two of string theory, as supersymmetry is a requirement of string theories that contain the standard model of particles

$G_{34}$  : Category three of string theory, as supersymmetry is a requirement of string theories that contain the standard model of particles

$T_{32}$  : Category one of absence of SUSY particles at all energies would be enough

$T_{33}$  : Category two of absence of SUSY particles at all energies would be enough

$T_{34}$  : Category three of absence of SUSY particles at all energies would be enough

### Module seven

technicolor is the term for physics theories that **require(e)** new gauge interactions

$G_{36}$  : Category one of new gauge interactions

$G_{37}$  : Category two of new gauge interactions

$G_{38}$  : Category three of new gauge interactions

$T_{36}$  : Category one of technicolor is the term for physics theories

$T_{37}$  : Category two of technicolor is the term for physics theories

$T_{38}$  : Category three of technicolor is the term for physics theories

### Module eight

Technicolor is the term for physics theories also **that (e) have** either no Higgs particles or unstable/unobservable (i.e., composite) Higgses

$G_{40}$  : Category one of either no Higgs particles or unstable/unobservable (i.e., composite) Higgses. Perspicuous forbearance, Sophisticated seasoning, constituent structure, concept formulation, concept formulation, beat cadence, interiority of perfect causes, incompatibility of conjunctions', resonantial resumptions, hypothetical propositions, logical compatibilities, on causal correspondences, predicational inherence, emphasis related phenomenological methodologies, Participational observation, Background actuality, Addressed aestheticism, polished perception, refined proficiency, Existential worldliness, Existential strife, Predicational anteriorities, apophthegm and axiom, brocard- gnome, implantation and inculcation, pronouncement and proposition, declaration dogma, Character consonance, shift and stratagem, Ontological consonance, Primordial exactitude, Phenomenological correlates, Accolytish representation, Atrophied asseveration, supposition, surmise, theorization Anamensial alienisms, Anchorite aperitif, Arcadian Atticism all delineated in the conditionalities and functionalities and orinetatilities of the system which incorporate the rules and regulations, axiomatic predications and postulation alcovishness of the foregoing state. **Characterstics of such systems are taken in to consideration in the complete**

**consolidation and consummation of the systems. This holds good for the entire monograph. We will not repeat the same**

$G_{41}$  : Category two of either no Higgs particles or unstable/unobservable (i.e., composite) Higgses

$G_{42}$  : Category three of either no Higgs particles or unstable/unobservable (i.e., composite) Higgses

$T_{40}$  : Category one of Technicolor is the term for physics theories

$T_{41}$  : Category two of Technicolor is the term for physics theories

$T_{42}$  : Category three of Technicolor is the term for physics theories

### Module Nine

There would also be an interesting new slew of observable particles. Although this could have been a plausible solution in principle, the recent discovery of what appears to be a fundamental; spin-0 scalar at the right energy to be the Higgs seems **to (e) invalidate** this possible solution to the hierarchy problem

$G_{44}$  : Category one of possible solution to the hierarchy problem. We are again talking of the systemic characteristics and not the fact there are three solutions. Characteristics of these innumerable systems form the basis of the classification scheme.

$G_{45}$  : Category two of possible solution to the hierarchy problem

$G_{46}$  : Category three of possible solution to the hierarchy problem

$T_{44}$  : Category one of recent discovery of what appears to be a fundamental; spin-0 scalar at the right energy to be the Higgs

$T_{45}$  : Category two of recent discovery of what appears to be a fundamental; spin-0 scalar at the right energy to be the Higgs

$T_{46}$  : Category three of recent discovery of what appears to be a fundamental; spin-0 scalar at the right energy to be the Higgs

### The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)},$   
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$   
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$   
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$   
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$   
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$   
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)},$   
 $(a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$   
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$

$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$   
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$ ,  
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$ ,  
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$ ,

are Dissipation coefficients

**Module Numbered One**

**The differential system of this model is now (Module Numbered one)**

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \\ &+(a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \\ &-(b''_{13})^{(1)}(G, t) = \text{First detritions factor} \end{aligned}$$

**Module Numbered Two**

**The differential system of this model is now (Module numbered two)**

$$\begin{aligned} \frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} & 12 \\ &+(a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor} \\ &-(b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor} \end{aligned}$$

**Module Numbered Three**

**The differential system of this model is now (Module numbered three)**

$$\begin{aligned} \frac{dG_{20}}{dt} &= (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} & 13 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} & 14 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} & 15 \\ \frac{dT_{20}}{dt} &= (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} & 16 \\ \frac{dT_{21}}{dt} &= (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} & 17 \\ \frac{dT_{22}}{dt} &= (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} & 18 \\ &+(a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor} \\ &-(b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor} \end{aligned}$$

**Module Numbered Four**

**The differential system of this model is now (Module numbered Four)**

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor}$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$$

**Module Numbered Five:**

**The differential system of this model is now (Module number five)**

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$$

**Module Numbered Six**

**The differential system of this model is now (Module numbered Six)**

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$$

**Module Numbered Seven:**

**The differential system of this model is now (Seventh Module)**

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)]T_{37} \tag{41}$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38} \tag{42}$$

$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$

**Module Numbered Eight**

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40} \tag{43}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41} \tag{44}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42} \tag{45}$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40} \tag{46}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41} \tag{47}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42} \tag{48}$$

**Module Numbered Nine**

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44} \tag{49}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45} \tag{50}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46} \tag{51}$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44} \tag{52}$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45} \tag{53}$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46} \tag{54}$$

$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$

$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$  55

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[ \begin{array}{l} (a'_{13})^{(1)} \boxed{+(a''_{13})^{(1)}(T_{14}, t)} \boxed{+(a''_{16})^{(2,2)}(T_{17}, t)} \boxed{+(a''_{20})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37}, t)} \boxed{+(a''_{40})^{(8,8)}(T_{41}, t)} \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{13} \tag{55}$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[ \begin{array}{l} (a'_{14})^{(1)} \boxed{+(a''_{14})^{(1)}(T_{14}, t)} \boxed{+(a''_{17})^{(2,2)}(T_{17}, t)} \boxed{+(a''_{21})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37}, t)} \boxed{+(a''_{41})^{(8,8)}(T_{41}, t)} \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{14} \tag{56}$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[ \begin{array}{l} (a'_{15})^{(1)} \boxed{+(a''_{15})^{(1)}(T_{14}, t)} \boxed{+(a''_{18})^{(2,2)}(T_{17}, t)} \boxed{+(a''_{22})^{(3,3)}(T_{21}, t)} \\ \boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37}, t)} \boxed{+(a''_{42})^{(8,8)}(T_{41}, t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{15} \tag{57}$$

Where  $\boxed{(a'_{13})^{(1)}(T_{14}, t)}$ ,  $\boxed{(a'_{14})^{(1)}(T_{14}, t)}$ ,  $\boxed{(a'_{15})^{(1)}(T_{14}, t)}$  are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2)}(T_{17}, t)}$  are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3)}(T_{21}, t)}$  are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)}$  are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)}$  are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)}$  are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{37})^{(7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{36})^{(7,7)}(T_{37}, t)}$  are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{41})^{(8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{42})^{(8,8)}(T_{41}, t)}$  are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$  are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[ \begin{array}{l} \boxed{(b'_{13})^{(1)}(G, t)} \quad \boxed{-(b''_{13})^{(1)}(G, t)} \quad \boxed{-(b''_{16})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{20})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{40})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[ \begin{array}{l} \boxed{(b'_{14})^{(1)}(G, t)} \quad \boxed{-(b''_{14})^{(1)}(G, t)} \quad \boxed{-(b''_{17})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{21})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{41})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[ \begin{array}{l} \boxed{(b'_{15})^{(1)}(G, t)} \quad \boxed{-(b''_{15})^{(1)}(G, t)} \quad \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} \quad \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7)}(G_{39}, t)} \quad \boxed{-(b''_{42})^{(8,8)}(G_{43}, t)} \quad \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15} \quad 60$$

Where  $\boxed{-(b''_{13})^{(1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1)}(G, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2)}(G_{19}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)}$  are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)}$  are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)}$  are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{36})^{(7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7)}(G_{39}, t)}$  are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{41})^{(8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{42})^{(8,8)}(G_{43}, t)}$  are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)}$  are ninth

detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[ \begin{array}{ccc} (a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) & + (a''_{13})^{(1,1)}(T_{14}, t) & + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[ \begin{array}{ccc} (a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t) & + (a''_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[ \begin{array}{ccc} (a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t) & + (a''_{15})^{(1,1)}(T_{14}, t) & + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where  $(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)$ ,  $(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)$ ,  $(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)$  are first augmentation coefficients for category 1, 2 and 3

$(a''_{13})^{(1,1)}(T_{14}, t)$ ,  $(a''_{14})^{(1,1)}(T_{14}, t)$ ,  $(a''_{15})^{(1,1)}(T_{14}, t)$  are second augmentation coefficient for category 1, 2 and 3

$(a''_{20})^{(3,3,3)}(T_{21}, t)$ ,  $(a''_{21})^{(3,3,3)}(T_{21}, t)$ ,  $(a''_{22})^{(3,3,3)}(T_{21}, t)$  are third augmentation coefficient for category 1, 2 and 3

$(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$ ,  $(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$ ,  $(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficient for category 1, 2 and 3

$(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$ ,  $(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$ ,  $(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficient for category 1, 2 and 3

$(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$ ,  $(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$ ,  $(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficient for category 1, 2 and 3

$(a''_{36})^{(7,7,7)}(T_{37}, t)$ ,  $(a''_{37})^{(7,7,7)}(T_{37}, t)$ ,  $(a''_{38})^{(7,7,7)}(T_{37}, t)$  are seventh augmentation coefficient for category 1, 2 and 3

$(a''_{40})^{(8,8,8)}(T_{41}, t)$ ,  $(a''_{41})^{(8,8,8)}(T_{41}, t)$ ,  $(a''_{42})^{(8,8,8)}(T_{41}, t)$  are eight augmentation coefficient for category 1, 2 and 3

$(a''_{44})^{(9,9)}(T_{45}, t)$ ,  $(a''_{45})^{(9,9)}(T_{45}, t)$ ,  $(a''_{46})^{(9,9)}(T_{45}, t)$  are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[ \begin{array}{ccc} (b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}, t) & - (b''_{13})^{(1,1)}(G, t) & - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9)}(G_{47}, t) \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[ \begin{array}{ccc} (b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}, t) & - (b''_{14})^{(1,1)}(G, t) & - (b''_{21})^{(3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9)}(G_{47}, t) \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[ \begin{array}{ccc} (b'_{18})^{(2)} \boxed{-(b''_{18})^{(2)}(G_{19}, t)} & \boxed{-(b'_{15})^{(1,1)}(G, t)} & \boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \tag{66}$$

where  $\boxed{-(b'_{16})^{(2)}(G_{19}, t)}$ ,  $\boxed{-(b'_{17})^{(2)}(G_{19}, t)}$ ,  $\boxed{-(b'_{18})^{(2)}(G_{19}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{13})^{(1,1)}(G, t)}$ ,  $\boxed{-(b'_{14})^{(1,1)}(G, t)}$ ,  $\boxed{-(b'_{15})^{(1,1)}(G, t)}$  are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)}$  are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$  are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$  are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$  are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)}$  are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)}$  are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{46})^{(9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{45})^{(9,9)}(G_{47}, t)}$  are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[ \begin{array}{ccc} (a'_{20})^{(3)} \boxed{+(a''_{20})^{(3)}(T_{21}, t)} & \boxed{+(a'_{16})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{13})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{20} \tag{67}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[ \begin{array}{ccc} (a'_{21})^{(3)} \boxed{+(a''_{21})^{(3)}(T_{21}, t)} & \boxed{+(a'_{17})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{14})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{21} \tag{68}$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[ \begin{array}{ccc} (a'_{22})^{(3)} \boxed{+(a''_{22})^{(3)}(T_{21}, t)} & \boxed{+(a'_{18})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{15})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{22} \tag{69}$$

$\boxed{+(a''_{20})^{(3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3)}(T_{21}, t)}$  are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a'_{16})^{(2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a'_{17})^{(2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a'_{18})^{(2,2,2)}(T_{17}, t)}$  are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a'_{13})^{(1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a'_{14})^{(1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a'_{15})^{(1,1,1)}(T_{14}, t)}$  are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)}$  are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)}$  are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)}$  are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37}, t)}$  are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41}, t)}$  are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{45})^{(9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{46})^{(9,9,9)}(T_{45}, t)}$  are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[ \begin{array}{ccc} \boxed{(b'_{20})^{(3)}(G_{23}, t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{13})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[ \begin{array}{ccc} \boxed{(b'_{21})^{(3)}(G_{23}, t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{14})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[ \begin{array}{ccc} \boxed{(b'_{22})^{(3)}(G_{23}, t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19}, t)} & \boxed{-(b'_{15})^{(1,1,1)}(G, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3)}(G_{23}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2)}(G_{19}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1)}(G, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)}$  are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)}$  are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)}$  are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39}, t)}$  are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43}, t)}$  are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$ ,  $-(b''_{45})^{(9,9,9)}(G_{47}, t)$ ,  $-(b''_{44})^{(9,9,9)}(G_{47}, t)$  are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[ \begin{array}{ccc} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) & + (a''_{28})^{(5,5)}(T_{29}, t) & + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[ \begin{array}{ccc} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) & + (a''_{29})^{(5,5)}(T_{29}, t) & + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[ \begin{array}{ccc} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) & + (a''_{30})^{(5,5)}(T_{29}, t) & + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$ ,  $(a''_{25})^{(4)}(T_{25}, t)$ ,  $(a''_{26})^{(4)}(T_{25}, t)$  are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$ ,  $+(a''_{29})^{(5,5)}(T_{29}, t)$ ,  $+(a''_{30})^{(5,5)}(T_{29}, t)$  are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$ ,  $+(a''_{33})^{(6,6)}(T_{33}, t)$ ,  $+(a''_{34})^{(6,6)}(T_{33}, t)$  are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$ ,  $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$ ,  $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$  are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{17})^{(2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{18})^{(2,2,2,2)}(T_{17}, t)$  are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3)}(T_{21}, t)$ ,  $+(a''_{21})^{(3,3,3,3)}(T_{21}, t)$ ,  $+(a''_{22})^{(3,3,3,3)}(T_{21}, t)$  are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$ ,  $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$ ,  $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$  are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$ ,  $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$ ,  $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$  are eighth augmentation coefficients for category 1, 2 and 3

$+(a''_{46})^{(9,9,9,9)}(T_{45}, t)$ ,  $+(a''_{45})^{(9,9,9,9)}(T_{45}, t)$ ,  $+(a''_{44})^{(9,9,9,9)}(T_{45}, t)$  are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[ \begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[ \begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[ \begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & - (b''_{30})^{(5,5)}(G_{31}, t) & - (b''_{34})^{(6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \tag{78}$$

Where  $-(b''_{24})^{(4)}(G_{27}, t)$ ,  $-(b''_{25})^{(4)}(G_{27}, t)$ ,  $-(b''_{26})^{(4)}(G_{27}, t)$  are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5)}(G_{31}, t)$  are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6,6)}(G_{35}, t)$  are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1,1,1)}(G, t)$ ,  $-(b''_{15})^{(1,1,1,1)}(G, t)$  are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2,2,2,2)}(G_{19}, t)$  are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3,3,3)}(G_{23}, t)$  are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{37})^{(7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{38})^{(7,7,7,7)}(G_{39}, t)$  are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{41})^{(8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{42})^{(8,8,8,8)}(G_{43}, t)$  are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{45})^{(9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{44})^{(9,9,9,9)}(G_{47}, t)$  are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[ \begin{array}{ccc} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) & + (a''_{24})^{(4,4)}(T_{25}, t) & + (a''_{32})^{(6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \tag{79}$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[ \begin{array}{ccc} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) & + (a''_{25})^{(4,4)}(T_{25}, t) & + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \tag{80}$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[ \begin{array}{ccc} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) & + (a''_{26})^{(4,4)}(T_{25}, t) & + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \tag{81}$$

Where  $+(a''_{28})^{(5)}(T_{29}, t)$ ,  $+(a''_{29})^{(5)}(T_{29}, t)$ ,  $+(a''_{30})^{(5)}(T_{29}, t)$  are first augmentation coefficients for category 1, 2 and 3

And  $+(a''_{24})^{(4,4)}(T_{25}, t)$ ,  $+(a''_{25})^{(4,4)}(T_{25}, t)$ ,  $+(a''_{26})^{(4,4)}(T_{25}, t)$  are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$ ,  $+(a''_{33})^{(6,6,6)}(T_{33}, t)$ ,  $+(a''_{34})^{(6,6,6)}(T_{33}, t)$  are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$ ,  $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$ ,  $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$  are fourth augmentation

coefficients for category 1,2, and 3

$$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$$
 are fifth augmentation

coefficients for category 1,2, and 3

$$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$$
 are sixth augmentation

coefficients for category 1,2, 3

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$$
 are seventh augmentation

coefficients for category 1,2, 3

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$$
 are eighth augmentation

coefficients for category 1,2, 3

$$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$$
 are ninth augmentation

coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[ \begin{array}{ccc} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) & - (b''_{24})^{(4,4)}(G_{27}, t) & - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \tag{82}$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[ \begin{array}{ccc} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) & - (b''_{25})^{(4,4)}(G_{27}, t) & - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \tag{83}$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[ \begin{array}{ccc} (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t) & - (b''_{26})^{(4,4)}(G_{27}, t) & - (b''_{34})^{(6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{30} \tag{84}$$

where  $-(b''_{28})^{(5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5)}(G_{31}, t)$  are first detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4)}(G_{27}, t)$ ,  $-(b''_{25})^{(4,4)}(G_{27}, t)$ ,  $-(b''_{26})^{(4,4)}(G_{27}, t)$  are second detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6,6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6,6,6)}(G_{35}, t)$  are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1,1,1,1)}(G, t)$ ,  $-(b''_{15})^{(1,1,1,1,1)}(G, t)$  are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)$  are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)$  are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$  are seventh detrition coefficients for category 1,2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$  are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$  are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \tag{85}$$

$$- \left[ \begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[ \begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \tag{86}$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[ \begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \tag{87}$$

$\boxed{+(a''_{32})^{(6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6)}(T_{33}, t)}$  are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5)}(T_{29}, t)}$  are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4)}(T_{25}, t)}$  are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)}$  - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)}$  - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)}$  sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)}$  seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)}$

Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)}$  ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[ \begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \tag{88}$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[ \begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \tag{89}$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[ \begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{34} \tag{90}$$

$-(b''_{32})^{(6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6)}(G_{35}, t)$  are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5,5)}(G_{31}, t)$  are second detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4)}(G_{27}, t)$ ,  $-(b''_{25})^{(4,4,4)}(G_{27}, t)$ ,  $-(b''_{26})^{(4,4,4)}(G_{27}, t)$  are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{15})^{(1,1,1,1,1,1)}(G, t)$  are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$  are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$  are sixth detrition coefficients for category 1, 2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$  are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)$  are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)$  are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[ \begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \tag{91}$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[ \begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \tag{92}$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[ \begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \tag{93}$$

Where  $(a''_{36})^{(7)}(T_{37}, t)$ ,  $(a''_{37})^{(7)}(T_{37}, t)$ ,  $(a''_{38})^{(7)}(T_{37}, t)$  are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$  are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$  are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$  are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$  are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$  are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$  are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$  are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$  are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} \boxed{(b'_{36})^{(7)}} & \boxed{-(b''_{36})^{(7)}(G_{39}, t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)} & \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)} & \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} \boxed{(b'_{37})^{(7)}} & \boxed{-(b''_{37})^{(7)}(G_{39}, t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)} & \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)} & \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} \boxed{(b'_{38})^{(7)}} & \boxed{-(b''_{38})^{(7)}(G_{39}, t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)} & \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)} & \end{bmatrix} T_{15}$$

Where  $\boxed{-(b''_{36})^{(7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7)}(G_{39}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$  are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$  are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$  are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)$  are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$  are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$  are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)} G_{41} - \left[ \begin{array}{l} (a''_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) + (a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)} G_{40} - \left[ \begin{array}{l} (a''_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) + (a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[ \begin{array}{l} (a''_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where  $+(a''_{40})^{(8)}(T_{41}, t)$ ,  $+(a''_{41})^{(8)}(T_{41}, t)$ ,  $+(a''_{42})^{(8)}(T_{41}, t)$  are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$  are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ ,  $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ ,  $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$  are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ ,  $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ ,  $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ ,  $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ ,  $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ ,  $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ ,  $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)}$  are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}$  are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$  are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[ \begin{array}{ccc} \boxed{(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t)} & \boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[ \begin{array}{ccc} \boxed{(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t)} & \boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[ \begin{array}{ccc} \boxed{(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t)} & \boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b'_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b'_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15}$$

Where  $\boxed{-(b'_{36})^{(7)}(G_{39}, t)}$ ,  $\boxed{-(b'_{37})^{(7)}(G_{39}, t)}$ ,  $\boxed{-(b'_{38})^{(7)}(G_{39}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b'_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b'_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b'_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b'_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b'_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b'_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}$  are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b'_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b'_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)}$  are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b'_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b'_{15})^{(1,1,1,1,1,1,1,1)}(G, t)}$  are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{13})^{(1,1,1,1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b'_{14})^{(1,1,1,1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b'_{38})^{(7,7)}(G_{39}, t)}$  are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b'_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b'_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)}$  are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b'_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b'_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b'_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)}$  are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[ \begin{array}{|c|} \hline (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \quad + (a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) \quad + (a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ \hline + (a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) \quad + (a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) \quad + (a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ \hline + (a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) \quad + (a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) \quad + (a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \\ \hline \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[ \begin{array}{|c|} \hline (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) \quad + (a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) \quad + (a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ \hline + (a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) \quad + (a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) \quad + (a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ \hline + (a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) \quad + (a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) \quad + (a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \\ \hline \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[ \begin{array}{|c|} \hline (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) \quad + (a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) \quad + (a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ \hline + (a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) \quad + (a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) \quad + (a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ \hline + (a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) \quad + (a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) \quad + (a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \\ \hline \end{array} \right] G_{15}$$

Where  $+(a'_{44})^{(9)}(T_{45}, t)$ ,  $+(a'_{45})^{(9)}(T_{45}, t)$ ,  $+(a'_{46})^{(9)}(T_{37}, t)$  are first augmentation coefficients for category 1, 2 and 3

$+(a'_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $+(a'_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $+(a'_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$  are second augmentation coefficient for category 1, 2 and 3

$+(a'_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$ ,  $+(a'_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$ ,  $+(a'_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$  are third augmentation coefficient for category 1, 2 and 3

$+(a'_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$ ,  $+(a'_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$ ,  $+(a'_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficient for category 1, 2 and 3

$+(a'_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$ ,  $+(a'_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$ ,  $+(a'_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficient for category 1, 2 and 3

$+(a'_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$ ,  $+(a'_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$ ,  $+(a'_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficient for category 1, 2 and 3

$+(a'_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$ ,  $+(a'_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$ ,  $+(a'_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$  are Seventh augmentation coefficient for category 1, 2 and 3

$+(a'_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$ ,  $+(a'_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$ ,  $+(a'_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$  are eighth augmentation coefficient for 1,2,3

$+(a'_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$ ,  $+(a'_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$ ,  $+(a'_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$  are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\left[ \begin{array}{ccc} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47}, t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)} T_{44} - \left[ \begin{array}{ccc} (b'_{45})^{(9)} \boxed{-(b''_{45})^{(9)}(G_{47}, t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)} T_{45} - \left[ \begin{array}{ccc} (b'_{46})^{(9)} \boxed{-(b''_{46})^{(9)}(G_{47}, t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{15}$$

Where  $\boxed{-(b''_{44})^{(9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{45})^{(9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{46})^{(9)}(G_{47}, t)}$  are first detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}$  are second detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}$  are third detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}$  are fourth detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)}$  are fifth detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)}$  are sixth detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)}$  are seventh detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)}$  are eighth detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)}$  are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15$$

The functions  $(a''_i)^{(1)}, (b''_i)^{(1)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(1)}, (r_i)^{(1)}$ :

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \tag{98}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

**Definition of**  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$  :

Where  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$  are positive constants and  $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(1)}(T'_{14}, t)$  and  $(a_i'')^{(1)}(T_{14}, t)$ .  $(T'_{14}, t)$  and  $(T_{14}, t)$  are points belonging to the interval  $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$ . It is to be noted that  $(a_i'')^{(1)}(T_{14}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{13})^{(1)} = 1$  then the function  $(a_i'')^{(1)}(T_{14}, t)$ , the first augmentation coefficient attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$  : 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$ , are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

**Definition of**  $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$  : 101

There exists two constants  $(\hat{P}_{13})^{(1)}$  and  $(\hat{Q}_{13})^{(1)}$  which together With  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$  and  $(\hat{B}_{13})^{(1)}$  and the constants  $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions  $(a_i'')^{(2)}, (b_i'')^{(2)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(2)}, (r_i)^{(2)}$ :

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \tag{102}$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

**Definition of**  $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$  : 106

Where  $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$  are positive constants and  $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19})' - (G_{19})| e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(2)}(T_{17}', t)$  and  $(a_i'')^{(2)}(T_{17}, t)$ .  $(T_{17}', t)$  and  $(T_{17}, t)$  are points belonging to the interval  $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$ . It is to be noted that  $(a_i'')^{(2)}(T_{17}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{16})^{(2)} = 1$  then the function  $(a_i'')^{(2)}(T_{17}, t)$ , the first augmentation coefficient attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$  :

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$ , are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

**Definition of**  $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$  :

There exists two constants  $(\hat{P}_{16})^{(2)}$  and  $(\hat{Q}_{16})^{(2)}$  which together with  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$  and  $(\hat{B}_{16})^{(2)}$  and the constants  $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$ ,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 112$$

The functions  $(a_i'')^{(3)}, (b_i'')^{(3)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(3)}, (r_i)^{(3)}$ :

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \tag{113}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

**Definition of**  $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$  :

Where  $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$  are positive constants and  $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T'_{21}, t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T'_{21} - T_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G'_{23}, t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G'_{23} - G_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(3)}(T'_{21}, t)$  and  $(a_i'')^{(3)}(T_{21}, t)$ .  $(T'_{21}, t)$  and  $(T_{21}, t)$  are points belonging to the interval  $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$ . It is to be noted that  $(a_i'')^{(3)}(T_{21}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{20})^{(3)} = 1$  then the function  $(a_i'')^{(3)}(T_{21}, t)$ , the first augmentation coefficient attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$  : 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$ , are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}} , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants  $(\hat{P}_{20})^{(3)}$  and  $(\hat{Q}_{20})^{(3)}$  which together with  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$  and  $(\hat{B}_{20})^{(3)}$  and the constants  $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$ , satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24, 25, 26 \tag{117}$$

The functions  $(a_i'')^{(4)}, (b_i'')^{(4)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(4)}, (r_i)^{(4)}$ :

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \tag{118}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

**Definition of**  $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$  :

Where  $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$  are positive constants and  $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T'_{25}, t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} \|(G_{27}) - (G_{27})'\| e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(4)}(T'_{25}, t)$  and  $(a_i'')^{(4)}(T_{25}, t)$ .  $(T'_{25}, t)$  and  $(T_{25}, t)$  are points belonging to the interval  $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$ . It is to be noted that  $(a_i'')^{(4)}(T_{25}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{24})^{(4)} = 1$  then the function  $(a_i'')^{(4)}(T_{25}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$  : 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$ , are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

**Definition of**  $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$  : 121

There exists two constants  $(\hat{P}_{24})^{(4)}$  and  $(\hat{Q}_{24})^{(4)}$  which together with  $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$  and  $(\hat{B}_{24})^{(4)}$  and the constants  $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30 \tag{122}$$

The functions  $(a_i'')^{(5)}, (b_i'')^{(5)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(5)}, (r_i)^{(5)}$ :

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \tag{123}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

**Definition of**  $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$  :

Where  $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$  are positive constants and  $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T'_{29}, t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T'_{29} - T_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} |(G_{31})' - (G_{31})| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(5)}(T'_{29}, t)$  and  $(a_i'')^{(5)}(T_{29}, t)$ .  $(T'_{29}, t)$  and  $(T_{29}, t)$  are points belonging to the interval  $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$ . It is to be noted that  $(a_i'')^{(5)}(T_{29}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{28})^{(5)} = 1$  then the function  $(a_i'')^{(5)}(T_{29}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$  : 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$ , are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

**Definition of**  $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$  : 126

There exists two constants  $(\hat{P}_{28})^{(5)}$  and  $(\hat{Q}_{28})^{(5)}$  which together with  $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$  and  $(\hat{B}_{28})^{(5)}$  and the constants  $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34 \tag{127}$$

The functions  $(a_i'')^{(6)}, (b_i'')^{(6)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(6)}, (r_i)^{(6)}$ :

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \tag{128}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

**Definition of**  $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$  :

Where  $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$  are positive constants and  $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T'_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T'_{33} - T_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} \|(G_{35}) - (G_{35})'\| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(6)}(T'_{33}, t)$  and  $(a_i'')^{(6)}(T_{33}, t)$ .  $(T'_{33}, t)$  and  $(T_{33}, t)$  are points belonging to the interval  $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$ . It is to be noted that  $(a_i'')^{(6)}(T_{33}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{32})^{(6)} = 1$  then the function  $(a_i'')^{(6)}(T_{33}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$  : 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$ , are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

**Definition of**  $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$  : 130

There exists two constants  $(\hat{P}_{32})^{(6)}$  and  $(\hat{Q}_{32})^{(6)}$  which together with  $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$  and  $(\hat{B}_{32})^{(6)}$  and the constants  $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

(A)  $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36, 37, 38$  131

(B) The functions  $(a_i'')^{(7)}, (b_i'')^{(7)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(7)}, (r_i)^{(7)}$ :

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

$$(C) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

(D)

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

**Definition of**  $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$  :

Where  $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$  are positive constants and  $\boxed{i = 36, 37, 38}$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}(G_{39}', t) - (b_i'')^{(7)}(G_{39}, t)| < (\hat{k}_{36})^{(7)} |(G_{39}) - (G_{39})'| e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(7)}(T_{37}', t)$  and  $(a_i'')^{(7)}(T_{37}, t)$ .  $(T_{37}', t)$  and  $(T_{37}, t)$  are points belonging to the interval  $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$ . It is to be noted that  $(a_i'')^{(7)}(T_{37}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{36})^{(7)} = 1$  then the function  $(a_i'')^{(7)}(T_{37}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$  :

134

(E)  $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$ , are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

**Definition of**  $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$  :

135

(F) There exists two constants  $(\hat{P}_{36})^{(7)}$  and  $(\hat{Q}_{36})^{(7)}$  which together with  $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$  and  $(\hat{B}_{36})^{(7)}$  and the constants  $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}$ ,  $i = 36, 37, 38$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40, 41, 42 \quad 136$$

The functions  $(a''_i)^{(8)}, (b''_i)^{(8)}$  are positive continuous increasing and bounded

**Definition of**  $(p_i)^{(8)}, (r_i)^{(8)}$ : 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

**Definition of**  $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$ :

Where  $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$  are positive constants and  $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} |(G_{43}) - (G_{43})'| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(8)}(T'_{41}, t)$  and  $(a'_i)^{(8)}(T_{41}, t)$ .  $(T'_{41}, t)$  and  $(T_{41}, t)$  are points belonging to the interval  $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$ . It is to be noted that  $(a''_i)^{(8)}(T_{41}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{40})^{(8)} = 1$  then the function  $(a''_i)^{(8)}(T_{41}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$ :

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$ , are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \quad 144$$

**Definition of**  $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$ :

There exists two constants  $(\hat{P}_{40})^{(8)}$  and  $(\hat{Q}_{40})^{(8)}$  which together with  $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$  and the constants  $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$ ,

Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44, 45, 46$$

146  
A

The functions  $(a''_i)^{(9)}, (b''_i)^{(9)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(9)}, (r_i)^{(9)}$ :

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

**Definition of**  $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$ :

Where  $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$  are positive constants and  $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T'_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(9)}(T'_{45}, t)$  and  $(a''_i)^{(9)}(T_{45}, t)$ .  $(T'_{45}, t)$  and  $(T_{45}, t)$  are points belonging to the interval  $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$ . It is to be noted that  $(a''_i)^{(9)}(T_{45}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{44})^{(9)} = 1$  then the function  $(a''_i)^{(9)}(T_{45}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$ :

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$ , are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

**Definition of**  $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$ :

There exists two constants  $(\hat{P}_{44})^{(9)}$  and  $(\hat{Q}_{44})^{(9)}$  which together with  $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$  and  $(\hat{B}_{44})^{(9)}$  and the constants  $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

**Theorem 1:** if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 2 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

**Definition of**  $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t} , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t} , \quad T_i(0) = T_i^0 > 0$$

**Theorem 3 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t} , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t} , \quad T_i(0) = T_i^0 > 0$$

**Theorem 4 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 5 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 6 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 7:** if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 8:** if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 9:** if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Proof:** Consider operator  $\mathcal{A}^{(1)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy 154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[ (a_{13})^{(1)} G_{14}(s_{(13)}) - \left( (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[ (a_{14})^{(1)} G_{13}(s_{(13)}) - \left( (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[ (a_{15})^{(1)} G_{14}(s_{(13)}) - \left( (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[ (b_{13})^{(1)} T_{14}(s_{(13)}) - \left( (b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[ (b_{14})^{(1)} T_{13}(s_{(13)}) - \left( (b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[ (b_{15})^{(1)} T_{14}(s_{(13)}) - \left( (b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where  $s_{(13)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

159

Consider operator  $\mathcal{A}^{(2)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[ (a_{16})^{(2)} G_{17}(s_{(16)}) - \left( (a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[ (a_{17})^{(2)} G_{16}(s_{(16)}) - \left( (a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[ (a_{18})^{(2)} G_{17}(s_{(16)}) - \left( (a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[ (b_{16})^{(2)} T_{17}(s_{(16)}) - \left( (b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[ (b_{17})^{(2)} T_{16}(s_{(16)}) - \left( (b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[ (b_{18})^{(2)} T_{17}(s_{(16)}) - \left( (b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where  $s_{(16)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(3)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[ (a_{20})^{(3)} G_{21}(s_{(20)}) - \left( (a'_{20})^{(3)} + a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[ (a_{21})^{(3)} G_{20}(s_{(20)}) - \left( (a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[ (a_{22})^{(3)} G_{21}(s_{(20)}) - \left( (a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[ (b_{20})^{(3)} T_{21}(s_{(20)}) - \left( (b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[ (b_{21})^{(3)} T_{20}(s_{(20)}) - \left( (b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[ (b_{22})^{(3)} T_{21}(s_{(20)}) - \left( (b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where  $s_{(20)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:** Consider operator  $\mathcal{A}^{(4)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[ (a_{24})^{(4)} G_{25}(s_{(24)}) - \left( (a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[ (a_{25})^{(4)} G_{24}(s_{(24)}) - \left( (a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[ (a_{26})^{(4)} G_{25}(s_{(24)}) - \left( (a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[ (b_{24})^{(4)} T_{25}(s_{(24)}) - \left( (b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[ (b_{25})^{(4)} T_{24}(s_{(24)}) - \left( (b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[ (b_{26})^{(4)} T_{25}(s_{(24)}) - \left( (b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where  $s_{(24)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:** Consider operator  $\mathcal{A}^{(5)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[ (a_{28})^{(5)} G_{29}(s_{(28)}) - \left( (a'_{28})^{(5)} + a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[ (a_{29})^{(5)} G_{28}(s_{(28)}) - \left( (a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[ (a_{30})^{(5)} G_{29}(s_{(28)}) - \left( (a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[ (b_{28})^{(5)} T_{29}(s_{(28)}) - \left( (b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[ (b_{29})^{(5)} T_{28}(s_{(28)}) - \left( (b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[ (b_{30})^{(5)} T_{29}(s_{(28)}) - \left( (b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where  $s_{(28)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(6)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[ (a_{32})^{(6)} G_{33}(s_{(32)}) - \left( (a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[ (a_{33})^{(6)} G_{32}(s_{(32)}) - \left( (a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[ (a_{34})^{(6)} G_{33}(s_{(32)}) - \left( (a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[ (b_{32})^{(6)} T_{33}(s_{(32)}) - \left( (b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[ (b_{33})^{(6)} T_{32}(s_{(32)}) - \left( (b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[ (b_{34})^{(6)} T_{33}(s_{(32)}) - \left( (b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where  $s_{(32)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(7)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[ (a_{36})^{(7)} G_{37}(s_{(36)}) - \left( (a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[ (a_{37})^{(7)} G_{36}(s_{(36)}) - \left( (a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[ (a_{38})^{(7)} G_{37}(s_{(36)}) - \left( (a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[ (b_{36})^{(7)} T_{37}(s_{(36)}) - \left( (b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[ (b_{37})^{(7)} T_{36}(s_{(36)}) - \left( (b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[ (b_{38})^{(7)} T_{37}(s_{(36)}) - \left( (b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where  $s_{(36)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(8)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[ (a_{40})^{(8)} G_{41}(s_{(40)}) - \left( (a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[ (a_{41})^{(8)} G_{40}(s_{(40)}) - \left( (a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[ (a_{42})^{(8)} G_{41}(s_{(40)}) - \left( (a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[ (b_{40})^{(8)} T_{41}(s_{(40)}) - \left( (b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[ (b_{41})^{(8)} T_{40}(s_{(40)}) - \left( (b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[ (b_{42})^{(8)} T_{41}(s_{(40)}) - \left( (b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where  $s_{(40)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(9)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[ (a_{44})^{(9)} G_{45}(s_{(44)}) - \left( (a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[ (a_{45})^{(9)} G_{44}(s_{(44)}) - \left( (a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[ (a_{46})^{(9)} G_{45}(s_{(44)}) - \left( (a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[ (b_{44})^{(9)} T_{45}(s_{(44)}) - \left( (b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[ (b_{45})^{(9)} T_{44}(s_{(44)}) - \left( (b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[ (b_{46})^{(9)} T_{45}(s_{(44)}) - \left( (b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where  $s_{(44)}$  is the integrand that is integrated over an interval  $(0, t)$

The operator  $\mathcal{A}^{(1)}$  maps the space of functions satisfying Equations into itself. Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[ (a_{13})^{(1)} \left( G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left( 1 + (a_{13})^{(1)}t \right) G_{14}^0 + \frac{(a_{13})^{(1)}(\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left( e^{(\hat{M}_{13})^{(1)}t} - 1 \right)$$

From which it follows that

168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)}t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[ \left( (\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{\left( -\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 1

Analogous inequalities hold also for  $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator  $\mathcal{A}^{(2)}$  maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[ (a_{16})^{(2)} \left( G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = \tag{169}$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left( e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[ \left( (\hat{P}_{16})^{(2)} + G_{17}^0 \right) e^{-\left( \frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for  $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator  $\mathcal{A}^{(3)}$  maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[ (a_{20})^{(3)} \left( G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left( e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[ \left( (\hat{P}_{20})^{(3)} + G_{21}^0 \right) e^{-\left( \frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0} \right)} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for  $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator  $\mathcal{A}^{(4)}$  maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[ (a_{24})^{(4)} \left( G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left( e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[ \left( (\hat{P}_{24})^{(4)} + G_{25}^0 \right) e^{-\left( \frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0} \right)} + (\hat{P}_{24})^{(4)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 4

The operator  $\mathcal{A}^{(5)}$  maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[ (a_{28})^{(5)} \left( G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left( e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\tilde{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\tilde{M}_{28})^{(5)}} \left[ ((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 5

The operator  $\mathcal{A}^{(6)}$  maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[ (a_{32})^{(6)} \left( G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\tilde{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\tilde{M}_{32})^{(6)}} \left( e^{(\tilde{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\tilde{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\tilde{M}_{32})^{(6)}} \left[ ((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 6

Analogous inequalities hold also for  $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(a) The operator  $\mathcal{A}^{(7)}$  maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[ (a_{36})^{(7)} \left( G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\tilde{M}_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left( e^{(\tilde{M}_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\tilde{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\tilde{M}_{36})^{(7)}} \left[ ((\hat{P}_{36})^{(7)} + G_{37}^0)e^{-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}} + (\hat{P}_{36})^{(7)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 7

The operator  $\mathcal{A}^{(8)}$  maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[ (a_{40})^{(8)} \left( G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\tilde{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left( e^{(\tilde{M}_{40})^{(8)}t} - 1 \right)$$

180

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\tilde{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\tilde{M}_{40})^{(8)}} \left[ ((\hat{P}_{40})^{(8)} + G_{41}^0)e^{-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}} + (\hat{P}_{40})^{(8)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 8

Analogous inequalities hold also for  $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator  $\mathcal{A}^{(9)}$  maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[ (a_{44})^{(9)} \left( G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$\left( 1 + (a_{44})^{(9)} t \right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left( e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[ \left( (\hat{P}_{44})^{(9)} + G_{45}^0 \right) e^{-\left( \frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0} \right)} + (\hat{P}_{44})^{(9)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 9

Analogous inequalities hold also for  $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take  $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$  and to choose 182

$(\hat{P}_{13})^{(1)}$  and  $(\hat{Q}_{13})^{(1)}$  large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[ (\hat{P}_{13})^{(1)} + \left( (\hat{P}_{13})^{(1)} + G_j^0 \right) e^{-\left( \frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{13})^{(1)} \tag{183}$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[ \left( (\hat{Q}_{13})^{(1)} + T_j^0 \right) e^{-\left( \frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \tag{184}$$

In order that the operator  $\mathcal{A}^{(1)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(1)}$  is a contraction with respect to the metric 185

$$d \left( (G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)} t} \right\}$$

Indeed if we denote

**Definition of  $\tilde{G}, \tilde{T}$  :**  $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} + (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} + G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)} s_{(13)}} e^{(\widehat{M}_{13})^{(1)} s_{(13)}} \} ds_{(13)}$$

Where  $s_{(13)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$|G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)} t} \leq \frac{1}{(\widehat{M}_{13})^{(1)}} \left( (a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)} \right) d \left( (G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}) \right) \tag{186}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 1:** The fact that we supposed  $(a''_{13})^{(1)}$  and  $(b''_{13})^{(1)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$  and  $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(1)}$  and  $(b''_i)^{(1)}$ ,  $i = 13, 14, 15$  depend only on  $T_{14}$  and respectively on  $G$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(1)} - (a''_i)^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)} t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$  and  $((\widehat{M}_{13})^{(1)})_3$  : 187

**Remark 3:** if  $G_{13}$  is bounded, the same property have also  $G_{14}$  and  $G_{15}$ . indeed if

$$G_{13} < (\widehat{M}_{13})^{(1)} \text{ it follows } \frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If  $G_{14}$  or  $G_{15}$  is bounded, the same property follows for  $G_{13}$ ,  $G_{15}$  and  $G_{13}$ ,  $G_{14}$  respectively.

**Remark 4:** If  $G_{13}$  is bounded, from below, the same property holds for  $G_{14}$  and  $G_{15}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{14}$  is bounded from below. 188

**Remark 5:** If  $T_{13}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$  then  $T_{14} \rightarrow \infty$ . 189

**Definition of**  $(m)^{(1)}$  and  $\varepsilon_1$  :

Indeed let  $t_1$  be so that for  $t > t_1$

$$(b_{14}')^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then  $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$  which leads to

$$T_{14} \geq \left( \frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left( \frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for  $T_{15}$  if  $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take  $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$  and to choose 190

$(\hat{P}_{16})^{(2)}$  and  $(\hat{Q}_{16})^{(2)}$  large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[ (\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left( \frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[ ((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left( \frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator  $\mathcal{A}^{(2)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself 193

The operator  $\mathcal{A}^{(2)}$  is a contraction with respect to the metric 194

$$d \left( ((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \right\}$$

Indeed if we denote 195

$$\text{Definition of } \tilde{G}_{19}, \tilde{T}_{19} : (\tilde{G}_{19}, \tilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$$

It results 196

$$|\tilde{G}_{16}^{(1)} - \tilde{G}_{16}^{(2)}| \leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} +$$

$$\int_0^t \{(a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)}(T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)} |(a''_{16})^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)}(T_{17}^{(2)}, s_{(16)})| e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} e^{(\widehat{M}_{16})^{(2)}s_{(16)}} ds_{(16)}$$

Where  $s_{(16)}$  represents integrand that is integrated over the interval  $[0, t]$  197

From the hypotheses it follows

$$|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)}t} \leq$$

$$\frac{1}{(\widehat{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows 198

**Remark 6:** The fact that we supposed  $(a''_{16})^{(2)}$  and  $(b''_{16})^{(2)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$  and  $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$  respectively of  $\mathbb{R}_+$ . 199

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(2)}$  and  $(b''_i)^{(2)}$ ,  $i = 16, 17, 18$  depend only on  $T_{17}$  and respectively on  $(G_{19})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 7:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  200

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t ((a'_i)^{(2)} - (a''_i)^{(2)}(T_{17}(s_{(16)}), s_{(16)})) ds_{(16)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(2)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{16})^{(2)})_1$ ,  $((\widehat{M}_{16})^{(2)})_2$  and  $((\widehat{M}_{16})^{(2)})_3$  : 201

**Remark 8:** if  $G_{16}$  is bounded, the same property have also  $G_{17}$  and  $G_{18}$ . indeed if

$$G_{16} < (\widehat{M}_{16})^{(2)} \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a'_{17})^{(2)}G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a'_{17})^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a'_{18})^{(2)}$$

If  $G_{17}$  or  $G_{18}$  is bounded, the same property follows for  $G_{16}$ ,  $G_{18}$  and  $G_{16}$ ,  $G_{17}$  respectively.

**Remark 9:** If  $G_{16}$  is bounded, from below, the same property holds for  $G_{17}$  and  $G_{18}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{17}$  is bounded from below. 202

**Remark 10:** If  $T_{16}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(2)}((G_{19})(t), t)) = (b'_{17})^{(2)}$  then  $T_{17} \rightarrow \infty$ . 203

**Definition of**  $(m)^{(2)}$  and  $\varepsilon_2$  :

Indeed let  $t_2$  be so that for  $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then  $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$  which leads to 204

$$T_{17} \geq \left( \frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$$T_{17} \geq \left( \frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2} \text{ By taking now } \varepsilon_2 \text{ sufficiently small one sees that } T_{17} \text{ is unbounded.} \quad 205$$

The same property holds for  $T_{18}$  if  $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take  $\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1$  and to choose 207

$(\hat{P}_{20})^{(3)}$  and  $(\hat{Q}_{20})^{(3)}$  large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[ (\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left( \frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[ ((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left( \frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator  $\mathcal{A}^{(3)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself 210

The operator  $\mathcal{A}^{(3)}$  is a contraction with respect to the metric 211

$$d \left( ((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)} t} \right\}$$

Indeed if we denote 212

**Definition of**  $\widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{(G_{23})}, \widetilde{(T_{23})}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned}
 |\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\
 &\int_0^t \{(a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} + \\
 &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}} + \\
 &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\widehat{M}_{20})^{(3)}s_{(20)}} e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\} ds_{(20)}
 \end{aligned}$$

Where  $s_{(20)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned}
 |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\widehat{M}_{20})^{(3)}t} &\leq \\
 \frac{1}{(\widehat{M}_{20})^{(3)}} &\left( (a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)} \right) d \left( ((G_{23})^{(1)}, (T_{23})^{(1)}); ((G_{23})^{(2)}, (T_{23})^{(2)}) \right)
 \end{aligned} \tag{214}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 11:** The fact that we supposed  $(a''_{20})^{(3)}$  and  $(b''_{20})^{(3)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t}$  and  $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t}$  respectively of  $\mathbb{R}_+$ . 215

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(3)}$  and  $(b''_i)^{(3)}$ ,  $i = 20, 21, 22$  depend only on  $T_{21}$  and respectively on  $(G_{23})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 12:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{20})^{(3)})_1, ((\widehat{M}_{20})^{(3)})_2$  and  $((\widehat{M}_{20})^{(3)})_3$  : 217

**Remark 13:** if  $G_{20}$  is bounded, the same property have also  $G_{21}$  and  $G_{22}$ . indeed if

$$G_{20} < (\widehat{M}_{20})^{(3)} \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If  $G_{21}$  or  $G_{22}$  is bounded, the same property follows for  $G_{20}$ ,  $G_{22}$  and  $G_{20}$ ,  $G_{21}$  respectively.

**Remark 14:** If  $G_{20}$  is bounded, from below, the same property holds for  $G_{21}$  and  $G_{22}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{21}$  is bounded from below. 218

**Remark 15:** If  $T_{20}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$  then  $T_{21} \rightarrow \infty$ . 219

**Definition of**  $(m)^{(3)}$  and  $\varepsilon_3$  :

Indeed let  $t_3$  be so that for  $t > t_3$

$$(b_{21}')^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then  $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$  which leads to 220

$$T_{21} \geq \left( \frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$$

If we take  $t$  such that  $e^{-\varepsilon_3 t} = \frac{1}{2}$  it results

$$T_{21} \geq \left( \frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3}$$

By taking now  $\varepsilon_3$  sufficiently small one sees that  $T_{21}$  is unbounded.

The same property holds for  $T_{22}$  if  $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take  $\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} < 1$  and to choose 221

$(\hat{P}_{24})^{(4)}$  and  $(\hat{Q}_{24})^{(4)}$  large to have

$$\frac{(a_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[ (\hat{P}_{24})^{(4)} + ((\hat{P}_{24})^{(4)} + G_j^0) e^{-\left( \frac{(\hat{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[ ((\hat{Q}_{24})^{(4)} + T_j^0) e^{-\left( \frac{(\hat{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{24})^{(4)} \right] \leq (\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator  $\mathcal{A}^{(4)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself 224

The operator  $\mathcal{A}^{(4)}$  is a contraction with respect to the metric 225

$$d \left( ((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)} t} \right\}$$

Indeed if we denote

$$\text{Definition of } (\widehat{G_{27}}, \widehat{T_{27}}) : (\widehat{G_{27}}, \widehat{T_{27}}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$$

It results

$$\begin{aligned}
 |\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} ds_{(24)} + \\
 &\int_0^t \{(a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} + \\
 &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} + \\
 &G_{24}^{(2)} |(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)})| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\} ds_{(24)}
 \end{aligned}$$

Where  $s_{(24)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned}
 |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)}t} &\leq & 226 \\
 \frac{1}{(\widehat{M}_{24})^{(4)}} ((a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)}) &d \left( ((G_{27})^{(1)}, (T_{27})^{(1)}); (G_{27})^{(2)}, (T_{27})^{(2)} \right)
 \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 16:** The fact that we supposed  $(a''_{24})^{(4)}$  and  $(b''_{24})^{(4)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$  and  $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$  respectively of  $\mathbb{R}_+$ . 227

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(4)}$  and  $(b''_i)^{(4)}$ ,  $i = 24, 25, 26$  depend only on  $T_{25}$  and respectively on  $(G_{27})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 17:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  228

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$  and  $((\widehat{M}_{24})^{(4)})_3$ : 229

**Remark 18:** if  $G_{24}$  is bounded, the same property have also  $G_{25}$  and  $G_{26}$ . indeed if

$$G_{24} < (\widehat{M}_{24})^{(4)} \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)} G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)} ((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)} ((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If  $G_{25}$  or  $G_{26}$  is bounded, the same property follows for  $G_{24}$ ,  $G_{26}$  and  $G_{24}$ ,  $G_{25}$  respectively.

**Remark 19:** If  $G_{24}$  is bounded, from below, the same property holds for  $G_{25}$  and  $G_{26}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{25}$  is bounded from below. 230

**Remark 20:** If  $T_{24}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$  then  $T_{25} \rightarrow \infty$ . 231

**Definition of**  $(m)^{(4)}$  and  $\varepsilon_4$  :

Indeed let  $t_4$  be so that for  $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then  $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$  which leads to 232

$$T_{25} \geq \left( \frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$$

If we take  $t$  such that  $e^{-\varepsilon_4 t} = \frac{1}{2}$  it results

$$T_{25} \geq \left( \frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4}$$

By taking now  $\varepsilon_4$  sufficiently small one sees that  $T_{25}$  is unbounded.

The same property holds for  $T_{26}$  if  $\lim_{t \rightarrow \infty} (b''_{26})^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for  $G_{29}$ ,  $G_{30}$ ,  $T_{28}$ ,  $T_{29}$ ,  $T_{30}$

It is now sufficient to take  $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}$ ,  $\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$  and to choose 233

$(\hat{P}_{28})^{(5)}$  and  $(\hat{Q}_{28})^{(5)}$  large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[ (\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left( \frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{28})^{(5)}$$
234

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} \left[ ((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left( \frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)}$$
235

In order that the operator  $\mathcal{A}^{(5)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(5)}$  is a contraction with respect to the metric 236

$$d \left( ((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{28})^{(5)} t} \right\}$$

Indeed if we denote

**Definition of**  $(\widehat{G}_{31}), (\widehat{T}_{31}) : ((\widehat{G}_{31}), (\widehat{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$|\widetilde{G}_{28}^{(1)} - \widetilde{G}_i^{(2)}| \leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} e^{(\widehat{M}_{28})^{(5)}s_{(28)}} ds_{(28)} +$$

$$\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} +$$

$$(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} e^{(\widehat{M}_{28})^{(5)}s_{(28)}} +$$

$$G_{28}^{(2)} | (a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)}) | e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\} ds_{(28)}$$

Where  $s_{(28)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses on it follows

$$|(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)}t} \leq \tag{237}$$

$$\frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)})$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 21:** The fact that we supposed  $(a''_{28})^{(5)}$  and  $(b''_{28})^{(5)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$  and  $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$  respectively of  $\mathbb{R}_+$ . 238

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(5)}$  and  $(b''_i)^{(5)}$ ,  $i = 28, 29, 30$  depend only on  $T_{29}$  and respectively on  $(G_{31})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 22:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  239

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$  and  $((\widehat{M}_{28})^{(5)})_3 :$  240

**Remark 23:** if  $G_{28}$  is bounded, the same property have also  $G_{29}$  and  $G_{30}$ . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)} G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If  $G_{29}$  or  $G_{30}$  is bounded, the same property follows for  $G_{28}$ ,  $G_{30}$  and  $G_{28}$ ,  $G_{29}$  respectively.

**Remark 24:** If  $G_{28}$  is bounded, from below, the same property holds for  $G_{29}$  and  $G_{30}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{29}$  is bounded from below. 241

**Remark 25:** If  $T_{28}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$  then  $T_{29} \rightarrow \infty$ . 242

**Definition of**  $(m)^{(5)}$  and  $\varepsilon_5$  :

Indeed let  $t_5$  be so that for  $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then  $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$  which leads to 243

$$T_{29} \geq \left( \frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$$

If we take  $t$  such that  $e^{-\varepsilon_5 t} = \frac{1}{2}$  it results

$$T_{29} \geq \left( \frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5}$$

By taking now  $\varepsilon_5$  sufficiently small one sees that  $T_{29}$  is unbounded.

The same property holds for  $T_{30}$  if  $\lim_{t \rightarrow \infty} (b'_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for  $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take  $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$  and to choose 244

$(\widehat{P}_{32})^{(6)}$  and  $(\widehat{Q}_{32})^{(6)}$  large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[ (\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[ ((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator  $\mathcal{A}^{(6)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(6)}$  is a contraction with respect to the metric 247

$$d \left( ((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)} t} \right\}$$

Indeed if we denote

**Definition of**  $(\widehat{G}_{35}), (\widehat{T}_{35}) : ( (\widehat{G}_{35}), (\widehat{T}_{35}) ) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$|\widehat{G}_{32}^{(1)} - \widehat{G}_i^{(2)}| \leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} e^{(\widehat{M}_{32})^{(6)}s_{(32)}} ds_{(32)} +$$

$$\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} +$$

$$(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} e^{(\widehat{M}_{32})^{(6)}s_{(32)}} +$$

$$G_{32}^{(2)} | (a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)}) | e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} e^{(\widehat{M}_{32})^{(6)}s_{(32)}} \} ds_{(32)}$$

Where  $s_{(32)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$|(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)}t} \leq \tag{248}$$

$$\frac{1}{(\widehat{M}_{32})^{(6)}} \left( (a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)} \right) d \left( ((G_{35})^{(1)}, (T_{35})^{(1)}); ((G_{35})^{(2)}, (T_{35})^{(2)}) \right)$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 26:** The fact that we supposed  $(a''_{32})^{(6)}$  and  $(b''_{32})^{(6)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$  and  $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$  respectively of  $\mathbb{R}_+$ . 249

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(6)}$  and  $(b''_i)^{(6)}, i = 32,33,34$  depend only on  $T_{33}$  and respectively on  $(G_{35})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 27:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  250

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(6)} - (a''_i)^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \} ds_{(32)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$  and  $((\widehat{M}_{32})^{(6)})_3 :$  251

**Remark 28:** if  $G_{32}$  is bounded, the same property have also  $G_{33}$  and  $G_{34}$ . indeed if

$$G_{32} < (\widehat{M}_{32})^{(6)} \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If  $G_{33}$  or  $G_{34}$  is bounded, the same property follows for  $G_{32}$ ,  $G_{34}$  and  $G_{32}$ ,  $G_{33}$  respectively.

**Remark 29:** If  $G_{32}$  is bounded, from below, the same property holds for  $G_{33}$  and  $G_{34}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{33}$  is bounded from below. 252

**Remark 30:** If  $T_{32}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b'_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$  then  $T_{33} \rightarrow \infty$ . 253

**Definition of**  $(m)^{(6)}$  and  $\varepsilon_6$  :

Indeed let  $t_6$  be so that for  $t > t_6$

$$(b_{33})^{(6)} - (b'_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then  $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$  which leads to 254

$$T_{33} \geq \left( \frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$$

If we take  $t$  such that  $e^{-\varepsilon_6 t} = \frac{1}{2}$  it results

$$T_{33} \geq \left( \frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6}$$

By taking now  $\varepsilon_6$  sufficiently small one sees that  $T_{33}$  is unbounded.

The same property holds for  $T_{34}$  if  $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for  $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$  255

It is now sufficient to take  $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$  and to choose

$(\widehat{P}_{36})^{(7)}$  and  $(\widehat{Q}_{36})^{(7)}$  large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[ (\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)}$$

256

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[ ((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)}$$

257

In order that the operator  $\mathcal{A}^{(7)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(7)}$  is a contraction with respect to the metric 258

$$d \left( ((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

**Definition of**  $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : ((\widetilde{G}_{39}), (\widetilde{T}_{39})) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\widetilde{G}_{36}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)}s_{(36)}} e^{(\widetilde{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{(a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)}s_{(36)}} e^{-(\widetilde{M}_{36})^{(7)}s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\widetilde{M}_{36})^{(7)}s_{(36)}} e^{(\widetilde{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} |(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)})| e^{-(\widetilde{M}_{36})^{(7)}s_{(36)}} e^{(\widetilde{M}_{36})^{(7)}s_{(36)}}\} ds_{(36)} \end{aligned}$$

Where  $s_{(36)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\widetilde{M}_{36})^{(7)}t} &\leq \tag{259} \\ \frac{1}{(\widetilde{M}_{36})^{(7)}} ((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)}) d &(((G_{39})^{(1)}, (T_{39})^{(1)}); (G_{39})^{(2)}, (T_{39})^{(2)}) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 31:** The fact that we supposed  $(a''_{36})^{(7)}$  and  $(b''_{36})^{(7)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)}t}$  and  $(\widehat{Q}_{36})^{(7)} e^{(\widetilde{M}_{36})^{(7)}t}$  respectively of  $\mathbb{R}_+$ . 260

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(7)}$  and  $(b''_i)^{(7)}, i = 36, 37, 38$  depend only on  $T_{37}$  and respectively on  $(G_{39})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 32:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  261

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widetilde{M}_{36})^{(7)})_1, ((\widetilde{M}_{36})^{(7)})_2$  and  $((\widetilde{M}_{36})^{(7)})_3$  : 262

**Remark 33:** if  $G_{36}$  is bounded, the same property have also  $G_{37}$  and  $G_{38}$ . indeed if

$$G_{36} < (\widetilde{M}_{36})^{(7)} \text{ it follows } \frac{dG_{37}}{dt} \leq ((\widetilde{M}_{36})^{(7)})_1 - (a'_{37})^{(7)} G_{37} \text{ and by integrating}$$

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If  $G_{37}$  or  $G_{38}$  is bounded, the same property follows for  $G_{36}$  ,  $G_{38}$  and  $G_{36}$  ,  $G_{37}$  respectively.

**Remark 34:** If  $G_{36}$  is bounded, from below, the same property holds for  $G_{37}$  and  $G_{38}$  . The proof is analogous with the preceding one. An analogous property is true if  $G_{37}$  is bounded from below. 263

**Remark 35:** If  $T_{36}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b'_{37})^{(7)}$  then  $T_{37} \rightarrow \infty$ . 264

**Definition of**  $(m)^{(7)}$  and  $\varepsilon_7$  :

Indeed let  $t_7$  be so that for  $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then  $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$  which leads to 265

$$T_{37} \geq \left( \frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$$

If we take  $t$  such that  $e^{-\varepsilon_7 t} = \frac{1}{2}$  it results

$$T_{37} \geq \left( \frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7}$$

By taking now  $\varepsilon_7$  sufficiently small one sees that  $T_{37}$  is unbounded.

The same property holds for  $T_{38}$  if  $\lim_{t \rightarrow \infty} (b_{38}'')^{(7)}((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take  $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$  and to choose  $(\widehat{P}_{40})^{(8)}$  and  $(\widehat{Q}_{40})^{(8)}$  large to have 266

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[ (\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)}$$
267

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[ ((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)}$$
268

In order that the operator  $\mathcal{A}^{(8)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(8)}$  is a contraction with respect to the metric

$$d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right), \left((G_{43})^{(2)}, (T_{43})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \tag{269}$$

Indeed if we denote 270

**Definition of**  $(\widetilde{G}_{43}), (\widetilde{T}_{43})$  :  $(\widetilde{G}_{43}), (\widetilde{T}_{43}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\widetilde{G}_{40}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{ (a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} | (a'_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)}) | e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} \} ds_{(40)} \end{aligned}$$

Where  $s_{(40)}$  represents integrand that is integrated over the interval  $[0, t]$  272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} \left( (a_{40})^{(8)} + (a'_{40})^{(8)} + (\bar{A}_{40})^{(8)} + (\bar{P}_{40})^{(8)} (\bar{k}_{40})^{(8)} \right) &d\left(\left((G_{43})^{(1)}, (T_{43})^{(1)}\right); (G_{43})^{(2)}, (T_{43})^{(2)}\right) \end{aligned} \tag{273}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 36:** The fact that we supposed  $(a''_{40})^{(8)}$  and  $(b''_{40})^{(8)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\bar{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$  and  $(\bar{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$  respectively of  $\mathbb{R}_+$ . 274

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a'_i)^{(8)}$  and  $(b'_i)^{(8)}$ ,  $i = 40, 41, 42$  depend only on  $T_{41}$  and respectively on  $(G_{43})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 37** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  275

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[ - \int_0^t \{ (a'_i)^{(8)} - (a''_i)^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \} ds_{(40)} \right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0 \end{aligned}$$

**Definition of**  $(\widetilde{M}_{40})^{(8)}_1, (\widetilde{M}_{40})^{(8)}_2$  and  $(\widetilde{M}_{40})^{(8)}_3$  : 276

**Remark 38:** if  $G_{40}$  is bounded, the same property have also  $G_{41}$  and  $G_{42}$  . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$  it follows  $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$  and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If  $G_{41}$  or  $G_{42}$  is bounded, the same property follows for  $G_{40}$  ,  $G_{42}$  and  $G_{40}$  ,  $G_{41}$  respectively.

**Remark 39:** If  $G_{40}$  is bounded, from below, the same property holds for  $G_{41}$  and  $G_{42}$  . The proof is 277  
 analogous with the preceding one. An analogous property is true if  $G_{41}$  is bounded from below.

**Remark 40:** If  $T_{40}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b'_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$  then  $T_{41} \rightarrow \infty$ . 278

**Definition of**  $(m)^{(8)}$  and  $\varepsilon_8$  :

Indeed let  $t_8$  be so that for  $t > t_8$

$$(b_{41})^{(8)} - (b'_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then  $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$  which leads to 279

$$T_{41} \geq \left( \frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$$

If we take  $t$  such that  $e^{-\varepsilon_8 t} = \frac{1}{2}$  it results

$$T_{41} \geq \left( \frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8}$$

By taking now  $\varepsilon_8$  sufficiently small one sees that  $T_{41}$  is unbounded.

The same property holds for  $T_{42}$  if  $\lim_{t \rightarrow \infty} ((b''_{42})^{(8)}((G_{43})(t), t(t), t)) = (b'_{42})^{(8)}$

It is now sufficient to take  $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$  ,  $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$  and to choose  $(\widehat{P}_{44})^{(9)}$  and  $(\widehat{Q}_{44})^{(9)}$  large to have 279  
 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[ (\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[ ((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator  $\mathcal{A}^{(9)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying 39,35,36 into itself

The operator  $\mathcal{A}^{(9)}$  is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

**Definition of**  $(\widehat{G_{47}}, \widehat{T_{47}}) : (\widehat{G_{47}}, \widehat{T_{47}}) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ G_{44}^{(2)} |(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a''_{44})^{(9)} (T_{45}^{(2)}, s_{(44)})| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)} \end{aligned}$$

Where  $s_{(44)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\bar{M}_{44})^{(9)}} &\left( (a_{44})^{(9)} + (a'_{44})^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{K}_{44})^{(9)} \right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis (39,35,36) the result follows

**Remark 41:** The fact that we supposed  $(a''_{44})^{(9)}$  and  $(b''_{44})^{(9)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$  and  $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(9)}$  and  $(b''_i)^{(9)}, i = 44,45,46$  depend only on  $T_{45}$  and respectively on  $(G_{47})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 42:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 99 to 44 it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0 \end{aligned}$$

**Definition of**  $(\bar{M}_{44})^{(9)}_1, (\bar{M}_{44})^{(9)}_2$  and  $(\bar{M}_{44})^{(9)}_3$  :

**Remark 43:** if  $G_{44}$  is bounded, the same property have also  $G_{45}$  and  $G_{46}$ . indeed if

$$G_{44} < (\bar{M}_{44})^{(9)} \text{ it follows } \frac{dG_{45}}{dt} \leq ((\bar{M}_{44})^{(9)}_1 - (a'_{45})^{(9)}) G_{45} \text{ and by integrating}$$

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If  $G_{45}$  or  $G_{46}$  is bounded, the same property follows for  $G_{44}$  ,  $G_{46}$  and  $G_{44}$  ,  $G_{45}$  respectively.

**Remark 44:** If  $G_{44}$  is bounded, from below, the same property holds for  $G_{45}$  and  $G_{46}$  . The proof is analogous with the preceding one. An analogous property is true if  $G_{45}$  is bounded from below.

**Remark 45:** If  $T_{44}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b'_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$  then  $T_{45} \rightarrow \infty$ .

**Definition of**  $(m)^{(9)}$  and  $\varepsilon_9$  :

Indeed let  $t_9$  be so that for  $t > t_9$

$$(b_{45})^{(9)} - (b'_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then  $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$  which leads to

$$T_{45} \geq \left( \frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left( \frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for  $T_{46}$  if  $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

**Behavior of the solutions of equation**

280

**Theorem** If we denote and define

**Definition of**  $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$  :

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$  four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

**Definition of**  $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$  :

281

By  $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$  and respectively  $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$  the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

**Definition of**  $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$  :

282

By  $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$  and respectively  $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$  the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

**Definition of**  $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$  :-

283

If we define  $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$  by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

284

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } \boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where  $(p_i)^{(1)}$  is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left( \frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[ e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \right) \leq G_{15}(t) \leq$$

286

$$\frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[ e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}}$$

287

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}$$

288

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[ e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq$$

289

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[ e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

**Definition of**  $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$ :-

290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

**Behavior of the solutions of equation**

291

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$  :

292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$  four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$$

293

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$$

294

**Definition of**  $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$  :

295

By  $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$  and respectively  $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$  the roots

296

of the equations  $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

297

and  $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$  and

298

**Definition of**  $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$  :

299

By  $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$  and respectively  $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$  the

300

roots of the equations  $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$

301

and  $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$

302

**Definition of**  $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$  :-

303

If we define  $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$  by

304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)}$$

305

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$$

306

and 
$$(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)}$$

307

and analogously

308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and 
$$\boxed{(u_0)^{(2)} = \begin{matrix} T_{16}^0 \\ T_{17}^0 \end{matrix}}$$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)}$  309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$  is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$$
 311

$$\left( \frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[ e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq$$
 312

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[ e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$$
 313

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$$
 314

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[ e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq$$
 315

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[ e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

**Definition of**  $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$ :- 316

Where  $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$  317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$$
 318

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

**Behavior of the solutions** 319

**Theorem 3:** If we denote and define

**Definition of**  $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$  :

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$  four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

**Definition of**  $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$  : 320

By  $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$  and respectively  $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$  the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By  $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$  and respectively  $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$  the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

**Definition of**  $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$  :- 321

If we define  $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$  by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } \boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously 322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } \boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$  is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left( \frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[ e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \leq \quad 324$$

$$\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[ e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})} \left[ e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})} \left[ e^{((R_1)^{(3)}+(r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t} \quad 328$$

**Definition of**  $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$ :-

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

**Behavior of the solutions of equation**

**Theorem:** If we denote and define

**Definition of**  $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$  :

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$  four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

**Definition of**  $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$  : 329

By  $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$  and respectively  $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$  the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$  : 330

By  $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$  and respectively  $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$  the

roots of the equations  $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and  $(b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

**Definition of**  $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$  :-

If we define  $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$  by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } \boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where  $(p_i)^{(4)}$  is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left( \frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \right) \left[ e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[ e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[ e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[ e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

**Definition of**  $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$ :-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

**Behavior of the solutions of equation**

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$  :

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$  four constants satisfying

$$\begin{aligned}
 -(\sigma_2)^{(5)} &\leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)} \\
 -(\tau_2)^{(5)} &\leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}
 \end{aligned}$$

**Definition of**  $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$  : 339

By  $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$  and respectively  $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$  the roots of the equations  
 $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$   
 and  $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$  and

**Definition of**  $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$  : 340

By  $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$  and respectively  $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$  the roots of the equations  $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$   
 and  $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

**Definition of**  $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$  :-

If we define  $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$  by

$$\begin{aligned}
 (m_2)^{(5)} &= (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)} \\
 (m_2)^{(5)} &= (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)}, \\
 \text{and } (v_0)^{(5)} &= \frac{G_{28}^0}{G_{29}^0}
 \end{aligned}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$\begin{aligned}
 (\mu_2)^{(5)} &= (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)} \\
 (\mu_2)^{(5)} &= (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}, \\
 \text{and } (u_0)^{(5)} &= \frac{T_{28}^0}{T_{29}^0}
 \end{aligned}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where  $(p_i)^{(5)}$  is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(s_1)^{(5)}t} \quad 343$$

$$\left( \frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((s_1)^{(5)} - (p_{28})^{(5)} - (s_2)^{(5)})} \right) \left[ e^{((s_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(s_2)^{(5)}t} \right] + G_{30}^0 e^{-(s_2)^{(5)}t} \leq G_{30}(t) \leq \quad 344$$

$$\frac{(a_{30})^{(5)}G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})} \left[ e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})} \left[ e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[ e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

**Definition of**  $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$ :- 348

Where  $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

**Behavior of the solutions of equation** 349

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$  :

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$  four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

**Definition of**  $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$  : 350

By  $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$  and respectively  $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$  the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$  : 351

By  $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$  and respectively  $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$  the

roots of the equations  $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$$

**Definition of**  $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$  :-

If we define  $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$  by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

and  $\boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

and  $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where  $(p_i)^{(6)}$  is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left( \frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \right) \left[ e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \leq \quad 355$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[ e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[ e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[ e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

**Definition of**  $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$ :- 359

$$\text{Where } (S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

**Behavior of the solutions of equation**

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$  :

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$  four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

**Definition of**  $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$  :

361

By  $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$  and respectively  $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$  the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

and  $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$  and

**Definition of**  $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$  :

362

By  $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$  and respectively  $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$  the

roots of the equations  $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and  $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

**Definition of**  $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$  :-

If we define  $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$  by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } (u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where  $(p_i)^{(7)}$  is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \tag{365}$$

$$\left( \frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[ e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \tag{366}$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[ e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \tag{367}$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \tag{368}$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[ e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \tag{369}$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[ e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

**Definition of**  $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$ :- 370

Where  $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

**Behavior of the solutions of equation** 371

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$  :

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$  four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

**Definition of**  $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$  : 372

By  $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$  and respectively  $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$  the roots of the equations  $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$  and  $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$  and

**Definition of**  $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$  :

By  $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$  and respectively  $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$  the

roots of the equations  $(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and  $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

**Definition of**  $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$  :-

If we define  $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$  by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where  $(p_i)^{(8)}$  is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t}$$

376

$$\left( \frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[ e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \leq$$

377

$$\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})} [e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}] + G_{42}^0 e^{-(a'_{42})^{(8)}t}$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})} [e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})} [e^{((R_1)^{(8)}+(r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t}] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

**Definition of**  $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$ :- 381

Where  $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

**Behavior of the solutions of equation 37 to 92** 382

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$  :

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$  four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

**Definition of**  $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$  :

By  $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$  and respectively  $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$  the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$  :

By  $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$  and respectively  $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$  the

$$\text{roots of the equations } (a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$$

**Definition of**  $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$  :-

If we define  $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$  by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$  where  $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$  are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where  $(p_i)^{(9)}$  is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)} ((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[ e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)} ((S_1)^{(9)} - (a'_{46})^{(9)})} \left[ e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)} ((R_1)^{(9)} - (b'_{46})^{(9)})} \left[ e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)} ((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[ e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

**Definition of**  $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$ :-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

**Proof:** From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left( (a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

**Definition of**  $v^{(1)}$  :- 
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left( (a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left( (a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$  :-

For  $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}(v_1)^{(1)} - (v_0)^{(1)}]t}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}(v_1)^{(1)} - (v_0)^{(1)}]t}}, \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

it follows  $(v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}}, \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}$$

From which we deduce  $(v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$

If  $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$  we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}(v_1)^{(1)} - (v_2)^{(1)}]t}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}(v_1)^{(1)} - (v_2)^{(1)}]t}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}} \leq (\bar{v}_1)^{(1)}$$

If  $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$ , we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(1)}(t)$  :-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(1)}(t)$  :-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If  $(a''_{13})^{(1)} = (a''_{14})^{(1)}$ , then  $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$  and in this case  $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$  if in addition  $(v_0)^{(1)} = (v_1)^{(1)}$  then  $v^{(1)}(t) = (v_0)^{(1)}$  and as a consequence  $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$  this also defines  $(v_0)^{(1)}$  for the special case

Analogously if  $(b''_{13})^{(1)} = (b''_{14})^{(1)}$ , then  $(\tau_1)^{(1)} = (\tau_2)^{(1)}$  and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$  if in addition  $(u_0)^{(1)} = (u_1)^{(1)}$  then  $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$  This is an important consequence of the relation between  $(v_1)^{(1)}$  and  $(\bar{v}_1)^{(1)}$ , and definition of  $(u_0)^{(1)}$ .

**Proof:** From global equations we obtain

387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left( (a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a'_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

**Definition of**  $v^{(2)}$  :-

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

388

It follows

389

$$- \left( (a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left( (a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains

390

**Definition of**  $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$  :-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows  $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} (\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}] t}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} (\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}] t}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce  $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$  392

If  $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$  we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)} (v_2)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)} ((v_1)^{(2)} - (v_2)^{(2)}) t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (\bar{v}_1)^{(2)}$$

If  $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$  , we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)} (\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)} ((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}) t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(2)}(t)$  :- 395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

**Definition of**  $u^{(2)}(t)$  :-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :** 397

If  $(a''_{16})^{(2)} = (a''_{17})^{(2)}$  , then  $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$  and in this case  $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$  if in addition  $(v_0)^{(2)} = (v_1)^{(2)}$  then  $v^{(2)}(t) = (v_0)^{(2)}$  and as a consequence  $G_{16}(t) = (v_0)^{(2)} G_{17}(t)$

Analogously if  $(b''_{16})^{(2)} = (b''_{17})^{(2)}$  , then  $(\tau_1)^{(2)} = (\tau_2)^{(2)}$  and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$  if in addition  $(u_0)^{(2)} = (u_1)^{(2)}$  then  $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$  This is an important consequence of the relation between  $(v_1)^{(2)}$  and  $(\bar{v}_1)^{(2)}$

**Proof :** From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left( (a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

**Definition of**  $v^{(3)}$  :- 
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$-\left( (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq -\left( (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

For  $0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

it follows  $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

**Definition of**  $(\bar{v}_1)^{(3)}$  :-

From which we deduce  $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If  $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$  we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If  $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$  , we obtain 403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(3)}(t)$  :-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(3)}(t)$  :-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{20})^{(3)} = (a''_{21})^{(3)}$ , then  $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$  and in this case  $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$  if in addition  $(v_0)^{(3)} = (v_1)^{(3)}$  then  $v^{(3)}(t) = (v_0)^{(3)}$  and as a consequence  $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if  $(b''_{20})^{(3)} = (b''_{21})^{(3)}$ , then  $(\tau_1)^{(3)} = (\tau_2)^{(3)}$  and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$  if in addition  $(u_0)^{(3)} = (u_1)^{(3)}$  then  $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$  This is an important consequence of the relation between  $(v_1)^{(3)}$  and  $(\bar{v}_1)^{(3)}$

**Proof:** From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left( (a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

**Definition of**  $v^{(4)}$  :-  $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$- \left( (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left( (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$  :-

For  $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}$$

it follows  $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}$$

From which we deduce  $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If  $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$  we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (\bar{C})^{(4)} (v_2)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (\bar{v}_1)^{(4)}$$

If  $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$ , we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(4)}(t)$  :-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(4)}(t)$  :-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}$ , then  $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$  and in this case  $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$  if in addition  $(v_0)^{(4)} = (v_1)^{(4)}$  then  $v^{(4)}(t) = (v_0)^{(4)}$  and as a consequence  $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$  **this also defines**  $(v_0)^{(4)}$  **for the special case .**

Analogously if  $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}$ , then  $(\tau_1)^{(4)} = (\tau_2)^{(4)}$  and then  $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$  if in addition  $(u_0)^{(4)} = (u_1)^{(4)}$  then  $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$  This is an important consequence of the relation between  $(v_1)^{(4)}$  and  $(\bar{v}_1)^{(4)}$ , **and definition of**  $(u_0)^{(4)}$ .

408

**Proof:** From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left( (a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t) \right) - (a_{29}'')^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

**Definition of**  $v^{(5)}$  :-  $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left( (a_{29})^{(5)} (v^{(5)})^2 + (\sigma_2)^{(5)} v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left( (a_{29})^{(5)} (v^{(5)})^2 + (\sigma_1)^{(5)} v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$  :-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}$$

it follows  $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}$$

From which we deduce  $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If  $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$  we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If  $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$  , we obtain

411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(5)}(t)$  :-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)} , \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(5)}(t)$  :-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)} , \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{28})^{(5)} = (a''_{29})^{(5)}$ , then  $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$  and in this case  $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$  if in addition  $(v_0)^{(5)} = (v_5)^{(5)}$  then  $v^{(5)}(t) = (v_0)^{(5)}$  and as a consequence  $G_{28}(t) = (v_0)^{(5)}G_{29}(t)$  **this also defines  $(v_0)^{(5)}$  for the special case .**

Analogously if  $(b''_{28})^{(5)} = (b''_{29})^{(5)}$ , then  $(\tau_1)^{(5)} = (\tau_2)^{(5)}$  and then  $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$  if in addition  $(u_0)^{(5)} = (u_1)^{(5)}$  then  $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$  This is an important consequence of the relation between  $(v_1)^{(5)}$  and  $(\bar{v}_1)^{(5)}$ , **and definition of  $(u_0)^{(5)}$ .**

**Proof:** From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left( (a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

**Definition of  $v^{(6)}$  :-** 
$$v^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

It follows

$$- \left( (a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left( (a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

**Definition of  $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$  :-**

For  $0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows  $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce  $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If  $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$  we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If  $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$ , we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of  $v^{(6)}(t)$  :-**

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(6)}(t)$  :-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{32})^{(6)} = (a''_{33})^{(6)}$ , then  $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$  and in this case  $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$  if in addition  $(v_0)^{(6)} = (v_1)^{(6)}$  then  $v^{(6)}(t) = (v_0)^{(6)}$  and as a consequence  $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$  **this also defines  $(v_0)^{(6)}$  for the special case.**

Analogously if  $(b''_{32})^{(6)} = (b''_{33})^{(6)}$ , then  $(\tau_1)^{(6)} = (\tau_2)^{(6)}$  and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$  if in addition  $(u_0)^{(6)} = (u_1)^{(6)}$  then  $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$  This is an important consequence of the relation between  $(v_1)^{(6)}$  and  $(\bar{v}_1)^{(6)}$ , **and definition of  $(u_0)^{(6)}$ .**

416

**Proof :** From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left( (a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

**Definition of**  $v^{(7)}$  :-  $\boxed{v^{(7)} = \frac{a_{36}}{a_{37}}}$

It follows

$$- \left( (a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left( (a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$  :-

For  $0 < \boxed{(v_0)^{(7)} = \frac{a_{36}^0}{a_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows  $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce  $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If  $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$  we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (\bar{C})^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If  $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$ , we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(7)}(t)$  :-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(7)}(t)$  :- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a_{36}'')^{(7)} = (a_{37}'')^{(7)}$ , then  $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$  and in this case  $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$  if in addition  $(v_0)^{(7)} = (v_1)^{(7)}$  then  $v^{(7)}(t) = (v_0)^{(7)}$  and as a consequence  $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$  **this also defines  $(v_0)^{(7)}$  for the special case.**

Analogously if  $(b_{36}'')^{(7)} = (b_{37}'')^{(7)}$ , then  $(\tau_1)^{(7)} = (\tau_2)^{(7)}$  and then  $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$  if in addition  $(u_0)^{(7)} = (u_1)^{(7)}$  then  $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$  This is an important consequence of the relation between  $(v_1)^{(7)}$  and  $(\bar{v}_1)^{(7)}$ , **and definition of  $(u_0)^{(7)}$ .**

**Proof :** From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left( (a_{40}')^{(8)} - (a_{41}')^{(8)} + (a_{40}'')^{(8)}(T_{41}, t) \right) - (a_{41}'')^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

**Definition of**  $v^{(8)}$  :- 
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$  :-

For  $0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}(v_1)^{(8)} - (v_0)^{(8)}]t}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}(v_1)^{(8)} - (v_0)^{(8)}]t}}, \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}$$

it follows  $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}}, \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}$$

From which we deduce  $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If  $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$  we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}(v_1)^{(8)} - (v_2)^{(8)}]t}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}(v_1)^{(8)} - (v_2)^{(8)}]t}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}} \leq (\bar{v}_1)^{(8)}$$

If  $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$ , we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(8)}(t)$  :-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(8)}(t)$  :-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{40})^{(8)} = (a''_{41})^{(8)}$ , then  $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$  and in this case  $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$  if in addition  $(v_0)^{(8)} = (v_1)^{(8)}$  then  $v^{(8)}(t) = (v_0)^{(8)}$  and as a consequence  $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$  **this also defines  $(v_0)^{(8)}$  for the special case .**

Analogously if  $(b''_{40})^{(8)} = (b''_{41})^{(8)}$ , then  $(\tau_1)^{(8)} = (\tau_2)^{(8)}$  and then  $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$  if in addition  $(u_0)^{(8)} = (u_1)^{(8)}$  then  $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$  This is an important consequence of the relation between  $(v_1)^{(8)}$  and  $(\bar{v}_1)^{(8)}$ , **and definition of  $(u_0)^{(8)}$ .**

**Proof :** From 99,20,44,22,23,44 we obtain

424  
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left( (a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

**Definition of  $v^{(9)}$  :-**  $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left( (a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left( (a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

**Definition of  $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$  :-**

For  $0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}$$

it follows  $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}$$

From which we deduce  $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If  $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$  we find like in the previous case,

$$(v_1)^{(9)} \leq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_2)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_2)^{(9)})t]}} \leq v^{(9)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} \leq (\bar{v}_1)^{(9)}$$

If  $0 < (v_1)^{(9)} \leq (\bar{v}_1)^{(9)} \leq \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$ , we obtain

$$(v_1)^{(9)} \leq v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} \leq (v_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(9)}(t)$  :-

$$(m_2)^{(9)} \leq v^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{v^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(9)}(t)$  :-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{44})^{(9)} = (a''_{45})^{(9)}$ , then  $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$  and in this case  $(v_1)^{(9)} = (\bar{v}_1)^{(9)}$  if in addition  $(v_0)^{(9)} = (v_1)^{(9)}$  then  $v^{(9)}(t) = (v_0)^{(9)}$  and as a consequence  $G_{44}(t) = (v_0)^{(9)}G_{45}(t)$  **this also defines  $(v_0)^{(9)}$  for the special case.**

Analogously if  $(b''_{44})^{(9)} = (b''_{45})^{(9)}$ , then  $(\tau_1)^{(9)} = (\tau_2)^{(9)}$  and then  $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$  if in addition  $(u_0)^{(9)} = (u_1)^{(9)}$  then  $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$  This is an important consequence of the relation between  $(v_1)^{(9)}$  and  $(\bar{v}_1)^{(9)}$ , **and definition of  $(u_0)^{(9)}$ .**

We can prove the following

425

**Theorem :** If  $(a'_i)^{(1)}$  and  $(b'_i)^{(1)}$  are independent on  $t$ , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with  $(p_{13})^{(1)}, (r_{14})^{(1)}$  as defined by equation are satisfied, then the system

**Theorem :** If  $(a'_i)^{(2)}$  and  $(b'_i)^{(2)}$  are independent on  $t$ , and the conditions with the notations

426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0, \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with  $(p_{16})^{(2)}, (r_{17})^{(2)}$  as defined by equation are satisfied, then the system

**Theorem :** If  $(a_i'')^{(3)}$  and  $(b_i'')^{(3)}$  are independent on  $t$ , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with  $(p_{20})^{(3)}, (r_{21})^{(3)}$  as defined by equation are satisfied, then the system

We can prove the following 432

**Theorem :** If  $(a_i'')^{(4)}$  and  $(b_i'')^{(4)}$  are independent on  $t$ , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with  $(p_{24})^{(4)}, (r_{25})^{(4)}$  as defined by equation are satisfied, then the system

**Theorem :** If  $(a_i'')^{(5)}$  and  $(b_i'')^{(5)}$  are independent on  $t$ , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with  $(p_{28})^{(5)}, (r_{29})^{(5)}$  as defined by equation are satisfied, then the system

**Theorem** If  $(a_i'')^{(6)}$  and  $(b_i'')^{(6)}$  are independent on  $t$ , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with  $(p_{32})^{(6)}, (r_{33})^{(6)}$  as defined by equation are satisfied , then the system

**Theorem :** If  $(a''_i)^{(7)}$  and  $(b''_i)^{(7)}$  are independent on  $t$  , and the conditions with the notations

435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with  $(p_{36})^{(7)}, (r_{37})^{(7)}$  as defined by equation are satisfied , then the system

**Theorem :** If  $(a''_i)^{(8)}$  and  $(b''_i)^{(8)}$  are independent on  $t$  , and the conditions with the notations

436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with  $(p_{40})^{(8)}, (r_{41})^{(8)}$  as defined by equation are satisfied , then the system

**Theorem :** If  $(a''_i)^{(9)}$  and  $(b''_i)^{(9)}$  are independent on  $t$  , and the conditions (with the notations 45,46,27,28)

436  
A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0 ,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with  $(p_{44})^{(9)}, (r_{45})^{(9)}$  as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0$$

437

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0$$

438

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

**Proof:** 485

(a) Indeed the first two equations have a nontrivial solution  $G_{13}, G_{14}$  if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

**Proof:** 486

(a) Indeed the first two equations have a nontrivial solution  $G_{16}, G_{17}$  if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

**Proof:** 487

(a) Indeed the first two equations have a nontrivial solution  $G_{20}, G_{21}$  if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

**Proof:** 488

(a) Indeed the first two equations have a nontrivial solution  $G_{24}, G_{25}$  if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

**Proof:**

489

(a) Indeed the first two equations have a nontrivial solution  $G_{28}, G_{29}$  if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

**Proof:**

490

(a) Indeed the first two equations have a nontrivial solution  $G_{32}, G_{33}$  if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

**Proof:**

491

(a) Indeed the first two equations have a nontrivial solution  $G_{36}, G_{37}$  if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

**Proof:**

492

(a) Indeed the first two equations have a nontrivial solution  $G_{40}, G_{41}$  if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

**Proof:**

492

A

(a) Indeed the first two equations have a nontrivial solution  $G_{44}, G_{45}$  if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

**Definition and uniqueness of  $T_{14}^*$  :-**

493

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(1)}(T_{14})$  being increasing, it follows that there exists a unique  $T_{14}^*$  for which  $f(T_{14}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

**Definition and uniqueness of  $T_{17}^*$  :-**

494

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(2)}(T_{17})$  being increasing, it follows that there exists a unique  $T_{17}^*$  for which  $f(T_{17}^*) = 0$ . With this value, we obtain from the three first

equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

**Definition and uniqueness of  $T_{21}^*$  :-** 496

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(1)}(T_{21})$  being increasing, it follows that there exists a unique  $T_{21}^*$  for which  $f(T_{21}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

**Definition and uniqueness of  $T_{25}^*$  :-** 497

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(4)}(T_{25})$  being increasing, it follows that there exists a unique  $T_{25}^*$  for which  $f(T_{25}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

**Definition and uniqueness of  $T_{29}^*$  :-** 498

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(5)}(T_{29})$  being increasing, it follows that there exists a unique  $T_{29}^*$  for which  $f(T_{29}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

**Definition and uniqueness of  $T_{33}^*$  :-** 499

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(6)}(T_{33})$  being increasing, it follows that there exists a unique  $T_{33}^*$  for which  $f(T_{33}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

**Definition and uniqueness of  $T_{37}^*$  :-** 500

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(7)}(T_{37})$  being increasing, it follows that there exists a unique  $T_{37}^*$  for which  $f(T_{37}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

**Definition and uniqueness of  $T_{41}^*$  :-** 501

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(8)}(T_{41})$  being increasing, it follows that there exists a unique  $T_{41}^*$  for which  $f(T_{41}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^*)]} , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^*)]}$$

**Definition and uniqueness of  $T_{45}^*$  :-**

501  
A

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(9)}(T_{45})$  being increasing, it follows that there exists a unique  $T_{45}^*$  for which  $f(T_{45}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^*)]} , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions  $G_{13}, G_{14}$  if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in  $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{14}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{14}^*$  such that  $\varphi(G^*) = 0$

By the same argument, the equations admit solutions  $G_{16}, G_{17}$  if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in  $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{17}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{17}^*$  such that  $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions  $G_{20}, G_{21}$  if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in  $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{21}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{21}^*$  such that  $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions  $G_{24}, G_{25}$  if

506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in  $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{25}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{25}^*$  such that  $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions  $G_{28}, G_{29}$  if 507  
 $\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in  $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{29}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{29}^*$  such that  $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions  $G_{32}, G_{33}$  if 508  
 $\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in  $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{33}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{33}^*$  such that  $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions  $G_{36}, G_{37}$  if 509  
 $\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in  $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{37}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{37}^*$  such that  $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions  $G_{40}, G_{41}$  if 510  
 $\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$

$$[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in  $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{41}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{41}^*$  such that  $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions  $G_{44}, G_{45}$  if  
 $\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in  $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{45}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{45}^*$  such that  $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

$G_{14}^*$  given by  $\varphi(G^*) = 0$ ,  $T_{14}^*$  given by  $f(T_{14}^*) = 0$  and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^*)]} , T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

$G_{17}^*$  given by  $\varphi((G_{19})^*) = 0$ ,  $T_{17}^*$  given by  $f(T_{17}^*) = 0$  and

512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} , G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]}$$

513

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^*)]} , T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^*)]}$$

514

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

515

$G_{21}^*$  given by  $\varphi((G_{23})^*) = 0$ ,  $T_{21}^*$  given by  $f(T_{21}^*) = 0$  and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} , G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

516

$G_{25}^*$  given by  $\varphi(G_{27}) = 0$ ,  $T_{25}^*$  given by  $f(T_{25}^*) = 0$  and

$$G_{24}^* = \frac{(a_{24})^{(4)}G_{25}^*}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} , G_{26}^* = \frac{(a_{26})^{(4)}G_{25}^*}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)}T_{25}^*}{[(b'_{24})^{(4)}-(b''_{24})^{(4)}((G_{27})^*)]} , T_{26}^* = \frac{(b_{26})^{(4)}T_{25}^*}{[(b'_{26})^{(4)}-(b''_{26})^{(4)}((G_{27})^*)]}$$

517

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

518

$G_{29}^*$  given by  $\varphi((G_{31})^*) = 0$ ,  $T_{29}^*$  given by  $f(T_{29}^*) = 0$  and

$$G_{28}^* = \frac{(a_{28})^{(5)}G_{29}^*}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} , G_{30}^* = \frac{(a_{30})^{(5)}G_{29}^*}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)}T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)}T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \quad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

$G_{33}^*$  given by  $\varphi((G_{35})^*) = 0$  ,  $T_{33}^*$  given by  $f(T_{33}^*) = 0$  and

$$G_{32}^* = \frac{(a_{32})^{(6)}G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)}G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)}T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)}T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

$G_{37}^*$  given by  $\varphi((G_{39})^*) = 0$  ,  $T_{37}^*$  given by  $f(T_{37}^*) = 0$  and

$$G_{36}^* = \frac{(a_{36})^{(7)}G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)}G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)}T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)}T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

$G_{41}^*$  given by  $\varphi((G_{43})^*) = 0$  ,  $T_{41}^*$  given by  $f(T_{41}^*) = 0$  and

$$G_{40}^* = \frac{(a_{40})^{(8)}G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)}G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)}T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)}T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99 523  
A

$G_{45}^*$  given by  $\varphi((G_{47})^*) = 0$  ,  $T_{45}^*$  given by  $f(T_{45}^*) = 0$  and

$$G_{44}^* = \frac{(a_{44})^{(9)}G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)}G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)}T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)}T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

**ASYMPTOTIC STABILITY ANALYSIS** 524

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a'_i)^{(1)}$  and  $(b'_i)^{(1)}$  belong to  $C^{(1)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

**Proof:** Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a'_{14})^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial(b'_i)^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})G_{13} + (a_{13})^{(1)}G_{14} - (q_{13})^{(1)}G_{13}^*T_{14} \quad 525$$

$$\frac{dG_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})G_{14} + (a_{14})^{(1)}G_{13} - (q_{14})^{(1)}G_{14}^*T_{14} \quad 526$$

$$\frac{dG_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})G_{15} + (a_{15})^{(1)}G_{14} - (q_{15})^{(1)}G_{15}^*T_{14} \quad 527$$

$$\frac{dT_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})T_{13} + (b_{13})^{(1)}T_{14} + \sum_{j=13}^{15} (s_{(13)(j)})T_{13}^*G_j \quad 528$$

$$\frac{dT_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})T_{14} + (b_{14})^{(1)}T_{13} + \sum_{j=13}^{15} (s_{(14)(j)})T_{14}^*G_j \quad 529$$

$$\frac{dT_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})T_{15} + (b_{15})^{(1)}T_{14} + \sum_{j=13}^{15} (s_{(15)(j)})T_{15}^*G_j \quad 530$$

**ASYMPTOTIC STABILITY ANALYSIS** 531

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a'_i)^{(2)}$  and  $(b'_i)^{(2)}$  belong to  $C^{(2)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

**Proof:** Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a'_{17})^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \quad \frac{\partial(b'_i)^{(2)}}{\partial G_j}(G_{19}^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \quad 534$$

$$\frac{dG_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \quad 535$$

$$\frac{dG_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \quad 536$$

$$\frac{dT_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(16)(j)})T_{16}^*G_j \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)})T_{17}^*G_j \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)})T_{18}^*G_j \quad 539$$

**ASYMPTOTIC STABILITY ANALYSIS** 540

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a'_i)^{(3)}$  and  $(b'_i)^{(3)}$  belong to  $C^{(3)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a''_i)^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b'_i)^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)})T_{20}^*G_j \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)})T_{21}^*G_j \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)})T_{22}^*G_j \quad 546$$

**ASYMPTOTIC STABILITY ANALYSIS** 547

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a'_i)^{(4)}$  and  $(b'_i)^{(4)}$  belong to  $C^{(4)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :- 548

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a''_i)^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b'_i)^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})G_{24} + (a_{24})^{(4)}G_{25} - (q_{24})^{(4)}G_{24}^*T_{25} \quad 549$$

$$\frac{dG_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})G_{25} + (a_{25})^{(4)}G_{24} - (q_{25})^{(4)}G_{25}^*T_{25} \quad 550$$

$$\frac{dG_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})G_{26} + (a_{26})^{(4)}G_{25} - (q_{26})^{(4)}G_{26}^*T_{25} \quad 551$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)})T_{24}^*G_j \quad 552$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)})T_{25}^*G_j \quad 553$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)})T_{26}^*G_j \quad 554$$

**ASYMPTOTIC STABILITY ANALYSIS** 555

**Theorem 5:** If the conditions of the previous theorem are satisfied and if the functions  $(a'_i)^{(5)}$  and  $(b''_i)^{(5)}$  belong to  $C^{(5)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :- 556

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial(b''_i)^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})G_{28} + (a_{28})^{(5)}G_{29} - (q_{28})^{(5)}G_{28}^*T_{29} \quad 557$$

$$\frac{dG_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})G_{29} + (a_{29})^{(5)}G_{28} - (q_{29})^{(5)}G_{29}^*T_{29} \quad 558$$

$$\frac{dG_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})G_{30} + (a_{30})^{(5)}G_{29} - (q_{30})^{(5)}G_{30}^*T_{29} \quad 559$$

$$\frac{dT_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})T_{28} + (b_{28})^{(5)}T_{29} + \sum_{j=28}^{30} (s_{(28)(j)})T_{28}^*G_j \quad 560$$

$$\frac{dT_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})T_{29} + (b_{29})^{(5)}T_{28} + \sum_{j=28}^{30} (s_{(29)(j)})T_{29}^*G_j \quad 561$$

$$\frac{dT_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})T_{30} + (b_{30})^{(5)}T_{29} + \sum_{j=28}^{30} (s_{(30)(j)})T_{30}^*G_j \quad 562$$

**ASYMPTOTIC STABILITY ANALYSIS** 563

**Theorem 6:** If the conditions of the previous theorem are satisfied and if the functions  $(a'_i)^{(6)}$  and  $(b''_i)^{(6)}$  belong to  $C^{(6)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :- 564

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(G_{35}^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \tag{565}$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \tag{566}$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \tag{567}$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(32)(j)})T_{32}^*G_j \tag{568}$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34}(s_{(33)(j)})T_{33}^*G_j \tag{569}$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34}(s_{(34)(j)})T_{34}^*G_j \tag{570}$$

**ASYMPTOTIC STABILITY ANALYSIS** 571

**Theorem 7:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(7)}$  and  $(b_i'')^{(7)}$  belong to  $C^{(7)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :- 572

$$G_i = G_i^* + \mathbb{G}_i , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} , \frac{\partial(b_i'')^{(7)}}{\partial G_j}(G_{39}^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \tag{573}$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \tag{574}$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \tag{575}$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(36)(j)})T_{36}^*G_j \tag{576}$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38}(s_{(37)(j)})T_{37}^*G_j \tag{578}$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38}(s_{(38)(j)})T_{38}^*G_j \tag{579}$$

Obviously, these values represent an equilibrium solution

**ASYMPTOTIC STABILITY ANALYSIS**

**Theorem 8:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(8)}$  and  $(b_i'')^{(8)}$  belong to  $C^{(8)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial(b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \tag{581}$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \tag{582}$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \tag{583}$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)})T_{40}^*G_j \tag{584}$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)})T_{41}^*G_j \tag{585}$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)})T_{42}^*G_j \tag{586}$$

**ASYMPTOTIC STABILITY ANALYSIS** 586  
A

**Theorem 9:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(9)}$  and  $(b_i'')^{(9)}$  belong to  $C^{(9)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \tag{586  
B}$$

$$\frac{dG_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})G_{45} + (a_{45})^{(9)}G_{44} - (q_{45})^{(9)}G_{45}^*T_{45} \quad 586$$

C

$$\frac{dG_{46}}{dt} = -((b'_{46})^{(9)} + (p_{46})^{(9)})G_{46} + (a_{46})^{(9)}G_{45} - (q_{46})^{(9)}G_{46}^*T_{45} \quad 586$$

D

$$\frac{dT_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})T_{44} + (b_{44})^{(9)}T_{45} + \sum_{j=44}^{46} (s_{(44)(j)})T_{44}^*G_j \quad 586$$

E

$$\frac{dT_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})T_{45} + (b_{45})^{(9)}T_{44} + \sum_{j=44}^{46} (s_{(45)(j)})T_{45}^*G_j \quad 586$$

F

$$\frac{dT_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})T_{46} + (b_{46})^{(9)}T_{45} + \sum_{j=44}^{46} (s_{(46)(j)})T_{46}^*G_j \quad 586$$

G

**The characteristic equation of this system is**

587

$$\begin{aligned} &((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ &\left[ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ &\left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \right) \\ &+ \left( ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \right) \\ &\left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \right) \\ &\left( ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &\left( ((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right) \\ &+ \left( ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)}G_{15} \\ &+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) \left( (a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^* \right) \\ &\left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \right) \} = 0 \\ &+ \\ &((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\ &\left[ ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\ &\left( ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \right) \\ &+ \left( ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \right) \\ &\left( ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \right) \\ &\left( ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \right) \end{aligned}$$

$$\begin{aligned}
 & \left( ((\lambda)^{(2)})^2 + ( (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} ) (\lambda)^{(2)} \right) \\
 & + \left( ((\lambda)^{(2)})^2 + ( (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} ) (\lambda)^{(2)} \right) (q_{18})^{(2)} G_{18} \\
 & + \left( (\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)} \right) \left( (a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^* \right) \\
 & \left. \left( ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}) S_{(17),(18)} T_{17}^* + (b_{17})^{(2)} S_{(16),(18)} T_{16}^* \right) \right\} = 0 \\
 & + \\
 & \left( (\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)} \right) \left\{ (\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)} \right\} \\
 & \left[ \left( (\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)} \right) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left( (\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) S_{(21),(21)} T_{21}^* + (b_{21})^{(3)} S_{(20),(21)} T_{21}^* \\
 & + \left( (\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)} \right) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \\
 & \left( (\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) S_{(21),(20)} T_{21}^* + (b_{21})^{(3)} S_{(20),(20)} T_{20}^* \\
 & \left( (\lambda)^{(3)} \right)^2 + ( (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} ) (\lambda)^{(3)} \\
 & \left( (\lambda)^{(3)} \right)^2 + ( (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} ) (\lambda)^{(3)} \\
 & + \left( (\lambda)^{(3)} \right)^2 + ( (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} ) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + \left( (\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)} \right) \left( (a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^* \right) \\
 & \left. \left( (\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) S_{(21),(22)} T_{21}^* + (b_{21})^{(3)} S_{(20),(22)} T_{20}^* \right\} = 0 \\
 & + \\
 & \left( (\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)} \right) \left\{ (\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)} \right\} \\
 & \left[ \left( (\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)} \right) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \\
 & \left( (\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)} \right) S_{(25),(25)} T_{25}^* + (b_{25})^{(4)} S_{(24),(25)} T_{25}^* \\
 & + \left( (\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)} \right) (q_{24})^{(4)} G_{24}^* + (a_{24})^{(4)} (q_{25})^{(4)} G_{25}^* \\
 & \left( (\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)} \right) S_{(25),(24)} T_{25}^* + (b_{25})^{(4)} S_{(24),(24)} T_{24}^* \\
 & \left( (\lambda)^{(4)} \right)^2 + ( (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} ) (\lambda)^{(4)}
 \end{aligned}$$

$$\begin{aligned}
 & \left( ((\lambda)^{(4)})^2 + ( (b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)} ) (\lambda)^{(4)} \right) \\
 & + \left( ((\lambda)^{(4)})^2 + ( (a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)} ) (\lambda)^{(4)} \right) (q_{26})^{(4)} G_{26} \\
 & + \left( (\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)} \right) \left( (a_{26})^{(4)} (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (a_{26})^{(4)} (q_{24})^{(4)} G_{24}^* \right) \\
 & \left. \left( ((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)}) s_{(25),(26)} T_{25}^* + (b_{25})^{(4)} s_{(24),(26)} T_{24}^* \right) \right\} = 0 \\
 & + \\
 & \left( (\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)} \right) \left\{ (\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)} \right\} \\
 & \left[ \left( (\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)} \right) (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (q_{28})^{(5)} G_{28}^* \right] \\
 & \left( (\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)} \right) s_{(29),(29)} T_{29}^* + (b_{29})^{(5)} s_{(28),(29)} T_{29}^* \\
 & + \left( (\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)} \right) (q_{28})^{(5)} G_{28}^* + (a_{28})^{(5)} (q_{29})^{(5)} G_{29}^* \\
 & \left( (\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)} \right) s_{(29),(28)} T_{29}^* + (b_{29})^{(5)} s_{(28),(28)} T_{28}^* \\
 & \left( (\lambda)^{(5)} \right)^2 + ( (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} ) (\lambda)^{(5)} \\
 & \left( (\lambda)^{(5)} \right)^2 + ( (b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)} ) (\lambda)^{(5)} \\
 & + \left( (\lambda)^{(5)} \right)^2 + ( (a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)} ) (\lambda)^{(5)} (q_{30})^{(5)} G_{30} \\
 & + \left( (\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)} \right) \left( (a_{30})^{(5)} (q_{29})^{(5)} G_{29}^* + (a_{29})^{(5)} (a_{30})^{(5)} (q_{28})^{(5)} G_{28}^* \right) \\
 & \left. \left( (\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)} \right) s_{(29),(30)} T_{29}^* + (b_{29})^{(5)} s_{(28),(30)} T_{28}^* \right\} = 0 \\
 & + \\
 & \left( (\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)} \right) \left\{ (\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)} \right\} \\
 & \left[ \left( (\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)} \right) (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (q_{32})^{(6)} G_{32}^* \right] \\
 & \left( (\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)} \right) s_{(33),(33)} T_{33}^* + (b_{33})^{(6)} s_{(32),(33)} T_{33}^* \\
 & + \left( (\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)} \right) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \\
 & \left( (\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)} \right) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \\
 & \left( (\lambda)^{(6)} \right)^2 + ( (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} ) (\lambda)^{(6)}
 \end{aligned}$$

$$\begin{aligned} & \left( ((\lambda)^{(6)})^2 + ( (b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)} ) (\lambda)^{(6)} \right) \\ & + \left( ((\lambda)^{(6)})^2 + ( (a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)} ) (\lambda)^{(6)} \right) (q_{34})^{(6)} G_{34} \\ & + \left( (\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)} \right) \left( (a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^* \right) \\ & \left. \left( ((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \right\} = 0 \end{aligned}$$

+

$$\begin{aligned} & \left( (\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)} \right) \left\{ \left( (\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)} \right) \right. \\ & \left. \left[ \left( (\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)} \right) (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (q_{36})^{(7)} G_{36}^* \right] \right. \\ & \left. \left( ((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(37)} T_{37}^* + (b_{37})^{(7)} s_{(36),(37)} T_{37}^* \right) \right. \\ & + \left( (\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)} \right) (q_{36})^{(7)} G_{36}^* + (a_{36})^{(7)} (q_{37})^{(7)} G_{37}^* \\ & \left. \left( ((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(36)} T_{37}^* + (b_{37})^{(7)} s_{(36),(36)} T_{36}^* \right) \right. \\ & \left. \left( ((\lambda)^{(7)})^2 + ( (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} ) (\lambda)^{(7)} \right) \right. \\ & \left. \left( ((\lambda)^{(7)})^2 + ( (b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)} ) (\lambda)^{(7)} \right) \right. \\ & + \left( ((\lambda)^{(7)})^2 + ( (a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)} ) (\lambda)^{(7)} \right) (q_{38})^{(7)} G_{38} \\ & + \left( (\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)} \right) \left( (a_{38})^{(7)} (q_{37})^{(7)} G_{37}^* + (a_{37})^{(7)} (a_{38})^{(7)} (q_{36})^{(7)} G_{36}^* \right) \\ & \left. \left( ((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)}) s_{(37),(38)} T_{37}^* + (b_{37})^{(7)} s_{(36),(38)} T_{36}^* \right) \right\} = 0 \end{aligned}$$

+

$$\begin{aligned} & \left( (\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)} \right) \left\{ \left( (\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)} \right) \right. \\ & \left. \left[ \left( (\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (q_{40})^{(8)} G_{40}^* \right] \right. \\ & \left. \left( ((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(41)} T_{41}^* + (b_{41})^{(8)} s_{(40),(41)} T_{41}^* \right) \right. \\ & + \left( (\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)} \right) (q_{40})^{(8)} G_{40}^* + (a_{40})^{(8)} (q_{41})^{(8)} G_{41}^* \\ & \left. \left( ((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)}) s_{(41),(40)} T_{41}^* + (b_{41})^{(8)} s_{(40),(40)} T_{40}^* \right) \right. \\ & \left. \left( ((\lambda)^{(8)})^2 + ( (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} ) (\lambda)^{(8)} \right) \right. \end{aligned}$$

$$\begin{aligned}
 & \left( ((\lambda)^{(8)})^2 + ( (b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)} ) (\lambda)^{(8)} \right) \\
 & + \left( ((\lambda)^{(8)})^2 + ( (a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)} ) (\lambda)^{(8)} \right) (q_{42})^{(8)} G_{42} \\
 & + \left( (\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)} \right) \left( (a_{42})^{(8)} (q_{41})^{(8)} G_{41}^* + (a_{41})^{(8)} (a_{42})^{(8)} (q_{40})^{(8)} G_{40}^* \right) \\
 & \left\{ \left( (\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)} \right) s_{(41),(42)} T_{41}^* + (b_{41})^{(8)} s_{(40),(42)} T_{40}^* \right\} = 0 \\
 & + \\
 & \left( (\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)} \right) \left\{ \left( (\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)} \right) \right. \\
 & \left[ \left( (\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (q_{44})^{(9)} G_{44}^* \right] \\
 & \left( (\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)} \right) s_{(45),(45)} T_{45}^* + (b_{45})^{(9)} s_{(44),(45)} T_{45}^* \right) \\
 & + \left( (\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)} \right) (q_{44})^{(9)} G_{44}^* + (a_{44})^{(9)} (q_{45})^{(9)} G_{45}^* \\
 & \left( (\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)} \right) s_{(45),(44)} T_{45}^* + (b_{45})^{(9)} s_{(44),(44)} T_{44}^* \right) \\
 & \left. \left( (\lambda)^{(9)} \right)^2 + ( (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} ) (\lambda)^{(9)} \right) \\
 & \left( (\lambda)^{(9)} \right)^2 + ( (b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)} ) (\lambda)^{(9)} \\
 & + \left( (\lambda)^{(9)} \right)^2 + ( (a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)} ) (\lambda)^{(9)} \right) (q_{46})^{(9)} G_{46} \\
 & + \left( (\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)} \right) \left( (a_{46})^{(9)} (q_{45})^{(9)} G_{45}^* + (a_{45})^{(9)} (a_{46})^{(9)} (q_{44})^{(9)} G_{44}^* \right) \\
 & \left\{ \left( (\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)} \right) s_{(45),(46)} T_{45}^* + (b_{45})^{(9)} s_{(44),(46)} T_{44}^* \right\} = 0
 \end{aligned}$$

**And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.**

## Warped extra Dimensions and Gravity: Tweedledee and Tweedledee

**Abstract:** Habit makes you and mars you also. Sometime I think repetitive thoughts make a person mad. I have apprehensions about the same with the particles. Think that your friend snubbed and nagged you ten times, it becomes thousand fold stronger. Are constellations and galaxies perforating their duty in grand design as a matter of habit? Is death also a habit? I donot know. It is true of human beings and also other systems. We talk of muscles having memory, fluids having memory and for that matter all autopoietic systems having one so they can practice. only politicians do not have memory! They preach something and practice another! And there is no guilty consciousness! With that all the deprivational feelings are dissolved. Jack the ripper, must also have found a sadistic pleasure in doing what he did. There is one thing I can be sure of: I am going to die. But what am I to make of that fact? This course will examine a number of issues that arise once we begin to reflect on our mortality. The possibility that death may not actually be the end is considered. Are we, in some sense, immortal? Would immortality be desirable? Also a clearer notion of what it is to die is examined. What does it mean to say that a person has died? What kind of fact is that? And, finally, different attitudes to death are evaluated. Is death an evil? How? Why? Is suicide morally permissible? Is it rational? How should the knowledge that I am going to die affect the way I live my life? (Shelly Kagan). Practice action and perform and just forget it as if you have done some office work. There shall be no ego or expectations. It becomes a habit. In the eventuality you have achieved sat Chit Ananda, although you might have that perpetual frown on the face. Believe me it is true. Tell baby from the beginning that this is space and this is time and you are on a perpetual search which nobody knows. A few days I was writing something and it became very clear God is not in an hat NAD or the space between ragas, but really cares two hoots for the language. That is the ultimate destiny. Experience the fact like Ramakrishna and you shall get the results.


### INTRODUCTION—VARIABLES USED

National Geographic Society: The Greatest Unsolved Problem in Theoretical Physics



Posted by Ethan on September 19, 2012

There are two other possibilities, one which is much more promising than the other, and both involving extra



dimensions.

Image credit: Cetin BAL, as far as I can tell.

- (1) Warped Extra Dimensions. This theory — pioneered by the aforementioned Lisa Randall along with Raman Sundrum — hold that gravity **is just** ( $\equiv$ ) **as** strong as the other forces, but not in our three-spatial-dimension Universe.
- (2) It lives in a different three-spatial-dimension Universe that's **offset** ( $\equiv$ , **e**, **eb**) by some tiny amount — like 10<sup>-31</sup> meters — from our own Universe in the fourth spatial dimension.
- (3) (Or, as the diagram above indicates, in the fifth dimension, once time is included.) This is interesting, because it would be stable, and it **could (eb) provide a** possible explanation as to why our Universe began expanding so rapidly at the beginning (warped spacetime can do that), so it's got some compelling perks.
- (4) What it **should also (e) include** are an extra set of particles; not supersymmetric particles, but Kaluza-Klein particles, which are a direct **consequence (e) of** there being extra dimensions.
- (5) For what it's worth, there has been a hint from one experiment in space that there **might (eb) be a** Kaluza-Klein particle at energy of about 600 GeV, or about 5 times the mass of the Higgs.




Image credit: J. Chang et al. (2008), Nature, from the Advanced Thin Ionization Calorimeter (ATIC).

- (6) This is by no means a certainty, **as it's(=) just** an excess of observed electrons **over(e) the** expected background, but it's worth keeping in mind as the LHC eventually ramps up to full energy; a new particle that's below 1,000 GeV in mass should be within range of this machine.

And finally...




Image credit: Universe-review.ca.

- (7) Large Extra Dimensions. Instead of being warped, the extra dimensions could be “large”, where large is only large **relative (e) to** the warped ones, which were 10<sup>-31</sup> meters in scale.
- (8) The “large” extra dimensions would be around millimeter-sized, which meant that new particles would start **showing (eb) up right** around the scale that the LHC is capable of probing.
- (9) Again, there would be new Kaluza-Klein particles, and this could be a possible **solution (e) to the** hierarchy problem.

## NOTATION

### Module One

Warped Extra Dimensions. This theory — pioneered by the aforementioned Lisa Randall along with Raman Sundrum — hold that gravity **is just (=) as** strong as the other forces, but not in our three-spatial-dimension Universe

$G_{13}$  : Category one of strong as the other forces, but not in our three-spatial-dimension Universe

$G_{14}$  : Category two of strong as the other forces, but not in our three-spatial-dimension Universe

$G_{15}$  : Category three of strong as the other forces, but not in our three-spatial-dimension Universe

$T_{13}$  : Category one of Warped Extra Dimensions. This theory — pioneered by the aforementioned Lisa Randall along with Raman Sundrum — hold that gravity

$T_{14}$  : Category two of Warped Extra Dimensions. This theory — pioneered by the aforementioned Lisa Randall along with Raman Sundrum — hold that gravity

$T_{15}$  : Category three of Warped Extra Dimensions. This theory — pioneered by the aforementioned Lisa Randall along with Raman Sundrum — hold that gravity

### Module Two(Care: both the cases are discussed)

It lives in a different three-spatial-dimension Universe that's **offset (=, e, eb)** by some tiny amount — like 10<sup>-31</sup> meters — from our own Universe in the fourth spatial dimension

$G_{16}$  : Category one of gravity lives in a different three-spatial-dimension Universe; some tiny amount — like 10<sup>-31</sup> meters — from our own Universe in the fourth spatial dimension

$G_{17}$  : Category two of gravity lives in a different three-spatial-dimension Universe; some tiny amount — like 10<sup>-31</sup> meters — from our own Universe in the fourth spatial dimension

$G_{18}$  : Category three of gravity lives in a different three-spatial-dimension Universe; some tiny amount — like 10<sup>-31</sup> meters — from our own Universe in the fourth spatial dimension

$T_{16}$  : Category one of some tiny amount — like 10<sup>-31</sup> meters — from our own Universe in the fourth spatial

dimension; gravity lives in a different three-spatial-dimension Universe

$T_{17}$  : Category two of some tiny amount — like 10-31 meters — from our own Universe in the fourth spatial dimension ;gravity lives in a different three-spatial-dimension Universe

$T_{18}$  : Category three of some tiny amount — like 10-31 meters — from our own Universe in the fourth spatial dimension; gravity lives in a different three-spatial-dimension Universe

### Module three

(Or, as the diagram above indicates, in the fifth dimension, once time is included.) This is interesting, because gravity it would be stable, and it **could (eb) provide a** possible explanation as to why our Universe began expanding so rapidly at the beginning (warped spacetime can do that), so it's got some compelling perks.

$G_{20}$  : Category one of gravity it would be stable (Please see the background above)

$G_{21}$  : Category two of gravity it would be stable

$G_{22}$  : Category three of gravity it would be stable

$T_{20}$  : Category one of possible explanation as to why our Universe began expanding so rapidly at the beginning (warped spacetime can do that), so it's got some compelling perks

$T_{21}$  : Category two of possible explanation as to why our Universe began expanding so rapidly at the beginning (warped spacetime can do that), so it's got some compelling perks

$T_{22}$  : Category three of possible explanation as to why our Universe began expanding so rapidly at the beginning (warped spacetime can do that), so it's got some compelling perks

### Module four

What it **should also (e) include** are an extra set of particles; not supersymmetric particles, but Kaluza-Klein particles, which are a direct **consequence (e) of** there being extra dimensions

$G_{24}$  : Category one of gravity and warped extra dimensions **should also include** are an extra set of particles; not supersymmetric particles, but Kaluza-Klein particles

$G_{25}$  : Category two of gravity and warped extra dimensions **should also include** are an extra set of particles; not supersymmetric particles, but Kaluza-Klein particles

$G_{26}$  : Category three of gravity and warped extra dimensions **should also include** are an extra set of particles; not supersymmetric particles, but Kaluza-Klein particles

$T_{24}$  : Category one of extra dimensions

$T_{25}$  : Category two of extra dimensions

$T_{26}$  : Category three of extra dimensions

### Module five

For what it's worth, there has been a hint from one experiment in space that there **might (eb) be a** Kaluza-Klein particle at energy of about 600 GeV, or about 5 times the mass of the Higgs

$G_{28}$  : Category one of experiment in space gives insinuations and innuendos

$G_{29}$  : Category two of experiment in space gives insinuations and innuendos

$G_{30}$  : Category three of experiment in space gives insinuations and innuendos

$T_{28}$  : Category one of Kaluza-Klein particle at energy of about 600 GeV, or about 5 times the mass of the Higgs

$T_{29}$  : Category two of Kaluza-Klein particle at energy of about 600 GeV, or about 5 times the mass of the Higgs

$T_{30}$  : Category three of Kaluza-Klein particle at energy of about 600 GeV, or about 5 times the mass of the Higgs

### Module six

This is by no means a certainty, **as it's(=) just** an excess of observed electrons **over(e) the** expected background, but it's worth keeping in mind as the LHC eventually ramps up to full energy; a new particle that's below 1,000 GeV in mass should be within range of this machine

$G_{32}$  : Category one of indeterminism of Kaluza-Klein particle at energy of about 600 GeV, or about 5 times the mass of the Higgs

$G_{33}$  : Category two of indeterminism of Kaluza-Klein particle at energy of about 600 GeV, or about 5 times the mass of the Higgs

$G_{34}$  : Category three of indeterminism of Kaluza-Klein particle at energy of about 600 GeV, or about 5 times the mass of the Higgs

$T_{32}$  : Category one of excess of observed electrons **over(e) the** expected background, but it's worth keeping in mind as the LHC eventually ramps up to full energy; a new particle that's below 1,000 GeV in mass should be within range of this machine

$T_{33}$  : Category two of excess of observed electrons **over(e) the** expected background, but it's worth keeping in mind as the LHC eventually ramps up to full energy; a new particle that's below 1,000 GeV in mass should be within range of this machine

$T_{34}$  : Category three of excess of observed electrons **over(e) the** expected background, but it's worth keeping in mind as the LHC eventually ramps up to full energy; a new particle that's below 1,000 GeV in mass should be within range of this machine

### Module seven

Large Extra Dimensions. Instead of being warped, the extra dimensions could be "large", where large is only large **relative (e) to** the warped ones, which were 10-31 meters in scale

$G_{36}$  : Category one of Instead of being warped, the extra dimensions could be "large", where large is only large; warped ones, which were 10-31 meters in scale

$G_{37}$  : Category two of Instead of being warped, the extra dimensions could be "large", where large is only large; warped ones, which were 10-31 meters in scale

$G_{38}$  : Category three of Instead of being warped, the extra dimensions could be "large", where large is only large; warped ones, which were 10-31 meters in scale

$T_{36}$  : Category one of warped ones, which were 10-31 meters in scale; Instead of being warped, the extra dimensions could be “large”, where large is only large

$T_{37}$  : Category two of warped ones, which were 10-31 meters in scale; Instead of being warped, the extra dimensions could be “large”, where large is only large

$T_{38}$  : Category three of warped ones, which were 10-31 meters in scale ;Instead of being warped, the extra dimensions could be “large”, where large is only large

### Module eight

The “large” extra dimensions would be around millimeter-sized, which meant that new particles would start **showing (eb) up right** around the scale that the LHC is capable of probing

$G_{40}$  : Category one of “large” extra dimensions would be around millimeter-sized, which meant that new particles

$G_{41}$  : Category two of “large” extra dimensions would be around millimeter-sized, which meant that new particles

$G_{42}$  : Category three of “large” extra dimensions would be around millimeter-sized, which meant that new particles

$T_{40}$  : Category one of **right** around the scale that the LHC is capable of probing. It is to be repeated that we are talking of system that satisfy the conditions of the so stated axiomatic predication in the foregoing.

$T_{41}$  : Category two of **right** around the scale that the LHC is capable of probing

$T_{42}$  : Category three of **right** around the scale that the LHC is capable of probing

### Module Nine

Again, there would be new Kaluza-Klein particles, and this could be a possible **solution (e) to the hierarchy problem**

$G_{44}$  : Category one of Kaluza-Klein particles, and this could be a possible **solution**

$G_{45}$  : Category two of Kaluza-Klein particles, and this could be a possible **solution**

$G_{46}$  : Category three of Kaluza-Klein particles, and this could be a possible **solution**

$T_{44}$  : Category one of hierarchy problem

$T_{45}$  : Category two of hierarchy problem

$T_{46}$  : Category three of hierarchy problem

### The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}:$   
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$   
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}, (a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)},$   
 $(a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$   
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$

$$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$$

$$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$$

are Accentuation coefficients

$$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}, (b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}$$

$$, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$$

$$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}, (a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)},$$

$$(a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$$

$$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$$

$$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$$

$$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$$

are Dissipation coefficients

**Module Numbered One**

**The differential system of this model is now (Module Numbered one)**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} \quad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$  First augmentation factor  
 $-(b''_{13})^{(1)}(G, t) =$  First detritions factor

**Module Numbered Two**

**The differential system of this model is now (Module numbered two)**

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$  First augmentation factor  
 $-(b''_{16})^{(2)}((G_{19}), t) =$  First detritions factor

**Module Numbered Three**

**The differential system of this model is now (Module numbered three)**

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$  First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$  First detritions factor

**Module Numbered Four**

**The differential system of this model is now (Module numbered Four)**

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}, t))]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}, t))]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}, t))]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$  First augmentation factor

$-(b''_{24})^{(4)}((G_{27}, t)) =$  First detritions factor

**Module Numbered Five:**

**The differential system of this model is now (Module number five)**

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}, t))]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}, t))]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}, t))]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$  First augmentation factor

$-(b''_{28})^{(5)}((G_{31}, t)) =$  First detritions factor

**Module Numbered Six**

**The differential system of this model is now (Module numbered Six)**

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}, t))]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}, t))]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}, t))]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$  First augmentation factor

**Module Numbered Seven:**

**The differential system of this model is now (Seventh Module)**

$$\begin{aligned} \frac{dG_{36}}{dt} &= (a_{36})^{(7)} G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)] G_{36} & 37 \\ \frac{dG_{37}}{dt} &= (a_{37})^{(7)} G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)] G_{37} & 38 \\ \frac{dG_{38}}{dt} &= (a_{38})^{(7)} G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)] G_{38} & 39 \\ \frac{dT_{36}}{dt} &= (b_{36})^{(7)} T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}, t))] T_{36} & 40 \\ \frac{dT_{37}}{dt} &= (b_{37})^{(7)} T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}, t))] T_{37} & 41 \\ \frac{dT_{38}}{dt} &= (b_{38})^{(7)} T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}, t))] T_{38} & 42 \\ &+ (a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor} \end{aligned}$$

**Module Numbered Eight**

The differential system of this model is now

$$\begin{aligned} \frac{dG_{40}}{dt} &= (a_{40})^{(8)} G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)] G_{40} & 43 \\ \frac{dG_{41}}{dt} &= (a_{41})^{(8)} G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)] G_{41} & 44 \\ \frac{dG_{42}}{dt} &= (a_{42})^{(8)} G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)] G_{42} & 45 \\ \frac{dT_{40}}{dt} &= (b_{40})^{(8)} T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}, t))] T_{40} & 46 \\ \frac{dT_{41}}{dt} &= (b_{41})^{(8)} T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}, t))] T_{41} & 47 \\ \frac{dT_{42}}{dt} &= (b_{42})^{(8)} T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}, t))] T_{42} & 48 \end{aligned}$$

**Module Numbered Nine**

The differential system of this model is now

$$\begin{aligned} \frac{dG_{44}}{dt} &= (a_{44})^{(9)} G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)] G_{44} & 49 \\ \frac{dG_{45}}{dt} &= (a_{45})^{(9)} G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)] G_{45} & 50 \\ \frac{dG_{46}}{dt} &= (a_{46})^{(9)} G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)] G_{46} & 51 \\ \frac{dT_{44}}{dt} &= (b_{44})^{(9)} T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}, t))] T_{44} & 52 \\ \frac{dT_{45}}{dt} &= (b_{45})^{(9)} T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}, t))] T_{45} & 53 \\ \frac{dT_{46}}{dt} &= (b_{46})^{(9)} T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}, t))] T_{46} & 54 \\ &+ (a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor} \\ &- (b''_{44})^{(9)}((G_{47}, t)) = \text{First detrition factor} \end{aligned}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[ \begin{array}{c} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)} G_{13} - \left[ \begin{array}{c} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)} G_{14} - \left[ \begin{array}{c} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \quad 57$$

Where  $(a''_{13})^{(1)}(T_{14}, t)$ ,  $(a''_{14})^{(1)}(T_{14}, t)$ ,  $(a''_{15})^{(1)}(T_{14}, t)$  are first augmentation coefficients for category 1, 2 and 3

$(a''_{16})^{(2,2)}(T_{17}, t)$ ,  $(a''_{17})^{(2,2)}(T_{17}, t)$ ,  $(a''_{18})^{(2,2)}(T_{17}, t)$  are second augmentation coefficient for category 1, 2 and 3

$(a''_{20})^{(3,3)}(T_{21}, t)$ ,  $(a''_{21})^{(3,3)}(T_{21}, t)$ ,  $(a''_{22})^{(3,3)}(T_{21}, t)$  are third augmentation coefficient for category 1, 2 and 3

$(a''_{24})^{(4,4,4,4)}(T_{25}, t)$ ,  $(a''_{25})^{(4,4,4,4)}(T_{25}, t)$ ,  $(a''_{26})^{(4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficient for category 1, 2 and 3

$(a''_{28})^{(5,5,5,5)}(T_{29}, t)$ ,  $(a''_{29})^{(5,5,5,5)}(T_{29}, t)$ ,  $(a''_{30})^{(5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficient for category 1, 2 and 3

$(a''_{32})^{(6,6,6,6)}(T_{33}, t)$ ,  $(a''_{33})^{(6,6,6,6)}(T_{33}, t)$ ,  $(a''_{34})^{(6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficient for category 1, 2 and 3

$(a''_{38})^{(7,7)}(T_{37}, t)$ ,  $(a''_{37})^{(7,7)}(T_{37}, t)$ ,  $(a''_{36})^{(7,7)}(T_{37}, t)$  are seventh augmentation coefficient for 1,2,3

$(a''_{40})^{(8,8)}(T_{41}, t)$ ,  $(a''_{41})^{(8,8)}(T_{41}, t)$ ,  $(a''_{42})^{(8,8)}(T_{41}, t)$  are eight augmentation coefficient for 1,2,3

$(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ ,  $(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ ,  $(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$  are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[ \begin{array}{l} (b'_{13})^{(1)}(G, t) \quad - (b'_{16})^{(2,2)}(G_{19}, t) \quad - (b'_{20})^{(3,3)}(G_{23}, t) \\ - (b'_{24})^{(4,4,4,4)}(G_{27}, t) \quad - (b'_{28})^{(5,5,5,5)}(G_{31}, t) \quad - (b'_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b'_{36})^{(7,7)}(G_{39}, t) \quad - (b'_{40})^{(8,8)}(G_{43}, t) \quad - (b'_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \tag{58}$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[ \begin{array}{l} (b'_{14})^{(1)}(G, t) \quad - (b'_{17})^{(2,2)}(G_{19}, t) \quad - (b'_{21})^{(3,3)}(G_{23}, t) \\ - (b'_{25})^{(4,4,4,4)}(G_{27}, t) \quad - (b'_{29})^{(5,5,5,5)}(G_{31}, t) \quad - (b'_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b'_{37})^{(7,7)}(G_{39}, t) \quad - (b'_{41})^{(8,8)}(G_{43}, t) \quad - (b'_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \tag{59}$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[ \begin{array}{l} (b'_{15})^{(1)}(G, t) \quad - (b'_{18})^{(2,2)}(G_{19}, t) \quad - (b'_{22})^{(3,3)}(G_{23}, t) \\ - (b'_{26})^{(4,4,4,4)}(G_{27}, t) \quad - (b'_{30})^{(5,5,5,5)}(G_{31}, t) \quad - (b'_{34})^{(6,6,6,6)}(G_{35}, t) \\ - (b'_{38})^{(7,7)}(G_{39}, t) \quad - (b'_{42})^{(8,8)}(G_{43}, t) \quad - (b'_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15} \tag{60}$$

Where  $(b'_{13})^{(1)}(G, t)$ ,  $(b'_{14})^{(1)}(G, t)$ ,  $(b'_{15})^{(1)}(G, t)$  are first detrition coefficients for category 1, 2 and 3

$(b'_{16})^{(2,2)}(G_{19}, t)$ ,  $(b'_{17})^{(2,2)}(G_{19}, t)$ ,  $(b'_{18})^{(2,2)}(G_{19}, t)$  are second detrition coefficients for category 1, 2 and 3

$(b'_{20})^{(3,3)}(G_{23}, t)$ ,  $(b'_{21})^{(3,3)}(G_{23}, t)$ ,  $(b'_{22})^{(3,3)}(G_{23}, t)$  are third detrition coefficients for category 1, 2 and 3

$(b'_{24})^{(4,4,4,4)}(G_{27}, t)$ ,  $(b'_{25})^{(4,4,4,4)}(G_{27}, t)$ ,  $(b'_{26})^{(4,4,4,4)}(G_{27}, t)$  are fourth detrition coefficients for category 1, 2 and 3

$(b'_{28})^{(5,5,5,5)}(G_{31}, t)$ ,  $(b'_{29})^{(5,5,5,5)}(G_{31}, t)$ ,  $(b'_{30})^{(5,5,5,5)}(G_{31}, t)$  are fifth detrition coefficients for category 1, 2 and 3

$(b'_{32})^{(6,6,6,6)}(G_{35}, t)$ ,  $(b'_{33})^{(6,6,6,6)}(G_{35}, t)$ ,  $(b'_{34})^{(6,6,6,6)}(G_{35}, t)$  are sixth detrition coefficients for

category 1, 2 and 3

$-(b''_{37})^{(7,7)}(G_{39}, t)$ ,  $-(b''_{36})^{(7,7)}(G_{39}, t)$ ,  $-(b''_{38})^{(7,7)}(G_{39}, t)$  are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43}, t)$ ,  $-(b''_{41})^{(8,8)}(G_{43}, t)$ ,  $-(b''_{42})^{(8,8)}(G_{43}, t)$  are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$  are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[ \begin{array}{l} (a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) + (a'_{13})^{(1,1)}(T_{14}, t) + (a'_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \tag{61}$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[ \begin{array}{l} (a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t) + (a'_{14})^{(1,1)}(T_{14}, t) + (a'_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \tag{62}$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[ \begin{array}{l} (a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t) + (a'_{15})^{(1,1)}(T_{14}, t) + (a'_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \tag{63}$$

Where  $+(a''_{16})^{(2)}(T_{17}, t)$ ,  $+(a''_{17})^{(2)}(T_{17}, t)$ ,  $+(a''_{18})^{(2)}(T_{17}, t)$  are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1)}(T_{14}, t)$ ,  $+(a''_{14})^{(1,1)}(T_{14}, t)$ ,  $+(a''_{15})^{(1,1)}(T_{14}, t)$  are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21}, t)$ ,  $+(a''_{21})^{(3,3,3)}(T_{21}, t)$ ,  $+(a''_{22})^{(3,3,3)}(T_{21}, t)$  are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)$ ,  $+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)$ ,  $+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)$ ,  $+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)$ ,  $+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)$ ,  $+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)$ ,  $+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37}, t)$ ,  $+(a''_{37})^{(7,7,7)}(T_{37}, t)$ ,  $+(a''_{38})^{(7,7,7)}(T_{37}, t)$  are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41}, t)$ ,  $+(a''_{41})^{(8,8,8)}(T_{41}, t)$ ,  $+(a''_{42})^{(8,8,8)}(T_{41}, t)$  are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45}, t)$ ,  $+(a''_{45})^{(9,9)}(T_{45}, t)$ ,  $+(a''_{46})^{(9,9)}(T_{45}, t)$  are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[ \begin{array}{l} (b'_{16})^{(2)} \boxed{-(b''_{16})^{(2)}(G_{19}, t)} \quad \boxed{-(b''_{13})^{(1,1)}(G, t)} \quad \boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)} \quad \boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)} \quad \boxed{-(b''_{44})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{16} \tag{64}$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[ \begin{array}{l} (b'_{17})^{(2)} \boxed{-(b''_{17})^{(2)}(G_{19}, t)} \quad \boxed{-(b''_{14})^{(1,1)}(G, t)} \quad \boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)} \quad \boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)} \quad \boxed{-(b''_{45})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{17} \tag{65}$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[ \begin{array}{l} (b'_{18})^{(2)} \boxed{-(b''_{18})^{(2)}(G_{19}, t)} \quad \boxed{-(b''_{15})^{(1,1)}(G, t)} \quad \boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} \quad \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} \quad \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)} \quad \boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)} \quad \boxed{-(b''_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \tag{66}$$

where  $\boxed{-(b''_{16})^{(2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2)}(G_{19}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1)}(G, t)}$  are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)}$  are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$  are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$  are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$  are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)}$  are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)}$  are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{46})^{(9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{45})^{(9,9)}(G_{47}, t)}$  are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[ \begin{array}{l} (a'_{20})^{(3)} \boxed{+(a''_{20})^{(3)}(T_{21}, t)} \quad \boxed{+(a''_{16})^{(2,2,2)}(T_{17}, t)} \quad \boxed{+(a''_{13})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)} \quad \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)} \quad \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)} \quad \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41}, t)} \quad \boxed{+(a''_{44})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{20} \tag{67}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[ \begin{array}{l} (a'_{21})^{(3)} \boxed{+(a''_{21})^{(3)}(T_{21}, t)} \quad \boxed{+(a''_{17})^{(2,2,2)}(T_{17}, t)} \quad \boxed{+(a''_{14})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)} \quad \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)} \quad \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)} \quad \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41}, t)} \quad \boxed{+(a''_{45})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{21} \tag{68}$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[ \begin{array}{ccc} (a'_{22})^{(3)} & + (a''_{22})^{(3)}(T_{21}, t) & + (a'_{18})^{(2,2,2)}(T_{17}, t) & + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ + (a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t) & \\ + (a'_{38})^{(7,7,7,7)}(T_{37}, t) & + (a'_{42})^{(8,8,8,8)}(T_{41}, t) & + (a'_{46})^{(9,9,9)}(T_{45}, t) & \end{array} \right] G_{22} \tag{69}$$

$+ (a'_{20})^{(3)}(T_{21}, t)$ ,  $+ (a'_{21})^{(3)}(T_{21}, t)$ ,  $+ (a'_{22})^{(3)}(T_{21}, t)$  are first augmentation coefficients for category 1, 2 and 3

$+ (a'_{16})^{(2,2,2)}(T_{17}, t)$ ,  $+ (a'_{17})^{(2,2,2)}(T_{17}, t)$ ,  $+ (a'_{18})^{(2,2,2)}(T_{17}, t)$  are second augmentation coefficients for category 1, 2 and 3

$+ (a'_{13})^{(1,1,1)}(T_{14}, t)$ ,  $+ (a'_{14})^{(1,1,1)}(T_{14}, t)$ ,  $+ (a'_{15})^{(1,1,1)}(T_{14}, t)$  are third augmentation coefficients for category 1, 2 and 3

$+ (a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t)$ ,  $+ (a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t)$ ,  $+ (a'_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficients for category 1, 2 and 3

$+ (a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t)$ ,  $+ (a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t)$ ,  $+ (a'_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficients for category 1, 2 and 3

$+ (a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t)$ ,  $+ (a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t)$ ,  $+ (a'_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficients for category 1, 2 and 3

$+ (a'_{36})^{(7,7,7,7)}(T_{37}, t)$ ,  $+ (a'_{37})^{(7,7,7,7)}(T_{37}, t)$ ,  $+ (a'_{38})^{(7,7,7,7)}(T_{37}, t)$  are seventh augmentation coefficients for category 1, 2 and 3

$+ (a'_{40})^{(8,8,8,8)}(T_{41}, t)$ ,  $+ (a'_{41})^{(8,8,8,8)}(T_{41}, t)$ ,  $+ (a'_{42})^{(8,8,8,8)}(T_{41}, t)$  are eight augmentation coefficients for category 1, 2 and 3

$+ (a'_{44})^{(9,9,9)}(T_{45}, t)$ ,  $+ (a'_{45})^{(9,9,9)}(T_{45}, t)$ ,  $+ (a'_{46})^{(9,9,9)}(T_{45}, t)$  are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[ \begin{array}{ccc} (b'_{20})^{(3)} & - (b''_{20})^{(3)}(G_{23}, t) & - (b'_{16})^{(2,2,2)}(G_{19}, t) & - (b'_{13})^{(1,1,1)}(G, t) \\ - (b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{28})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{32})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{36})^{(7,7,7,7)}(G_{39}, t) & - (b'_{40})^{(8,8,8,8)}(G_{43}, t) & - (b'_{44})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{20} \tag{70}$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[ \begin{array}{ccc} (b'_{21})^{(3)} & - (b''_{21})^{(3)}(G_{23}, t) & - (b'_{17})^{(2,2,2)}(G_{19}, t) & - (b'_{14})^{(1,1,1)}(G, t) \\ - (b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{29})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{33})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{37})^{(7,7,7,7)}(G_{39}, t) & - (b'_{41})^{(8,8,8,8)}(G_{43}, t) & - (b'_{45})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{21} \tag{71}$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[ \begin{array}{ccc} (b'_{22})^{(3)} & - (b''_{22})^{(3)}(G_{23}, t) & - (b'_{18})^{(2,2,2)}(G_{19}, t) & - (b'_{15})^{(1,1,1)}(G, t) \\ - (b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t) & - (b'_{30})^{(5,5,5,5,5,5)}(G_{31}, t) & - (b'_{34})^{(6,6,6,6,6,6)}(G_{35}, t) & \\ - (b'_{38})^{(7,7,7,7)}(G_{39}, t) & - (b'_{42})^{(8,8,8,8)}(G_{43}, t) & - (b'_{46})^{(9,9,9)}(G_{47}, t) & \end{array} \right] T_{22} \tag{72}$$

$- (b''_{20})^{(3)}(G_{23}, t)$ ,  $- (b''_{21})^{(3)}(G_{23}, t)$ ,  $- (b''_{22})^{(3)}(G_{23}, t)$  are first detrition coefficients for category 1, 2 and 3

$- (b'_{16})^{(2,2,2)}(G_{19}, t)$ ,  $- (b'_{17})^{(2,2,2)}(G_{19}, t)$ ,  $- (b'_{18})^{(2,2,2)}(G_{19}, t)$  are second detrition coefficients for category 1, 2 and 3

$- (b'_{13})^{(1,1,1)}(G, t)$ ,  $- (b'_{14})^{(1,1,1)}(G, t)$ ,  $- (b'_{15})^{(1,1,1)}(G, t)$  are third detrition coefficients for category 1, 2 and 3

$- (b'_{24})^{(4,4,4,4,4,4)}(G_{27}, t)$ ,  $- (b'_{25})^{(4,4,4,4,4,4)}(G_{27}, t)$ ,  $- (b'_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$  are fourth detrition

coefficients for category 1, 2 and 3

$$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}, \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}, \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} \text{ are fifth detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}, \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}, \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \text{ are sixth detrition}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39}, t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39}, t)}, \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39}, t)} \text{ are seventh detrition coefficients}$$

for category 1, 2 and 3

$$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43}, t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43}, t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43}, t)} \text{ are eight detrition coefficients for}$$

category 1, 2 and 3

$$\boxed{-(b''_{46})^{(9,9,9)}(G_{47}, t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47}, t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47}, t)} \text{ are ninth detrition coefficients for}$$

category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[ \begin{array}{ccc} \boxed{(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)} & \boxed{+(a''_{28})^{(5,5)}(T_{29}, t)} & \boxed{+(a''_{32})^{(6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[ \begin{array}{ccc} \boxed{(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)} & \boxed{+(a''_{29})^{(5,5)}(T_{29}, t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33}, t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[ \begin{array}{ccc} \boxed{(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)} & \boxed{+(a''_{30})^{(5,5)}(T_{29}, t)} & \boxed{+(a''_{34})^{(6,6)}(T_{33}, t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14}, t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17}, t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37}, t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41}, t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{26} \quad 75$$

$$\boxed{(a''_{24})^{(4)}(T_{25}, t)}, \boxed{(a''_{25})^{(4)}(T_{25}, t)}, \boxed{(a''_{26})^{(4)}(T_{25}, t)} \text{ are first augmentation coefficients}$$

category 1, 2 3

$$\boxed{+(a''_{28})^{(5,5)}(T_{29}, t)}, \boxed{+(a''_{29})^{(5,5)}(T_{29}, t)}, \boxed{+(a''_{30})^{(5,5)}(T_{29}, t)} \text{ are second augmentation}$$

coefficient for category 1, 2 and 3

$$\boxed{+(a''_{32})^{(6,6)}(T_{33}, t)}, \boxed{+(a''_{33})^{(6,6)}(T_{33}, t)}, \boxed{+(a''_{34})^{(6,6)}(T_{33}, t)} \text{ are third augmentation}$$

coefficient for category 1, 2 and 3

$$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14}, t)}$$

are fourth augmentation coefficients for category 1, 2 and 3

$$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17}, t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17}, t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17}, t)}$$

are fifth augmentation coefficients for category 1, 2 and 3

$$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21}, t)}$$

are sixth augmentation coefficients for category 1, 2 and 3

$$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37}, t)}$$

are seventh augmentation coefficients for category 1, 2 and 3

$$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41}, t)}$$

are eighth augmentation coefficients for category 1, 2 and 3

$$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45}, t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45}, t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45}, t)} \text{ are ninth detrition coefficients for}$$

category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[ \begin{array}{ccc} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) & - (b''_{28})^{(5,5)}(G_{31}, t) & - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \tag{76}$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[ \begin{array}{ccc} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) & - (b''_{29})^{(5,5)}(G_{31}, t) & - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \tag{77}$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[ \begin{array}{ccc} (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}, t) & - (b''_{30})^{(5,5)}(G_{31}, t) & - (b''_{34})^{(6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{26} \tag{78}$$

Where  $-(b''_{24})^{(4)}(G_{27}, t)$ ,  $-(b''_{25})^{(4)}(G_{27}, t)$ ,  $-(b''_{26})^{(4)}(G_{27}, t)$  are first detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5)}(G_{31}, t)$  are second detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6,6)}(G_{35}, t)$  are third detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1,1,1)}(G, t)$ ,  $-(b''_{15})^{(1,1,1,1)}(G, t)$  are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2,2,2,2)}(G_{19}, t)$  are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3,3,3)}(G_{23}, t)$  are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)$  are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)$  are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{45})^{(9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{44})^{(9,9,9,9)}(G_{47}, t)$  are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[ \begin{array}{ccc} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) & + (a''_{24})^{(4,4)}(T_{25}, t) & + (a''_{32})^{(6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \tag{79}$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[ \begin{array}{ccc} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) & + (a''_{25})^{(4,4)}(T_{25}, t) & + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \tag{80}$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[ \begin{array}{ccc} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) & + (a''_{26})^{(4,4)}(T_{25}, t) & + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \tag{81}$$

Where  $\boxed{+(a''_{28})^{(5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5)}(T_{29}, t)}$  are first augmentation coefficients for category 1, 2 and 3

And  $\boxed{+(a''_{24})^{(4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4)}(T_{25}, t)}$  are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6)}(T_{33}, t)}$  are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)}$  are fourth augmentation coefficients for category 1, 2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)}$  are fifth augmentation coefficients for category 1, 2, and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)}$  are sixth augmentation coefficients for category 1, 2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)}$  are seventh augmentation coefficients for category 1, 2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)}$  are eighth augmentation coefficients for category 1, 2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)}$  are ninth augmentation coefficients for category 1, 2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[ \begin{array}{ccc} \boxed{(b'_{28})^{(5)}} & \boxed{-(b''_{28})^{(5)}(G_{31}, t)} & \boxed{-(b''_{24})^{(4,4)}(G_{27}, t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)} & \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)} & \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[ \begin{array}{ccc} \boxed{(b'_{29})^{(5)}} & \boxed{-(b''_{29})^{(5)}(G_{31}, t)} & \boxed{-(b''_{25})^{(4,4)}(G_{27}, t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)} & \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)} & \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[ \begin{array}{ccc} \boxed{(b'_{30})^{(5)}} & \boxed{-(b''_{30})^{(5)}(G_{31}, t)} & \boxed{-(b''_{26})^{(4,4)}(G_{27}, t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)} & \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)} & \end{array} \right] T_{30} \quad 84$$

where  $\boxed{-(b''_{28})^{(5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5)}(G_{31}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4)}(G_{27}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)}$  are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)}$  are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)}$  are sixth detrition coefficients

for category 1,2, and 3

$$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t), -(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)$$

are seventh detrition coefficients for category 1,2, and 3

$$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t), -(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)$$

are eighth detrition coefficients for category 1,2, and 3

$$-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t), -(b''_{45})^{(9,9,9,9,9)}(G_{47}, t), -(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)$$

are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} \tag{85}$$

$$- \left[ \begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[ \begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \tag{86}$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[ \begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) & + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \tag{87}$$

$$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$$

are first augmentation coefficients for category 1, 2 and 3

$$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$$

are second augmentation coefficients for category 1, 2 and 3

$$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$$

are third augmentation coefficients for category 1, 2 and 3

$$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$$

- are fourth augmentation coefficients

$$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$$

- fifth augmentation coefficients

$$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$$

sixth augmentation coefficients

$$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$$

seventh augmentation coefficients

$$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$$

Eighth augmentation coefficients

$$+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$$

ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[ \begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[ \begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[ \begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t) & - (b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{34} \quad 90$$

$[-(b''_{32})^{(6)}(G_{35}, t)]$ ,  $[-(b''_{33})^{(6)}(G_{35}, t)]$ ,  $[-(b''_{34})^{(6)}(G_{35}, t)]$  are first detrition coefficients for category 1, 2 and 3

$[-(b''_{28})^{(5,5,5)}(G_{31}, t)]$ ,  $[-(b''_{29})^{(5,5,5)}(G_{31}, t)]$ ,  $[-(b''_{30})^{(5,5,5)}(G_{31}, t)]$  are second detrition coefficients for category 1, 2 and 3

$[-(b''_{24})^{(4,4,4)}(G_{27}, t)]$ ,  $[-(b''_{25})^{(4,4,4)}(G_{27}, t)]$ ,  $[-(b''_{26})^{(4,4,4)}(G_{27}, t)]$  are third detrition coefficients for category 1, 2 and 3

$[-(b''_{13})^{(1,1,1,1,1,1)}(G, t)]$ ,  $[-(b''_{14})^{(1,1,1,1,1,1)}(G, t)]$ ,  $[-(b''_{15})^{(1,1,1,1,1,1)}(G, t)]$  are fourth detrition coefficients for category 1, 2, and 3

$[-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)]$ ,  $[-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)]$ ,  $[-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)]$  are fifth detrition coefficients for category 1, 2, and 3

$[-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)]$ ,  $[-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)]$ ,  $[-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)]$  are sixth detrition coefficients for category 1, 2, and 3

$[-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)]$ ,  $[-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)]$ ,  $[-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)]$  are seventh detrition coefficients for category 1, 2, and 3

$[-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)]$ ,  $[-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)]$ ,  $[-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)]$  are eighth detrition coefficients for category 1, 2, and 3

$[-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)]$ ,  $[-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)]$ ,  $[-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)]$  are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \left[ \begin{array}{ccc} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \left[ \begin{array}{ccc} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt}$$

93

$$= (a_{38})^{(7)} G_{37} - \left[ \begin{array}{ccc} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) & + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where  $(a''_{36})^{(7)}(T_{37}, t)$ ,  $(a''_{37})^{(7)}(T_{37}, t)$ ,  $(a''_{38})^{(7)}(T_{37}, t)$  are first augmentation coefficients for category 1, 2 and 3

$(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$  are second augmentation coefficient for category 1, 2 and 3

$(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ ,  $(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ ,  $(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$  are third augmentation coefficient for category 1, 2 and 3

$(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ ,  $(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ ,  $(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficient for category 1, 2 and 3

$(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ ,  $(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ ,  $(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficient for category 1, 2 and 3

$(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ ,  $(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ ,  $(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficient for category 1, 2 and 3

$(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ ,  $(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ ,  $(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$  are seventh augmentation coefficient for category 1, 2 and 3

$(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ ,  $(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ ,  $(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$  are eighth augmentation coefficient for 1,2,3

$(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ ,  $(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ ,  $(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$  are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)} T_{37} - \left[ \begin{array}{ccc} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) & - (b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)} T_{36} - \left[ \begin{array}{ccc} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) & - (b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)} T_{37} - \left[ \begin{array}{ccc} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) & - (b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1)}(G, t) & - (b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) & - (b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where  $-(b''_{36})^{(7)}(G_{39}, t)$ ,  $-(b''_{37})^{(7)}(G_{39}, t)$ ,  $-(b''_{38})^{(7)}(G_{39}, t)$  are first detrition coefficients for

category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$  are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$  are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$  are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$  are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$  are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)$  are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$  are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$  are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

$$= (a_{40})^{(8)}G_{41} - \left[ \begin{array}{ccc} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) & + (a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40} - \left[ \begin{array}{ccc} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) & + (a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt}$$

$$= (a_{42})^{(8)}G_{41} - \left[ \begin{array}{ccc} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) & + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) & + (a''_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t) & + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where  $+(a''_{40})^{(8)}(T_{41}, t)$ ,  $+(a''_{41})^{(8)}(T_{41}, t)$ ,  $+(a''_{42})^{(8)}(T_{41}, t)$  are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$  are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ ,  $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ ,  $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$  are third

augmentation coefficient for category 1, 2 and 3

$$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$$

augmentation coefficient for category 1, 2 and 3

$$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$$

augmentation coefficient for category 1, 2 and 3

$$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t), +(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t), +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$$

augmentation coefficient for category 1, 2 and 3

$$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$$

augmentation coefficient for 1,2,3

$$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$$

augmentation coefficient for 1,2,3

$$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t), +(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$$

augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$

$$(b_{40})^{(8)}T_{41} - \left[ \begin{array}{ccc} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) & - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$

$$(b_{41})^{(8)}T_{40} - \left[ \begin{array}{ccc} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) & - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$

$$(b_{42})^{(8)}T_{41} - \left[ \begin{array}{ccc} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) & - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where  $-(b''_{36})^{(7)}(G_{39}, t), -(b''_{37})^{(7)}(G_{39}, t), -(b''_{38})^{(7)}(G_{39}, t)$  are first detrition coefficients for category 1, 2 and 3

$$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$$

are second detrition coefficients for category 1, 2 and 3

$$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$$

are third detrition coefficients for category 1, 2 and 3

$$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$$

are fourth detrition coefficients for category 1, 2 and 3

$$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$$

are fifth detrition coefficients for category 1, 2 and 3

$$-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t), -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t), -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$$

are sixth detrition coefficients

for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{38})^{(7,7)}(G_{39}, t)$  are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$  are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$  are ninth detrition coefficients for category 1, 2 and 3

$$\begin{aligned} \frac{dG_{44}}{dt} &= (a_{44})^{(9)}G_{45} \\ &- \left[ \begin{array}{l} (a''_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) + (a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13} \end{aligned}$$

$$\begin{aligned} \frac{dG_{45}}{dt} &= (a_{45})^{(9)}G_{44} \\ &- \left[ \begin{array}{l} (a''_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) + (a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14} \end{aligned}$$

$$\begin{aligned} \frac{dG_{46}}{dt} &= (a_{46})^{(9)}G_{45} \\ &- \left[ \begin{array}{l} (a''_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15} \end{aligned}$$

Where  $+(a''_{44})^{(9)}(T_{45}, t)$ ,  $+(a''_{45})^{(9)}(T_{45}, t)$ ,  $+(a''_{46})^{(9)}(T_{37}, t)$  are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$  are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ ,  $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ ,  $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$  are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ ,  $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ ,  $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ ,  $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ ,  $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ ,  $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ ,  $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ ,  $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ ,  $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$  are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)}$  are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t)}$  are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[ \begin{array}{ccc} \boxed{(b'_{44})^{(9)} - \boxed{(b''_{44})^{(9)}(G_{47}, t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{13}$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[ \begin{array}{ccc} \boxed{(b'_{45})^{(9)} - \boxed{(b''_{45})^{(9)}(G_{47}, t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{14}$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[ \begin{array}{ccc} \boxed{(b'_{46})^{(9)} - \boxed{(b''_{46})^{(9)}(G_{47}, t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{15}$$

Where  $\boxed{-(b''_{44})^{(9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{45})^{(9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{46})^{(9)}(G_{47}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)}$  are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)}$  are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)}$  are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)}$  are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)}$  are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)}$  are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i, j = 13, 14, 15 \quad 97$$

The functions  $(a''_i)^{(1)}, (b''_i)^{(1)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(1)}, (r_i)^{(1)}$ :

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)} \quad 98$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

**Definition of**  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$ :

Where  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$  are positive constants and  $i = 13, 14, 15$

They satisfy Lipschitz condition: 99

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T_{14} - T'_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(1)}(T'_{14}, t)$  and  $(a''_i)^{(1)}(T_{14}, t)$ .  $(T'_{14}, t)$  and  $(T_{14}, t)$  are points belonging to the interval  $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$ . It is to be noted that  $(a''_i)^{(1)}(T_{14}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{13})^{(1)} = 1$  then the function  $(a''_i)^{(1)}(T_{14}, t)$ , the first augmentation coefficient attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$ : 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$ , are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

**Definition of**  $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$ : 101

There exists two constants  $(\hat{P}_{13})^{(1)}$  and  $(\hat{Q}_{13})^{(1)}$  which together With  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$  and  $(\hat{B}_{13})^{(1)}$  and the constants  $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a'_i)^{(2)}, (a''_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (b''_i)^{(2)} > 0, \quad i, j = 16, 17, 18$$

The functions  $(a''_i)^{(2)}, (b''_i)^{(2)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(2)}, (r_i)^{(2)}$ :

$$(a''_i)^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \tag{102}$$

$$(b''_i)^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b'_i)^{(2)} \leq (\hat{B}_{16})^{(2)} \tag{103}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(2)}(T_{17}, t) = (p_i)^{(2)} \tag{104}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(2)}(G_{19}, t) = (r_i)^{(2)} \tag{105}$$

**Definition of**  $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$  : 106

Where  $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$  are positive constants and  $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a''_i)^{(2)}(T'_{17}, t) - (a''_i)^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T'_{17} - T_{17}| e^{-(\hat{M}_{16})^{(2)}t} \tag{107}$$

$$|(b''_i)^{(2)}((G_{19})', t) - (b''_i)^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G_{19})' - (G_{19})| e^{-(\hat{M}_{16})^{(2)}t} \tag{108}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(2)}(T'_{17}, t)$  and  $(a''_i)^{(2)}(T_{17}, t)$ .  $(T'_{17}, t)$  and  $(T_{17}, t)$  are points belonging to the interval  $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$ . It is to be noted that  $(a''_i)^{(2)}(T_{17}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{16})^{(2)} = 1$  then the function  $(a''_i)^{(2)}(T_{17}, t)$ , the first augmentation coefficient attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$  :

$$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, \text{ are positive constants} \tag{109}$$

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

**Definition of**  $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$  :

There exists two constants  $(\hat{P}_{16})^{(2)}$  and  $(\hat{Q}_{16})^{(2)}$  which together with  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$  and  $(\hat{B}_{16})^{(2)}$  and the constants  $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \tag{110}$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [ (b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)} ] < 1 \tag{111}$$

Where we suppose

$$(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22 \tag{112}$$

The functions  $(a''_i)^{(3)}, (b''_i)^{(3)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(3)}, (r_i)^{(3)}$ :

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \tag{113}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

**Definition of**  $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$  :

Where  $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$  are positive constants and  $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G_{23}', t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} ||G_{23} - G_{23}'|| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(3)}(T'_{21}, t)$  and  $(a''_i)^{(3)}(T_{21}, t)$ .  $(T'_{21}, t)$  and  $(T_{21}, t)$  are points belonging to the interval  $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$ . It is to be noted that  $(a''_i)^{(3)}(T_{21}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{20})^{(3)} = 1$  then the function  $(a''_i)^{(3)}(T_{21}, t)$ , the first augmentation coefficient attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$  : 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$ , are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants  $(\hat{P}_{20})^{(3)}$  and  $(\hat{Q}_{20})^{(3)}$  which together with  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$  and  $(\hat{B}_{20})^{(3)}$  and the constants  $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$ , satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [ (a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)} ] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [ (b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)} ] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 117$$

The functions  $(a''_i)^{(4)}, (b''_i)^{(4)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(4)}, (r_i)^{(4)}$ :

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

**Definition of**  $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$  :

Where  $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$  are positive constants and  $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25} - T'_{25}| e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b''_i)^{(4)}((G_{27})', t) - (b''_i)^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} |(G_{27}) - (G_{27})'| e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(4)}(T'_{25}, t)$  and  $(a''_i)^{(4)}(T_{25}, t)$ .  $(T'_{25}, t)$  and  $(T_{25}, t)$  are points belonging to the interval  $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$ . It is to be noted that  $(a''_i)^{(4)}(T_{25}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{24})^{(4)} = 1$  then the function  $(a''_i)^{(4)}(T_{25}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$  : 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$ , are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} + \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

**Definition of**  $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$  : 121

There exists two constants  $(\hat{P}_{24})^{(4)}$  and  $(\hat{Q}_{24})^{(4)}$  which together with  $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$  and  $(\hat{B}_{24})^{(4)}$  and the constants  $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, \quad i, j = 28, 29, 30 \tag{122}$$

The functions  $(a''_i)^{(5)}, (b''_i)^{(5)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(5)}, (r_i)^{(5)}$ :

$$\begin{aligned} (a''_i)^{(5)}(T_{29}, t) &\leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)} \\ (b''_i)^{(5)}((G_{31}), t) &\leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)} \\ \lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) &= (p_i)^{(5)} \\ \lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) &= (r_i)^{(5)} \end{aligned} \tag{123}$$

**Definition of**  $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$  :

Where  $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$  are positive constants and  $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$\begin{aligned} |(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| &\leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)}t} \\ |(b''_i)^{(5)}((G_{31})', t) - (b''_i)^{(5)}((G_{31}), t)| &< (\hat{k}_{28})^{(5)} |(G_{31}) - (G_{31})'| e^{-(\hat{M}_{28})^{(5)}t} \end{aligned}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(5)}(T'_{29}, t)$  and  $(a''_i)^{(5)}(T_{29}, t)$ .  $(T'_{29}, t)$  and  $(T_{29}, t)$  are points belonging to the interval  $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$ . It is to be noted that  $(a''_i)^{(5)}(T_{29}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{28})^{(5)} = 1$  then the function  $(a''_i)^{(5)}(T_{29}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$  : 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$ , are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}} , \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

**Definition of**  $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$  : 126

There exists two constants  $(\hat{P}_{28})^{(5)}$  and  $(\hat{Q}_{28})^{(5)}$  which together with  $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$  and  $(\hat{B}_{28})^{(5)}$  and the constants  $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$ , satisfy the inequalities

$$\begin{aligned} \frac{1}{(\hat{M}_{28})^{(5)}} [ (a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)} ] &< 1 \\ \frac{1}{(\hat{M}_{28})^{(5)}} [ (b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)} ] &< 1 \end{aligned}$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, \quad i, j = 32, 33, 34 \quad 127$$

The functions  $(a''_i)^{(6)}, (b''_i)^{(6)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(6)}, (r_i)^{(6)}$ :

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

**Definition of**  $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$  :

Where  $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$  are positive constants and  $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b''_i)^{(6)}((G_{35})', t) - (b''_i)^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} |(G_{35}) - (G_{35})'| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(6)}(T'_{33}, t)$  and  $(a''_i)^{(6)}(T_{33}, t)$ .  $(T'_{33}, t)$  and  $(T_{33}, t)$  are points belonging to the interval  $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$ . It is to be noted that  $(a''_i)^{(6)}(T_{33}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{32})^{(6)} = 1$  then the function  $(a''_i)^{(6)}(T_{33}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$  : 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$ , are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

**Definition of**  $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$  : 130

There exists two constants  $(\hat{P}_{32})^{(6)}$  and  $(\hat{Q}_{32})^{(6)}$  which together with  $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$  and  $(\hat{B}_{32})^{(6)}$  and the constants  $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [ (b_i)^{(6)} + (b'_i)^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)} ] < 1$$

Where we suppose

(G)  $(a_i)^{(7)}, (a'_i)^{(7)}, (a''_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (b''_i)^{(7)} > 0, \quad i, j = 36, 37, 38$  131

(H) The functions  $(a''_i)^{(7)}, (b''_i)^{(7)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(7)}, (r_i)^{(7)}$ :

$$(a''_i)^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b''_i)^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b'_i)^{(7)} \leq (\hat{B}_{36})^{(7)}$$

132

(I)  $\lim_{T_2 \rightarrow \infty} (a''_i)^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(J)

$$\lim_{G \rightarrow \infty} (b''_i)^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

**Definition of**  $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$  :

Where  $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}$  are positive constants and  $i = 36, 37, 38$

They satisfy Lipschitz condition:

133

$$|(a''_i)^{(7)}(T'_{37}, t) - (a''_i)^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T'_{37}| e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b''_i)^{(7)}((G_{39})', t) - (b''_i)^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} |(G_{39})' - (G_{39})| e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(7)}(T'_{37}, t)$  and  $(a''_i)^{(7)}(T_{37}, t)$ .  $(T'_{37}, t)$  and  $(T_{37}, t)$  are points belonging to the interval  $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$ . It is to be noted that  $(a''_i)^{(7)}(T_{37}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{36})^{(7)} = 1$  then the function  $(a''_i)^{(7)}(T_{37}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$  :

134

(K)  $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$ , are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

**Definition of**  $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$  :

135

(L) There exists two constants  $(\hat{P}_{36})^{(7)}$  and  $(\hat{Q}_{36})^{(7)}$  which together with  $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$  and  $(\hat{B}_{36})^{(7)}$  and the constants

$(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, \quad i, j = 40,41,42 \tag{136}$$

The functions  $(a''_i)^{(8)}, (b''_i)^{(8)}$  are positive continuous increasing and bounded

**Definition of**  $(p_i)^{(8)}, (r_i)^{(8)}$ : 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \tag{138}$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \tag{139}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \tag{140}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \tag{141}$$

**Definition of**  $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$  :

Where  $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$  are positive constants and  $i = 40,41,42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T'_{41}| e^{-(\hat{M}_{40})^{(8)}t} \tag{142}$$

$$|(b''_i)^{(8)}((G_{43})', t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} |(G_{43}) - (G_{43})'| e^{-(\hat{M}_{40})^{(8)}t} \tag{143}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(8)}(T'_{41}, t)$  and  $(a''_i)^{(8)}(T_{41}, t)$ .  $(T'_{41}, t)$  and  $(T_{41}, t)$  are points belonging to the interval  $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$ . It is to be noted that  $(a''_i)^{(8)}(T_{41}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{40})^{(8)} = 1$  then the function  $(a''_i)^{(8)}(T_{41}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$  :

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$ , are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}} , \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \tag{144}$$

**Definition of**  $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$  :

There exists two constants  $(\hat{P}_{40})^{(8)}$  and  $(\hat{Q}_{40})^{(8)}$  which together with  $(\hat{M}_{40})^{(8)}$ ,  $(\hat{k}_{40})^{(8)}$ ,  $(\hat{A}_{40})^{(8)}$ ,  $(\hat{B}_{40})^{(8)}$  and the constants  $(a_i)^{(8)}$ ,  $(a'_i)^{(8)}$ ,  $(b_i)^{(8)}$ ,  $(b'_i)^{(8)}$ ,  $(p_i)^{(8)}$ ,  $(r_i)^{(8)}$ ,  $i = 40,41,42$ ,

Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a'_i)^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \tag{145}$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b'_i)^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \tag{146}$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, \quad i, j = 44,45,46 \tag{146}$$

A

The functions  $(a''_i)^{(9)}$ ,  $(b''_i)^{(9)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(9)}$ ,  $(r_i)^{(9)}$ :

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

**Definition of**  $(\hat{A}_{44})^{(9)}$ ,  $(\hat{B}_{44})^{(9)}$  :

Where  $(\hat{A}_{44})^{(9)}$ ,  $(\hat{B}_{44})^{(9)}$ ,  $(p_i)^{(9)}$ ,  $(r_i)^{(9)}$  are positive constants and  $i = 44,45,46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T'_{45} - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G_{47})', t) - (b''_i)^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(9)}(T'_{45}, t)$  and  $(a''_i)^{(9)}(T_{45}, t)$ .  $(T'_{45}, t)$  and  $(T_{45}, t)$  are points belonging to the interval  $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$ . It is to be noted that  $(a''_i)^{(9)}(T_{45}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{44})^{(9)} = 1$  then the function  $(a''_i)^{(9)}(T_{45}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{44})^{(9)}$ ,  $(\hat{k}_{44})^{(9)}$  :

$(\hat{M}_{44})^{(9)}$ ,  $(\hat{k}_{44})^{(9)}$ , are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

**Definition of**  $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$  :

There exists two constants  $(\hat{P}_{44})^{(9)}$  and  $(\hat{Q}_{44})^{(9)}$  which together with  $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$  and  $(\hat{B}_{44})^{(9)}$  and the constants  $(a_i)^{(9)}, (a'_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46,$  satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a'_i)^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b'_i)^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

**Theorem 1:** if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 2 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

**Definition of**  $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

**Theorem 3 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

**Theorem 4 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 5 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 6 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 7:** if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 8:** if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 9:** if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Proof:** Consider operator  $\mathcal{A}^{(1)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy 154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[ (a_{13})^{(1)} G_{14}(s_{(13)}) - \left( (a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[ (a_{14})^{(1)} G_{13}(s_{(13)}) - \left( (a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[ (a_{15})^{(1)} G_{14}(s_{(13)}) - \left( (a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[ (b_{13})^{(1)} T_{14}(s_{(13)}) - \left( (b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[ (b_{14})^{(1)} T_{13}(s_{(13)}) - \left( (b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[ (b_{15})^{(1)} T_{14}(s_{(13)}) - \left( (b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where  $s_{(13)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

159

Consider operator  $\mathcal{A}^{(2)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[ (a_{16})^{(2)} G_{17}(s_{(16)}) - \left( (a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[ (a_{17})^{(2)} G_{16}(s_{(16)}) - \left( (a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[ (a_{18})^{(2)} G_{17}(s_{(16)}) - \left( (a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[ (b_{16})^{(2)} T_{17}(s_{(16)}) - \left( (b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[ (b_{17})^{(2)} T_{16}(s_{(16)}) - \left( (b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[ (b_{18})^{(2)} T_{17}(s_{(16)}) - \left( (b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where  $s_{(16)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(3)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[ (a_{20})^{(3)} G_{21}(s_{(20)}) - \left( (a'_{20})^{(3)} + a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[ (a_{21})^{(3)} G_{20}(s_{(20)}) - \left( (a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[ (a_{22})^{(3)} G_{21}(s_{(20)}) - \left( (a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[ (b_{20})^{(3)} T_{21}(s_{(20)}) - \left( (b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[ (b_{21})^{(3)} T_{20}(s_{(20)}) - \left( (b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[ (b_{22})^{(3)} T_{21}(s_{(20)}) - \left( (b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where  $s_{(20)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:** Consider operator  $\mathcal{A}^{(4)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[ (a_{24})^{(4)} G_{25}(s_{(24)}) - \left( (a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[ (a_{25})^{(4)} G_{24}(s_{(24)}) - \left( (a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[ (a_{26})^{(4)} G_{25}(s_{(24)}) - \left( (a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[ (b_{24})^{(4)} T_{25}(s_{(24)}) - \left( (b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[ (b_{25})^{(4)} T_{24}(s_{(24)}) - \left( (b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[ (b_{26})^{(4)} T_{25}(s_{(24)}) - \left( (b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where  $s_{(24)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:** Consider operator  $\mathcal{A}^{(5)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[ (a_{28})^{(5)} G_{29}(s_{(28)}) - \left( (a'_{28})^{(5)} + a''_{28}(s_{(28)}, s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[ (a_{29})^{(5)} G_{28}(s_{(28)}) - \left( (a'_{29})^{(5)} + a''_{29}(s_{(28)}, s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[ (a_{30})^{(5)} G_{29}(s_{(28)}) - \left( (a'_{30})^{(5)} + a''_{30}(s_{(28)}, s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[ (b_{28})^{(5)} T_{29}(s_{(28)}) - \left( (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}, s_{(28)})) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[ (b_{29})^{(5)} T_{28}(s_{(28)}) - \left( (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}, s_{(28)})) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[ (b_{30})^{(5)} T_{29}(s_{(28)}) - \left( (b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}, s_{(28)})) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where  $s_{(28)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(6)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[ (a_{32})^{(6)} G_{33}(s_{(32)}) - \left( (a'_{32})^{(6)} + a''_{32}(s_{(32)}, s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[ (a_{33})^{(6)} G_{32}(s_{(32)}) - \left( (a'_{33})^{(6)} + a''_{33}(s_{(32)}, s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[ (a_{34})^{(6)} G_{33}(s_{(32)}) - \left( (a'_{34})^{(6)} + a''_{34}(s_{(32)}, s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[ (b_{32})^{(6)} T_{33}(s_{(32)}) - \left( (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}, s_{(32)})) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[ (b_{33})^{(6)} T_{32}(s_{(32)}) - \left( (b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[ (b_{34})^{(6)} T_{33}(s_{(32)}) - \left( (b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where  $s_{(32)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(7)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[ (a_{36})^{(7)} G_{37}(s_{(36)}) - \left( (a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[ (a_{37})^{(7)} G_{36}(s_{(36)}) - \left( (a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[ (a_{38})^{(7)} G_{37}(s_{(36)}) - \left( (a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[ (b_{36})^{(7)} T_{37}(s_{(36)}) - \left( (b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[ (b_{37})^{(7)} T_{36}(s_{(36)}) - \left( (b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[ (b_{38})^{(7)} T_{37}(s_{(36)}) - \left( (b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where  $s_{(36)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(8)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(M_{40})^{(8)}t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[ (a_{40})^{(8)} G_{41}(s_{(40)}) - \left( (a'_{40})^{(8)} + (a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[ (a_{41})^{(8)} G_{40}(s_{(40)}) - \left( (a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[ (a_{42})^{(8)} G_{41}(s_{(40)}) - \left( (a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[ (b_{40})^{(8)} T_{41}(s_{(40)}) - \left( (b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[ (b_{41})^{(8)} T_{40}(s_{(40)}) - \left( (b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[ (b_{42})^{(8)} T_{41}(s_{(40)}) - \left( (b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where  $s_{(40)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

166

Consider operator  $\mathcal{A}^{(9)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(M_{44})^{(9)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(M_{44})^{(9)}t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[ (a_{44})^{(9)} G_{45}(s_{(44)}) - \left( (a'_{44})^{(9)} + (a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[ (a_{45})^{(9)} G_{44}(s_{(44)}) - \left( (a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[ (a_{46})^{(9)} G_{45}(s_{(44)}) - \left( (a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[ (b_{44})^{(9)} T_{45}(s_{(44)}) - \left( (b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[ (b_{45})^{(9)} T_{44}(s_{(44)}) - \left( (b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[ (b_{46})^{(9)} T_{45}(s_{(44)}) - \left( (b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where  $s_{(44)}$  is the integrand that is integrated over an interval  $(0, t)$

The operator  $\mathcal{A}^{(1)}$  maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t [(a_{13})^{(1)} (G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}})] ds_{(13)} =$$

$$(1 + (a_{13})^{(1)} t) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} (e^{(\hat{M}_{13})^{(1)} t} - 1)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[ ((\hat{P}_{13})^{(1)} + G_{14}^0) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 1

Analogous inequalities hold also for  $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator  $\mathcal{A}^{(2)}$  maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t [(a_{16})^{(2)} (G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} s_{(16)}})] ds_{(16)} =$$

$$(1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} (e^{(\hat{M}_{16})^{(2)} t} - 1)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[ ((\hat{P}_{16})^{(2)} + G_{17}^0) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for  $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator  $\mathcal{A}^{(3)}$  maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t [(a_{20})^{(3)} (G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}})] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} (e^{(\hat{M}_{20})^{(3)} t} - 1)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[ ((\hat{P}_{20})^{(3)} + G_{21}^0) e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for  $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator  $\mathcal{A}^{(4)}$  maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t [(a_{24})^{(4)} (G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}})] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} (e^{(\hat{M}_{24})^{(4)} t} - 1)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[ ((\hat{P}_{24})^{(4)} + G_{25}^0)e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)} \right]$$

$(G_t^0)$  is as defined in the statement of theorem 4

The operator  $\mathcal{A}^{(5)}$  maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t [(a_{28})^{(5)} (G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}s_{(28)}})] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} (e^{(\hat{M}_{28})^{(5)}t} - 1)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[ ((\hat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

$(G_t^0)$  is as defined in the statement of theorem 5

The operator  $\mathcal{A}^{(6)}$  maps the space of functions satisfying Equations into itself .Indeed it is obvious that

176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t [(a_{32})^{(6)} (G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}s_{(32)}})] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} (e^{(\hat{M}_{32})^{(6)}t} - 1)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[ ((\hat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

$(G_t^0)$  is as defined in the statement of theorem 6

Analogous inequalities hold also for  $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(b) The operator  $\mathcal{A}^{(7)}$  maps the space of functions satisfying Equations into itself .Indeed it is obvious that

178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t [(a_{36})^{(7)} (G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}})] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} (e^{(\hat{M}_{36})^{(7)}t} - 1)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[ ((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 7

The operator  $\mathcal{A}^{(8)}$  maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[ (a_{40})^{(8)} \left( G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} = \tag{180}$$

$$\left( 1 + (a_{40})^{(8)} t \right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left( e^{(\hat{M}_{40})^{(8)} t} - 1 \right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[ \left( (\hat{P}_{40})^{(8)} + G_{41}^0 \right) e^{\left( -\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0} \right)} + (\hat{P}_{40})^{(8)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 8

Analogous inequalities hold also for  $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator  $\mathcal{A}^{(9)}$  maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[ (a_{44})^{(9)} \left( G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$\left( 1 + (a_{44})^{(9)} t \right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left( e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[ \left( (\hat{P}_{44})^{(9)} + G_{45}^0 \right) e^{\left( -\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0} \right)} + (\hat{P}_{44})^{(9)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 9

Analogous inequalities hold also for  $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take  $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$  and to choose 182

$(\hat{P}_{13})^{(1)}$  and  $(\hat{Q}_{13})^{(1)}$  large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[ (\hat{P}_{13})^{(1)} + \left( (\hat{P}_{13})^{(1)} + G_j^0 \right) e^{-\left( \frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{13})^{(1)} \tag{183}$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[ \left( (\hat{Q}_{13})^{(1)} + T_j^0 \right) e^{-\left( \frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \tag{184}$$

In order that the operator  $\mathcal{A}^{(1)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(1)}$  is a contraction with respect to the metric 185

$$d \left( (G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{13})^{(1)}t} \}$$

Indeed if we denote

**Definition of  $\tilde{G}, \tilde{T} : (\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$**

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{ (a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} | (a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)}) | e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} \} ds_{(13)} \end{aligned}$$

Where  $s_{(13)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)}t} &\leq \\ \frac{1}{(\widehat{M}_{13})^{(1)}} &((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d \left( (G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}) \right) \end{aligned}$$

186

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 1:** The fact that we supposed  $(a''_{13})^{(1)}$  and  $(b''_{13})^{(1)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$  and  $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(1)}$  and  $(b''_i)^{(1)}, i = 13, 14, 15$  depend only on  $T_{14}$  and respectively on  $G$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t ((a'_i)^{(1)} - (a''_i)^{(1)}(T_{14}(s_{(13)}), s_{(13)})) ds_{(13)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)}t} > 0 \text{ for } t > 0$$

**Definition of  $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$  and  $((\widehat{M}_{13})^{(1)})_3$ :**

187

**Remark 3:** if  $G_{13}$  is bounded, the same property have also  $G_{14}$  and  $G_{15}$ . indeed if

$$G_{13} < (\widehat{M}_{13})^{(1)} \text{ it follows } \frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If  $G_{14}$  or  $G_{15}$  is bounded, the same property follows for  $G_{13}$  ,  $G_{15}$  and  $G_{13}$  ,  $G_{14}$  respectively.

**Remark 4:** If  $G_{13}$  is bounded, from below, the same property holds for  $G_{14}$  and  $G_{15}$  . The proof is analogous with the preceding one. An analogous property is true if  $G_{14}$  is bounded from below. 188

**Remark 5:** If  $T_{13}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b'_{14})^{(1)}$  then  $T_{14} \rightarrow \infty$ . 189

**Definition of**  $(m)^{(1)}$  and  $\varepsilon_1$  :

Indeed let  $t_1$  be so that for  $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then  $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$  which leads to

$$T_{14} \geq \left( \frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left( \frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for  $T_{15}$  if  $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take  $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$  and to choose 190

$(\widehat{P}_{16})^{(2)}$  and  $(\widehat{Q}_{16})^{(2)}$  large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} \left[ (\widehat{P}_{16})^{(2)} + ((\widehat{P}_{16})^{(2)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} \left[ ((\widehat{Q}_{16})^{(2)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{16})^{(2)} \right] \leq (\widehat{Q}_{16})^{(2)} \quad 192$$

In order that the operator  $\mathcal{A}^{(2)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself 193

The operator  $\mathcal{A}^{(2)}$  is a contraction with respect to the metric 194

$$d\left( ((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote

195

**Definition of**  $\widetilde{G}_{19}, \widetilde{T}_{19} : (\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results

196

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} | (a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)}) | e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} \} ds_{(16)} \end{aligned}$$

Where  $s_{(16)}$  represents integrand that is integrated over the interval  $[0, t]$

197

From the hypotheses it follows

$$\begin{aligned} |(\widetilde{G}_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} &\leq \\ \frac{1}{(\bar{M}_{16})^{(2)}} &((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)}) d \left( ((G_{19})^{(1)}, (T_{19})^{(1)}); (G_{19})^{(2)}, (T_{19})^{(2)} \right) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

198

**Remark 6:** The fact that we supposed  $(a''_{16})^{(2)}$  and  $(b''_{16})^{(2)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$  and  $(\widehat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$  respectively of  $\mathbb{R}_+$ .

199

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(2)}$  and  $(b''_i)^{(2)}$ ,  $i = 16, 17, 18$  depend only on  $T_{17}$  and respectively on  $(G_{19})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 7:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

200

it results

$$G_i(t) \geq G_i^0 e^{\left[ - \int_0^t \{ (a'_i)^{(2)} - (a''_i)^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \} ds_{(16)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(2)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{16})^{(2)})_1, ((\widehat{M}_{16})^{(2)})_2$  and  $((\widehat{M}_{16})^{(2)})_3 :$

201

**Remark 8:** if  $G_{16}$  is bounded, the same property have also  $G_{17}$  and  $G_{18}$ . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$  it follows  $\frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a'_{17})^{(2)} G_{17}$  and by integrating

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a'_{17})^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a'_{18})^{(2)}$$

If  $G_{17}$  or  $G_{18}$  is bounded, the same property follows for  $G_{16}$  ,  $G_{18}$  and  $G_{16}$  ,  $G_{17}$  respectively.

**Remark 9:** If  $G_{16}$  is bounded, from below, the same property holds for  $G_{17}$  and  $G_{18}$  . The proof is 202  
 analogous with the preceding one. An analogous property is true if  $G_{17}$  is bounded from below.

**Remark 10:** If  $T_{16}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b'_i)^{(2)} ((G_{19})(t), t)) = (b'_{17})^{(2)}$  then  $T_{17} \rightarrow \infty$ . 203

**Definition of**  $(m)^{(2)}$  and  $\varepsilon_2$  :

Indeed let  $t_2$  be so that for  $t > t_2$

$$(b_{17})^{(2)} - (b'_i)^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then  $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$  which leads to 204

$$T_{17} \geq \left( \frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$$

If we take  $t$  such that  $e^{-\varepsilon_2 t} = \frac{1}{2}$  it results

$$T_{17} \geq \left( \frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_2}$$

By taking now  $\varepsilon_2$  sufficiently small one sees that  $T_{17}$  is unbounded. 205

The same property holds for  $T_{18}$  if  $\lim_{t \rightarrow \infty} (b''_{18})^{(2)} ((G_{19})(t), t) = (b'_{18})^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take  $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$  and to choose 207

$(\widehat{P}_{20})^{(3)}$  and  $(\widehat{Q}_{20})^{(3)}$  large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} \left[ (\widehat{P}_{20})^{(3)} + ((\widehat{P}_{20})^{(3)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{20})^{(3)}$$

208

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} \left[ ((\widehat{Q}_{20})^{(3)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{20})^{(3)} \right] \leq (\widehat{Q}_{20})^{(3)}$$

209

In order that the operator  $\mathcal{A}^{(3)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying 210  
 Equations into itself

The operator  $\mathcal{A}^{(3)}$  is a contraction with respect to the metric 211

$$d \left( ((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \}$$

Indeed if we denote

212

**Definition of**  $\widetilde{G}_{23}, \widetilde{T}_{23} : ( (\widetilde{G}_{23}), (\widetilde{T}_{23}) ) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results

213

$$\begin{aligned} |\widetilde{G}_{20}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} | (a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)}) | e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned}$$

Where  $s_{(20)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \\ \frac{1}{(\bar{M}_{20})^{(3)}} &((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)} (\widehat{k}_{20})^{(3)}) d \left( ((G_{23})^{(1)}, (T_{23})^{(1)}); (G_{23})^{(2)}, (T_{23})^{(2)} \right) \end{aligned}$$

214

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 11:** The fact that we supposed  $(a''_{20})^{(3)}$  and  $(b''_{20})^{(3)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$  and  $(\widehat{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$  respectively of  $\mathbb{R}_+$ .

215

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(3)}$  and  $(b''_i)^{(3)}$ ,  $i = 20, 21, 22$  depend only on  $T_{21}$  and respectively on  $(G_{23})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 12:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

216

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \} ds_{(20)}]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(3)}t} > 0 \text{ for } t > 0 \end{aligned}$$

**Definition of**  $((\bar{M}_{20})^{(3)})_1, ((\bar{M}_{20})^{(3)})_2$  and  $((\bar{M}_{20})^{(3)})_3 :$

217

**Remark 13:** if  $G_{20}$  is bounded, the same property have also  $G_{21}$  and  $G_{22}$  . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$  it follows  $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)}G_{21}$  and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If  $G_{21}$  or  $G_{22}$  is bounded, the same property follows for  $G_{20}$  ,  $G_{22}$  and  $G_{20}$  ,  $G_{21}$  respectively.

**Remark 14:** If  $G_{20}$  is bounded, from below, the same property holds for  $G_{21}$  and  $G_{22}$  . The proof is analogous with the preceding one. An analogous property is true if  $G_{21}$  is bounded from below. 218

**Remark 15:** If  $T_{20}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b'_i)^{(3)} ((G_{23})(t), t)) = (b'_{21})^{(3)}$  then  $T_{21} \rightarrow \infty$ . 219

**Definition of**  $(m)^{(3)}$  and  $\varepsilon_3$  :

Indeed let  $t_3$  be so that for  $t > t_3$

$$(b_{21})^{(3)} - (b'_i)^{(3)} ((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then  $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$  which leads to 220

$$T_{21} \geq \left( \frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$$

If we take  $t$  such that  $e^{-\varepsilon_3 t} = \frac{1}{2}$  it results

$$T_{21} \geq \left( \frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3}$$

By taking now  $\varepsilon_3$  sufficiently small one sees that  $T_{21}$  is unbounded.

The same property holds for  $T_{22}$  if  $\lim_{t \rightarrow \infty} (b''_{22})^{(3)} ((G_{23})(t), t) = (b'_{22})^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take  $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$  and to choose 221

$(\widehat{P}_{24})^{(4)}$  and  $(\widehat{Q}_{24})^{(4)}$  large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \left[ (\widehat{P}_{24})^{(4)} + ((\widehat{P}_{24})^{(4)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \left[ ((\widehat{Q}_{24})^{(4)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{24})^{(4)} \right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator  $\mathcal{A}^{(4)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself 224

The operator  $\mathcal{A}^{(4)}$  is a contraction with respect to the metric 225

$$d\left(\left((G_{27})^{(1)}, (T_{27})^{(1)}\right), \left((G_{27})^{(2)}, (T_{27})^{(2)}\right)\right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{24})^{(4)}t} \}$$

Indeed if we denote

**Definition of**  $(\widehat{G}_{27}), (\widehat{T}_{27}) : ((\widehat{G}_{27}), (\widehat{T}_{27})) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\widetilde{G}_{24}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} ds_{(24)} + \\ &\int_0^t \{ (a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} + \\ &G_{24}^{(2)} | (a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)}) | e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} e^{(\widehat{M}_{24})^{(4)}s_{(24)}} \} ds_{(24)} \end{aligned}$$

Where  $s_{(24)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)}t} &\leq \frac{1}{(\widehat{M}_{24})^{(4)}} \left( (a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)} (\widehat{K}_{24})^{(4)} \right) d\left(\left((G_{27})^{(1)}, (T_{27})^{(1)}\right), \left((G_{27})^{(2)}, (T_{27})^{(2)}\right)\right) \end{aligned} \tag{226}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 16:** The fact that we supposed  $(a''_{24})^{(4)}$  and  $(b''_{24})^{(4)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$  and  $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$  respectively of  $\mathbb{R}_+$ . 227

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a'_i)^{(4)}$  and  $(b'_i)^{(4)}$ ,  $i = 24, 25, 26$  depend only on  $T_{25}$  and respectively on  $(G_{27})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 17:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  228

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[ - \int_0^t \{ (a'_i)^{(4)} - (a''_i)^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \} ds_{(24)} \right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(4)}t} > 0 \text{ for } t > 0 \end{aligned}$$

**Definition of**  $(\widehat{M}_{24})^{(4)}_1, (\widehat{M}_{24})^{(4)}_2$  and  $(\widehat{M}_{24})^{(4)}_3 :$  229

**Remark 18:** if  $G_{24}$  is bounded, the same property have also  $G_{25}$  and  $G_{26}$ . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$  it follows  $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$  and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If  $G_{25}$  or  $G_{26}$  is bounded, the same property follows for  $G_{24}$ ,  $G_{26}$  and  $G_{24}$ ,  $G_{25}$  respectively.

**Remark 19:** If  $G_{24}$  is bounded, from below, the same property holds for  $G_{25}$  and  $G_{26}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{25}$  is bounded from below. 230

**Remark 20:** If  $T_{24}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b'_i)^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$  then  $T_{25} \rightarrow \infty$ . 231

**Definition of**  $(m)^{(4)}$  and  $\varepsilon_4$ :

Indeed let  $t_4$  be so that for  $t > t_4$

$$(b_{25})^{(4)} - (b'_i)^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then  $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$  which leads to 232

$$T_{25} \geq \left( \frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$$

If we take  $t$  such that  $e^{-\varepsilon_4 t} = \frac{1}{2}$  it results

$$T_{25} \geq \left( \frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4}$$

By taking now  $\varepsilon_4$  sufficiently small one sees that  $T_{25}$  is unbounded.

The same property holds for  $T_{26}$  if  $\lim_{t \rightarrow \infty} (b''_{26})^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for  $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take  $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$  and to choose 233

$(\widehat{P}_{28})^{(5)}$  and  $(\widehat{Q}_{28})^{(5)}$  large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} \left[ (\widehat{P}_{28})^{(5)} + ((\widehat{P}_{28})^{(5)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{28})^{(5)}$$
234

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} \left[ ((\widehat{Q}_{28})^{(5)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{28})^{(5)} \right] \leq (\widehat{Q}_{28})^{(5)}$$
235

In order that the operator  $\mathcal{A}^{(5)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(5)}$  is a contraction with respect to the metric

236

$$d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)}\right), \left((G_{31})^{(2)}, (T_{31})^{(2)}\right)\right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{28})^{(5)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{28})^{(5)}t} \}$$

Indeed if we denote

**Definition of**  $(\widehat{G}_{31}), (\widehat{T}_{31}) : (\widehat{G}_{31}), (\widehat{T}_{31}) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\widetilde{G}_{28}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} e^{(\widehat{M}_{28})^{(5)}s_{(28)}} ds_{(28)} + \\ &\int_0^t \{(a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} e^{(\widehat{M}_{28})^{(5)}s_{(28)}} + \\ &G_{28}^{(2)} |(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\} ds_{(28)} \end{aligned}$$

Where  $s_{(28)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)}t} &\leq \\ \frac{1}{(\widehat{M}_{28})^{(5)}} &\left( (a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)} \right) d\left(\left((G_{31})^{(1)}, (T_{31})^{(1)}\right); (G_{31})^{(2)}, (T_{31})^{(2)}\right) \end{aligned} \tag{237}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 21:** The fact that we supposed  $(a''_{28})^{(5)}$  and  $(b''_{28})^{(5)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$  and  $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$  respectively of  $\mathbb{R}_+$ . 238

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(5)}$  and  $(b''_i)^{(5)}, i = 28, 29, 30$  depend only on  $T_{29}$  and respectively on  $(G_{31})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 22:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  239

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(5)}t} > 0 \text{ for } t > 0 \end{aligned}$$

**Definition of**  $(\widehat{M}_{28})^{(5)}_1, (\widehat{M}_{28})^{(5)}_2$  and  $(\widehat{M}_{28})^{(5)}_3 :$  240

**Remark 23:** if  $G_{28}$  is bounded, the same property have also  $G_{29}$  and  $G_{30}$ . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)}G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If  $G_{29}$  or  $G_{30}$  is bounded, the same property follows for  $G_{28}$ ,  $G_{30}$  and  $G_{28}$ ,  $G_{29}$  respectively.

**Remark 24:** If  $G_{28}$  is bounded, from below, the same property holds for  $G_{29}$  and  $G_{30}$ . The proof is 241  
 analogous with the preceding one. An analogous property is true if  $G_{29}$  is bounded from below.

**Remark 25:** If  $T_{28}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b'_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$  then  $T_{29} \rightarrow \infty$ . 242

**Definition of**  $(m)^{(5)}$  and  $\varepsilon_5$  :

Indeed let  $t_5$  be so that for  $t > t_5$

$$(b_{29})^{(5)} - (b'_i)^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then  $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$  which leads to 243

$$T_{29} \geq \left( \frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left( \frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for  $T_{30}$  if  $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for  $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take  $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$  and to choose 244

$(\widehat{P}_{32})^{(6)}$  and  $(\widehat{Q}_{32})^{(6)}$  large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[ (\widehat{P}_{32})^{(6)} + ((\widehat{P}_{32})^{(6)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[ ((\widehat{Q}_{32})^{(6)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{32})^{(6)} \right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator  $\mathcal{A}^{(6)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(6)}$  is a contraction with respect to the metric

247

$$d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}\right), \left((G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{32})^{(6)}t} \right\}$$

Indeed if we denote

**Definition of**  $(\widehat{G}_{35}), (\widehat{T}_{35}) : ((\widehat{G}_{35}), (\widehat{T}_{35})) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\widetilde{G}_{32}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} e^{(\widehat{M}_{32})^{(6)}s_{(32)}} ds_{(32)} + \\ &\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} + \\ &(a''_{32})^{(6)}(T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} e^{(\widehat{M}_{32})^{(6)}s_{(32)}} + \\ &G_{32}^{(2)} |(a''_{32})^{(6)}(T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)}(T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\} ds_{(32)} \end{aligned}$$

Where  $s_{(32)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)}t} &\leq \\ \frac{1}{(\widehat{M}_{32})^{(6)}} &\left( (a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)} \right) d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}\right), \left((G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) \end{aligned}$$

248

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 26:** The fact that we supposed  $(a''_{32})^{(6)}$  and  $(b''_{32})^{(6)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$  and  $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$  respectively of  $\mathbb{R}_+$ .

249

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a'_i)^{(6)}$  and  $(b'_i)^{(6)}$ ,  $i = 32, 33, 34$  depend only on  $T_{33}$  and respectively on  $(G_{35})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 27:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

250

it results

$$\begin{aligned} G_i(t) &\geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(6)} - (a''_i)^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0 \\ T_i(t) &\geq T_i^0 e^{-(b'_i)^{(6)}t} > 0 \text{ for } t > 0 \end{aligned}$$

**Definition of**  $(\widehat{M}_{32})^{(6)}_1, (\widehat{M}_{32})^{(6)}_2$  and  $(\widehat{M}_{32})^{(6)}_3 :$

251

**Remark 28:** if  $G_{32}$  is bounded, the same property have also  $G_{33}$  and  $G_{34}$  . indeed if

$$G_{32} < (\widehat{M}_{32})^{(6)} \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If  $G_{33}$  or  $G_{34}$  is bounded, the same property follows for  $G_{32}$  ,  $G_{34}$  and  $G_{32}$  ,  $G_{33}$  respectively.

**Remark 29:** If  $G_{32}$  is bounded, from below, the same property holds for  $G_{33}$  and  $G_{34}$  . The proof is 252  
 analogous with the preceding one. An analogous property is true if  $G_{33}$  is bounded from below.

**Remark 30:** If  $T_{32}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(6)} ((G_{35})(t), t)) = (b'_{33})^{(6)}$  then  $T_{33} \rightarrow \infty$ . 253

**Definition of**  $(m)^{(6)}$  and  $\varepsilon_6$  :

Indeed let  $t_6$  be so that for  $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then  $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$  which leads to 254

$$T_{33} \geq \left( \frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$$T_{33} \geq \left( \frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6} \text{ By taking now } \varepsilon_6 \text{ sufficiently small one sees that } T_{33} \text{ is unbounded.}$$

The same property holds for  $T_{34}$  if  $\lim_{t \rightarrow \infty} (b_{34}'')^{(6)} ((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for  $G_{37}$  ,  $G_{38}$  ,  $T_{36}$  ,  $T_{37}$  ,  $T_{38}$  255

It is now sufficient to take  $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$  and to choose

$(\widehat{P}_{36})^{(7)}$  and  $(\widehat{Q}_{36})^{(7)}$  large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[ (\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[ ((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator  $\mathcal{A}^{(7)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying

Equations into itself

The operator  $\mathcal{A}^{(7)}$  is a contraction with respect to the metric

258

$$d\left(\left((G_{39})^{(1)}, (T_{39})^{(1)}\right), \left((G_{39})^{(2)}, (T_{39})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

**Definition of**  $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : ((\widetilde{G}_{39}), (\widetilde{T}_{39})) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{(a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{-(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} |(a''_{36})^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)} (T_{37}^{(2)}, s_{(36)})| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}}\} ds_{(36)} \end{aligned}$$

Where  $s_{(36)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)}t} &\leq \\ \frac{1}{(\bar{M}_{36})^{(7)}} &\left( (a_{36})^{(7)} + (a'_{36})^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)} \right) d\left(\left((G_{39})^{(1)}, (T_{39})^{(1)}\right); (G_{39})^{(2)}, (T_{39})^{(2)}\right) \end{aligned}$$

259

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 31:** The fact that we supposed  $(a''_{36})^{(7)}$  and  $(b''_{36})^{(7)}$  depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$  and  $(\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$  respectively of  $\mathbb{R}_+$ .

260

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(7)}$  and  $(b''_i)^{(7)}$ ,  $i = 36, 37, 38$  depend only on  $T_{37}$  and respectively on  $(G_{39})$  (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 32:** There does not exist any t where  $G_i(t) = 0$  and  $T_i(t) = 0$

261

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(7)} - (a''_i)^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{36})^{(7)})_1, ((\widehat{M}_{36})^{(7)})_2$  and  $((\widehat{M}_{36})^{(7)})_3$  : 262

**Remark 33:** if  $G_{36}$  is bounded, the same property have also  $G_{37}$  and  $G_{38}$  . indeed if

$G_{36} < ((\widehat{M}_{36})^{(7)})$  it follows  $\frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a'_{37})^{(7)}G_{37}$  and by integrating

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If  $G_{37}$  or  $G_{38}$  is bounded, the same property follows for  $G_{36}$  ,  $G_{38}$  and  $G_{36}$  ,  $G_{37}$  respectively.

**Remark 34:** If  $G_{36}$  is bounded, from below, the same property holds for  $G_{37}$  and  $G_{38}$  . The proof is 263  
analogous with the preceding one. An analogous property is true if  $G_{37}$  is bounded from below.

**Remark 35:** If  $T_{36}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(7)}((G_{39})(t), t)) = (b'_{37})^{(7)}$  then  $T_{37} \rightarrow \infty$ . 264

**Definition of**  $(m)^{(7)}$  and  $\varepsilon_7$  :

Indeed let  $t_7$  be so that for  $t > t_7$

$$(b_{37})^{(7)} - (b''_i)^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then  $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$  which leads to 265

$$T_{37} \geq \left( \frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$$

If we take  $t$  such that  $e^{-\varepsilon_7 t} = \frac{1}{2}$  it results

$$T_{37} \geq \left( \frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7}$$

By taking now  $\varepsilon_7$  sufficiently small one sees that  $T_{37}$  is unbounded.

The same property holds for  $T_{38}$  if  $\lim_{t \rightarrow \infty} (b''_i)^{(7)}((G_{39})(t), t) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take  $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$  and to choose 266

$(\widehat{P}_{40})^{(8)}$  and  $(\widehat{Q}_{40})^{(8)}$  large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[ (\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)}$$
267

$$\frac{(b_i)^{(8)}}{(\bar{M}_{40})^{(8)}} \left[ ((\hat{Q}_{40})^{(8)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{40})^{(8)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)} \right] \leq (\hat{Q}_{40})^{(8)}$$

In order that the operator  $\mathcal{A}^{(8)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(8)}$  is a contraction with respect to the metric

$$d\left( (G_{43})^{(1)}, (T_{43})^{(1)}, (G_{43})^{(2)}, (T_{43})^{(2)} \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \right\} \tag{269}$$

Indeed if we denote 270

**Definition of**  $(\widehat{G_{43}}, \widehat{T_{43}})$  :  $(\widehat{G_{43}}, \widehat{T_{43}}) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t (a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ &G_{40}^{(2)} |(a''_{40})^{(8)} (T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)} (T_{41}^{(2)}, s_{(40)})| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} \end{aligned}$$

Where  $s_{(40)}$  represents integrand that is integrated over the interval  $[0, t]$  272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} &\leq \\ \frac{1}{(\bar{M}_{40})^{(8)}} &\left( (a_{40})^{(8)} + (a'_{40})^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)} \right) d\left( (G_{43})^{(1)}, (T_{43})^{(1)}; (G_{43})^{(2)}, (T_{43})^{(2)} \right) \end{aligned} \tag{273}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 36:** The fact that we supposed  $(a''_{40})^{(8)}$  and  $(b''_{40})^{(8)}$  depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$  and  $(\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$  respectively of  $\mathbb{R}_+$ . 274

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(8)}$  and  $(b''_i)^{(8)}$ ,  $i = 40, 41, 42$  depend only on  $T_{41}$  and respectively on  $(G_{43})$  (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 37** There does not exist any t where  $G_i(t) = 0$  and  $T_i(t) = 0$  275

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(8)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{40})^{(8)})_1, ((\widehat{M}_{40})^{(8)})_2$  and  $((\widehat{M}_{40})^{(8)})_3$  : 276

**Remark 38:** if  $G_{40}$  is bounded, the same property have also  $G_{41}$  and  $G_{42}$  . indeed if

$$G_{40} < (\widehat{M}_{40})^{(8)} \text{ it follows } \frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a_{41}')^{(8)}G_{41} \text{ and by integrating}$$

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a_{42}')^{(8)}$$

If  $G_{41}$  or  $G_{42}$  is bounded, the same property follows for  $G_{40}$  ,  $G_{42}$  and  $G_{40}$  ,  $G_{41}$  respectively.

**Remark 39:** If  $G_{40}$  is bounded, from below, the same property holds for  $G_{41}$  and  $G_{42}$  . The proof is 277  
 analogous with the preceding one. An analogous property is true if  $G_{41}$  is bounded from below.

**Remark 40:** If  $T_{40}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$  then  $T_{41} \rightarrow \infty$ . 278

**Definition of**  $(m)^{(8)}$  and  $\varepsilon_8$  :

Indeed let  $t_8$  be so that for  $t > t_8$

$$(b_{41}')^{(8)} - (b_i'')^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then  $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$  which leads to 279

$$T_{41} \geq \left( \frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_8 t} = \frac{1}{2} \text{ it results}$$

$$T_{41} \geq \left( \frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8} \text{ By taking now } \varepsilon_8 \text{ sufficiently small one sees that } T_{41} \text{ is unbounded.}$$

The same property holds for  $T_{42}$  if  $\lim_{t \rightarrow \infty} (b_{42}'')^{(8)}((G_{43})(t), t(t), t) = (b_{42}')^{(8)}$

It is now sufficient to take  $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} , \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$  and to choose  $(\widehat{P}_{44})^{(9)}$  and  $(\widehat{Q}_{44})^{(9)}$  large to have 279  
 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[ (\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[ ((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator  $\mathcal{A}^{(9)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying 39,35,36 into itself

The operator  $\mathcal{A}^{(9)}$  is a contraction with respect to the metric

$$d\left( ((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{44})^{(9)}t} \}$$

Indeed if we denote

**Definition of**  $(\widehat{G}_{47}, \widehat{T}_{47}) : (\widehat{G}_{47}, \widehat{T}_{47}) = \mathcal{A}^{(9)}((G_{47}, T_{47}))$

It results

$$\begin{aligned} |\widehat{G}_{44}^{(1)} - \widehat{G}_{44}^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} e^{(\widehat{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{ (a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} e^{(\widehat{M}_{44})^{(9)}s_{(44)}} + \\ G_{44}^{(2)} | (a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a''_{44})^{(9)} (T_{45}^{(2)}, s_{(44)}) | e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} e^{(\widehat{M}_{44})^{(9)}s_{(44)}} \} ds_{(44)} \end{aligned}$$

Where  $s_{(44)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\widehat{M}_{44})^{(9)}t} &\leq \\ \frac{1}{(\widehat{M}_{44})^{(9)}} &((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)} (\widehat{k}_{44})^{(9)}) d\left( ((G_{47})^{(1)}, (T_{47})^{(1)}); (G_{47})^{(2)}, (T_{47})^{(2)} \right) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis (39,35,36) the result follows

**Remark 41:** The fact that we supposed  $(a''_{44})^{(9)}$  and  $(b''_{44})^{(9)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{44})^{(9)} e^{(\widehat{M}_{44})^{(9)}t}$  and  $(\widehat{Q}_{44})^{(9)} e^{(\widehat{M}_{44})^{(9)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(9)}$  and  $(b''_i)^{(9)}, i = 44,45,46$  depend only on  $T_{45}$  and respectively on  $(G_{47})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 42:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(9)} - (a''_i)^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \} ds_{(44)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{44})^{(9)})_1, ((\widehat{M}_{44})^{(9)})_2$  and  $((\widehat{M}_{44})^{(9)})_3$  :

**Remark 43:** if  $G_{44}$  is bounded, the same property have also  $G_{45}$  and  $G_{46}$  . indeed if

$G_{44} < ((\widehat{M}_{44})^{(9)})$  it follows  $\frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)}G_{45}$  and by integrating

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If  $G_{45}$  or  $G_{46}$  is bounded, the same property follows for  $G_{44}$ ,  $G_{46}$  and  $G_{44}$ ,  $G_{45}$  respectively.

**Remark 44:** If  $G_{44}$  is bounded, from below, the same property holds for  $G_{45}$  and  $G_{46}$  . The proof is analogous with the preceding one. An analogous property is true if  $G_{45}$  is bounded from below.

**Remark 45:** If  $T_{44}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b'_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$  then  $T_{45} \rightarrow \infty$ .

**Definition of**  $(m)^{(9)}$  and  $\varepsilon_9$  :

Indeed let  $t_9$  be so that for  $t > t_9$

$$(b_{45})^{(9)} - (b'_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then  $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$  which leads to

$$T_{45} \geq \left( \frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$$

If we take  $t$  such that  $e^{-\varepsilon_9 t} = \frac{1}{2}$  it results

$$T_{45} \geq \left( \frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), t = \log \frac{2}{\varepsilon_9}$$

By taking now  $\varepsilon_9$  sufficiently small one sees that  $T_{45}$  is unbounded.

The same property holds for  $T_{46}$  if  $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

**Behavior of the solutions of equation**

280

**Theorem** If we denote and define

**Definition of**  $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$  :

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$  four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

**Definition of**  $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$  :

281

By  $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$  and respectively  $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$  the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

**Definition of**  $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$  : 282

By  $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$  and respectively  $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$  the roots of the equations  $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$  and  $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

**Definition of**  $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$  :- 283

If we define  $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$  by  
 $(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}$ , if  $(v_0)^{(1)} < (v_1)^{(1)}$

$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}$ , if  $(v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)}$ ,

and  $\boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}$ , if  $(\bar{v}_1)^{(1)} < (v_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}$ , if  $(u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}$ , if  $(u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}$ ,

and  $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}$ , if  $(\bar{u}_1)^{(1)} < (u_0)^{(1)}$  where  $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where  $(p_i)^{(1)}$  is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left( \frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[ e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \right) \leq G_{15}(t) \leq$$
 286

$$\frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[ e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}}$$
 287

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}$$
 288

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[ e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq$$
 289

$$\frac{(a_{15})^{(1)}T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})} \left[ e^{((R_1)^{(1)}+(r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

**Definition of**  $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$ :- 290

$$\text{Where } (S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

**Behavior of the solutions of equation** 291

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$  : 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$  four constants satisfying  
 $-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)}$  293

$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)}$  294

**Definition of**  $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$  : 295

By  $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$  and respectively  $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$  the roots 296

of the equations  $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$  297

and  $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$  and 298

**Definition of**  $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$  : 299

By  $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$  and respectively  $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$  the 300

roots of the equations  $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$  301

and  $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$  302

**Definition of**  $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$  :- 303

If we define  $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$  by 304

$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}$ , **if**  $(v_0)^{(2)} < (v_1)^{(2)}$  305

$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}$ , **if**  $(v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}$ , 306

and  $(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 307$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and  $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$  is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 311$$

$$\left( \frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \left[ e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \right) \leq G_{18}(t) \leq \quad 312$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[ e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[ e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[ e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

**Definition of**  $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$ :- 316

Where  $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$  317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

**Behavior of the solutions** 319

**Theorem 3:** If we denote and define

**Definition of**  $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$  :

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$  four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

**Definition of**  $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$  :

320

By  $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$  and respectively  $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$  the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By  $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$  and respectively  $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$  the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

**Definition of**  $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$  :-

321

If we define  $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$  by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } (u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((s_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(s_1)^{(3)}t}$$

$(p_i)^{(3)}$  is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((s_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(s_1)^{(3)}t}$$

323

$$\left( \frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[ e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \leq \right. \tag{324}$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})} \left[ e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \tag{325}$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \tag{326}$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})} \left[ e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \tag{327}$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[ e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

**Definition of**  $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$ :- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

**Behavior of the solutions of equation**

**Theorem:** If we denote and define

**Definition of**  $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$  :

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$  four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

**Definition of**  $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$  : 329

By  $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$  and respectively  $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$  the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$  : 330

By  $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$  and respectively  $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$  the

roots of the equations  $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and  $(b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

**Definition of**  $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$  :-

If we define  $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$  by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where  $(p_i)^{(4)}$  is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left( \frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[ e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \right) \leq G_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[ e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[ e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[ e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

**Definition of**  $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$ :-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

331

332

333

334

335

336

337

**Behavior of the solutions of equation**

338

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$  :

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$  four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

**Definition of**  $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$  :

339

By  $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$  and respectively  $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$  the roots of the equations

$$(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$$

$$\text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$  :

340

By  $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$  and respectively  $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$  the

roots of the equations  $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and  $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

**Definition of**  $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$  :-

If we define  $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$  by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously

341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities

342

$$G_{28}^0 e^{((s_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(s_1)^{(5)}t}$$

where  $(p_i)^{(5)}$  is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \tag{343}$$

$$\left( \frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \right) \left[ e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \leq \tag{344}$$

$$\frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[ e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \tag{345}$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \tag{346}$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[ e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \tag{347}$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[ e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

**Definition of**  $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$ :- 348

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)} (\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

**Behavior of the solutions of equation** 349

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$  :

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$  four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

**Definition of**  $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$  : 350

By  $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$  and respectively  $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$  the roots of the equations

$$(a_{33})^{(6)} (v^{(6)})^2 + (\sigma_1)^{(6)} v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)} (u^{(6)})^2 + (\tau_1)^{(6)} u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$  : 351

By  $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$  and respectively  $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$  the

roots of the equations  $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and  $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

**Definition of**  $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$  :-

If we define  $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$  by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities

353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where  $(p_i)^{(6)}$  is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left( \frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[ e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \right) \leq G_{34}(t) \leq \quad 355$$

$$\frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[ e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 356}$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[ e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[ e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

**Definition of**  $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$  :- 359

Where  $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

**Behavior of the solutions of equation**

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$  :

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$  four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

**Definition of**  $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$  :

361

By  $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$  and respectively  $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$  the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

and  $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$  and

**Definition of**  $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$  :

362

By  $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$  and respectively  $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$  the

roots of the equations  $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and  $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

**Definition of**  $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$  :-

If we define  $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$  by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \text{ if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \text{ if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

and  $(v_0)^{(7)} = \frac{a_{36}^0}{a_{37}^0}$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \text{ if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \text{ if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \text{ if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and  $(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \text{ if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where  $(p_i)^{(7)}$  is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \tag{365}$$

$$\left( \frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[ e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \leq \tag{366}$$

$$\frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[ e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \tag{367}$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \tag{368}$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[ e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \tag{369}$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[ e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

**Definition of  $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$ :-** 370

Where  $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

**Behavior of the solutions of equation** 371

**Theorem 2:** If we denote and define

**Definition of  $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$  :**

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$  four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) - (b''_{41})^{(8)}(G_{43}, t) \leq -(\tau_1)^{(8)}$$

**Definition of**  $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$  : 372

By  $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$  and respectively  $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$  the roots of the equations

$$(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$  :

By  $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$  and respectively  $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$  the

$$\text{roots of the equations } (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$$

**Definition of**  $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$  :-

If we define  $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$  by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \text{ if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \text{ if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \text{ if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \text{ if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \text{ if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \text{ if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((s_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(s_1)^{(8)}t}$$

where  $(p_i)^{(8)}$  is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \tag{376}$$

$$\left( \frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \left[ e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \right) \leq G_{42}(t) \leq \tag{377}$$

$$\frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})} \left[ e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \tag{378}$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \tag{379}$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})} \left[ e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \tag{380}$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[ e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

**Definition of**  $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$ :- 381

Where  $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

**Behavior of the solutions of equation 37 to 92** 382

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$  :

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$  four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

**Definition of**  $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$  :

By  $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$  and respectively  $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$  the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$  :

By  $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$  and respectively  $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$  the roots of the equations  $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$  and  $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

**Definition of**  $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$  :-

If we define  $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$  by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \text{ if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \text{ if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \text{ if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \text{ if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \text{ if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

$$\text{and } (u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \text{ if } (\bar{u}_1)^{(9)} < (u_0)^{(9)}$  where  $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$  are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((s_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(s_1)^{(9)}t}$$

where  $(p_i)^{(9)}$  is defined by equation 45

$$\frac{1}{(m_2)^{(9)}} G_{44}^0 e^{((s_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(s_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((s_1)^{(9)} - (p_{44})^{(9)} - (s_2)^{(9)})} \left[ e^{((s_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(s_2)^{(9)}t} \right] + G_{46}^0 e^{-(s_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((s_1)^{(9)} - (a'_{46})^{(9)})} \left[ e^{(s_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[ e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[ e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

**Definition of**  $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$ :-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

**Proof:** From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left( (a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

**Definition of**  $v^{(1)}$  :- 
$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$- \left( (a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left( (a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$  :-

$$\text{For } 0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}(v_1)^{(1)} - (v_0)^{(1)}]t}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}(v_1)^{(1)} - (v_0)^{(1)}]t}}, \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

$$\text{it follows } (v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}}, \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}$$

$$\text{From which we deduce } (v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$$

If  $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$  we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}(v_1)^{(1)} - (v_2)^{(1)}]t}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}(v_1)^{(1)} - (v_2)^{(1)}]t}} \leq v^{(1)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}} \leq (\bar{v}_1)^{(1)}$$

If  $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$ , we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{c})^{(1)} (\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)} ((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}) t]}}{1 + (\bar{c})^{(1)} e^{[-(a_{14})^{(1)} ((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}) t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(1)}(t)$  :-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(1)}(t)$  :-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If  $(a'_{13})^{(1)} = (a'_{14})^{(1)}$ , then  $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$  and in this case  $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$  if in addition  $(v_0)^{(1)} = (v_1)^{(1)}$  then  $v^{(1)}(t) = (v_0)^{(1)}$  and as a consequence  $G_{13}(t) = (v_0)^{(1)} G_{14}(t)$  this also defines  $(v_0)^{(1)}$  for the special case

Analogously if  $(b'_{13})^{(1)} = (b'_{14})^{(1)}$ , then  $(\tau_1)^{(1)} = (\tau_2)^{(1)}$  and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$  if in addition  $(u_0)^{(1)} = (u_1)^{(1)}$  then  $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$  This is an important consequence of the relation between  $(v_1)^{(1)}$  and  $(\bar{v}_1)^{(1)}$ , and definition of  $(u_0)^{(1)}$ .

**Proof:** From global equations we obtain

387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left( (a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)} (T_{17}, t) \right) - (a''_{17})^{(2)} (T_{17}, t) v^{(2)} - (a_{17})^{(2)} v^{(2)}$$

**Definition of**  $v^{(2)}$  :-

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

388

It follows

389

$$- \left( (a_{17})^{(2)} (v^{(2)})^2 + (\sigma_2)^{(2)} v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left( (a_{17})^{(2)} (v^{(2)})^2 + (\sigma_1)^{(2)} v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains

390

**Definition of**  $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$  :-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows  $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner , we get

391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce  $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$

392

If  $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$  we find like in the previous case,

393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}$$

If  $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$  , we obtain

394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(2)}(t)$  :-

395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)} , \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain

396

**Definition of**  $u^{(2)}(t)$  :-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)} , \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

397

If  $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}$  , then  $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$  and in this case  $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$  if in addition  $(v_0)^{(2)} = (v_1)^{(2)}$  then  $v^{(2)}(t) = (v_0)^{(2)}$  and as a consequence  $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if  $(b''_{16})^{(2)} = (b''_{17})^{(2)}$ , then  $(\tau_1)^{(2)} = (\tau_2)^{(2)}$  and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$  if in addition  $(u_0)^{(2)} = (u_1)^{(2)}$  then  $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$  This is an important consequence of the relation between  $(v_1)^{(2)}$  and  $(\bar{v}_1)^{(2)}$

**Proof:** From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left( (a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

**Definition of  $v^{(3)}$  :-** 399

$$v^{(3)} = \frac{G_{20}^0}{G_{21}^0}$$

It follows

$$- \left( (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left( (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

it follows  $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

**Definition of  $(\bar{v}_1)^{(3)}$  :-**

From which we deduce  $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If  $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$  we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If  $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$ , we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)})t]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $\nu^{(3)}(t)$  :-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(3)}(t)$  :-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If  $(a''_{20})^{(3)} = (a''_{21})^{(3)}$ , then  $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$  and in this case  $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$  if in addition  $(\nu_0)^{(3)} = (\nu_1)^{(3)}$  then  $\nu^{(3)}(t) = (\nu_0)^{(3)}$  and as a consequence  $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if  $(b''_{20})^{(3)} = (b''_{21})^{(3)}$ , then  $(\tau_1)^{(3)} = (\tau_2)^{(3)}$  and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$  if in addition  $(u_0)^{(3)} = (u_1)^{(3)}$  then  $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$  This is an important consequence of the relation between  $(\nu_1)^{(3)}$  and  $(\bar{\nu}_1)^{(3)}$

**Proof :** From global equations we obtain

404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left( (a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

**Definition of**  $\nu^{(4)}$  :-  $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$- \left( (a_{25})^{(4)}(\nu^{(4)})^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} \right) \leq \frac{d\nu^{(4)}}{dt} \leq - \left( (a_{25})^{(4)}(\nu^{(4)})^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

**Definition of**  $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$  :-

$$\text{For } 0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{[-(a_{25})^{(4)}((\nu_1)^{(4)} - (\nu_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((\nu_1)^{(4)} - (\nu_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}$$

it follows  $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{c})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} (\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}] t}}{4 + (\bar{c})^{(4)} e^{[-(a_{25})^{(4)} (\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}] t}} , \quad \boxed{(\bar{c})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}$$

From which we deduce  $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If  $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$  we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (c)^{(4)} (v_2)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}}{1 + (c)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{c})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{c})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (\bar{v}_1)^{(4)}$$

If  $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$ , we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{c})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}}{1 + (\bar{c})^{(4)} e^{[-(a_{25})^{(4)} ((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}) t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(4)}(t)$  :-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)} , \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(4)}(t)$  :-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)} , \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{24})^{(4)} = (a''_{25})^{(4)}$ , then  $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$  and in this case  $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$  if in addition  $(v_0)^{(4)} = (v_1)^{(4)}$  then  $v^{(4)}(t) = (v_0)^{(4)}$  and as a consequence  $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$  **this also defines  $(v_0)^{(4)}$  for the special case .**

Analogously if  $(b''_{24})^{(4)} = (b''_{25})^{(4)}$ , then  $(\tau_1)^{(4)} = (\tau_2)^{(4)}$  and then

$(u_1)^{(4)} = (\bar{u}_4)^{(4)}$  if in addition  $(u_0)^{(4)} = (u_1)^{(4)}$  then  $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$  This is an important consequence of the relation between  $(v_1)^{(4)}$  and  $(\bar{v}_1)^{(4)}$ , **and definition of  $(u_0)^{(4)}$ .**

408

**Proof:** From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left( (a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)} (T_{29}, t) \right) - (a''_{29})^{(5)} (T_{29}, t) v^{(5)} - (a_{29})^{(5)} v^{(5)}$$

**Definition of**  $v^{(5)}$  :- 
$$v^{(5)} = \frac{G_{28}}{G_{29}}$$

It follows

$$- \left( (a_{29})^{(5)} (v^{(5)})^2 + (\sigma_2)^{(5)} v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left( (a_{29})^{(5)} (v^{(5)})^2 + (\sigma_1)^{(5)} v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$  :-

For  $0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)} (v_2)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (v_0)^{(5)}) t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (v_0)^{(5)}) t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}$$

it follows  $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}$$

From which we deduce  $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If  $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$  we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)} (v_2)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (v_2)^{(5)}) t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (v_2)^{(5)}) t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} \leq (\bar{v}_1)^{(5)}$$

If  $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$  , we obtain

411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(5)}(t)$  :-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)} , \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(5)}(t)$  :-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)} , \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{28})^{(5)} = (a''_{29})^{(5)}$ , then  $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$  and in this case  $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$  if in addition  $(v_0)^{(5)} = (v_5)^{(5)}$  then  $v^{(5)}(t) = (v_0)^{(5)}$  and as a consequence  $G_{28}(t) = (v_0)^{(5)}G_{29}(t)$  **this also defines  $(v_0)^{(5)}$  for the special case .**

Analogously if  $(b''_{28})^{(5)} = (b''_{29})^{(5)}$ , then  $(\tau_1)^{(5)} = (\tau_2)^{(5)}$  and then  $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$  if in addition  $(u_0)^{(5)} = (u_1)^{(5)}$  then  $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$  This is an important consequence of the relation between  $(v_1)^{(5)}$  and  $(\bar{v}_1)^{(5)}$ , **and definition of  $(u_0)^{(5)}$ .**

**Proof :** From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left( (a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

**Definition of  $v^{(6)}$  :-** 
$$v^{(6)} = \frac{G_{32}}{G_{33}}$$

It follows

$$- \left( (a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left( (a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

**Definition of  $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$  :-**

For  $0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows  $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce  $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If  $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$  we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If  $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$ , we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (C)^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(6)}(t)$  :-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(6)}(t)$  :-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{32})^{(6)} = (a''_{33})^{(6)}$ , then  $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$  and in this case  $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$  if in addition  $(v_0)^{(6)} = (v_1)^{(6)}$  then  $v^{(6)}(t) = (v_0)^{(6)}$  and as a consequence  $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$  **this also defines**  $(v_0)^{(6)}$  **for the special case**.

Analogously if  $(b''_{32})^{(6)} = (b''_{33})^{(6)}$ , then  $(\tau_1)^{(6)} = (\tau_2)^{(6)}$  and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$  if in addition  $(u_0)^{(6)} = (u_1)^{(6)}$  then  $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$  This is an important consequence of the relation between  $(v_1)^{(6)}$  and  $(\bar{v}_1)^{(6)}$ , **and definition of**  $(u_0)^{(6)}$ .

416

**Proof :** From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left( (a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

**Definition of**  $v^{(7)}$  :-  $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left( (a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left( (a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$  :-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows  $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{c})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} (\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}] t}}{1 + (\bar{c})^{(7)} e^{[-(a_{37})^{(7)} (\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}] t}} , \quad \boxed{(\bar{c})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce  $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If  $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$  we find like in the previous case,

418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (c)^{(7)} (v_2)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}}{1 + (c)^{(7)} e^{[-(a_{37})^{(7)} ((v_1)^{(7)} - (v_2)^{(7)}) t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{c})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} (\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}] t}}{1 + (\bar{c})^{(7)} e^{[-(a_{37})^{(7)} (\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}] t}} \leq (\bar{v}_1)^{(7)}$$

If  $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$  , we obtain

419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{c})^{(7)} (\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)} (\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}] t}}{1 + (\bar{c})^{(7)} e^{[-(a_{37})^{(7)} (\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}] t}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(7)}(t)$  :-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)} , \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(7)}(t)$  :-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)} , \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{36})^{(7)} = (a''_{37})^{(7)}$ , then  $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$  and in this case  $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$  if in addition  $(v_0)^{(7)} = (v_1)^{(7)}$  then  $v^{(7)}(t) = (v_0)^{(7)}$  and as a consequence  $G_{36}(t) = (v_0)^{(7)} G_{37}(t)$  **this also defines  $(v_0)^{(7)}$  for the special case .**

Analogously if  $(b''_{36})^{(7)} = (b''_{37})^{(7)}$ , then  $(\tau_1)^{(7)} = (\tau_2)^{(7)}$  and then  $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$  if in addition

$(u_0)^{(7)} = (u_1)^{(7)}$  then  $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$  This is an important consequence of the relation between  $(v_1)^{(7)}$  and  $(\bar{v}_1)^{(7)}$ , and definition of  $(u_0)^{(7)}$ .

**Proof:** From global equations we obtain

421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left( (a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

**Definition of**  $v^{(8)}$  :- 
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$- \left( (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq - \left( (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$  :-

For  $0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}(v_1)^{(8)} - (v_0)^{(8)}]t}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}(v_1)^{(8)} - (v_0)^{(8)}]t}}, \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}$$

it follows  $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner, we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}}, \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}$$

From which we deduce  $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If  $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$  we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}(v_1)^{(8)} - (v_2)^{(8)}]t}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}(v_1)^{(8)} - (v_2)^{(8)}]t}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}} \leq (\bar{v}_1)^{(8)}$$

If  $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$ , we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(8)}(t)$  :-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(8)}(t)$  :-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{40})^{(8)} = (a''_{41})^{(8)}$ , then  $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$  and in this case  $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$  if in addition  $(v_0)^{(8)} = (v_1)^{(8)}$  then  $v^{(8)}(t) = (v_0)^{(8)}$  and as a consequence  $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$  **this also defines  $(v_0)^{(8)}$  for the special case .**

Analogously if  $(b''_{40})^{(8)} = (b''_{41})^{(8)}$ , then  $(\tau_1)^{(8)} = (\tau_2)^{(8)}$  and then  $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$  if in addition  $(u_0)^{(8)} = (u_1)^{(8)}$  then  $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$  This is an important consequence of the relation between  $(v_1)^{(8)}$  and  $(\bar{v}_1)^{(8)}$ , **and definition of  $(u_0)^{(8)}$ .**

**Proof :** From 99,20,44,22,23,44 we obtain

424  
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left( (a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

**Definition of**  $v^{(9)}$  :-  $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left( (a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left( (a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$  :-

$$\text{For } 0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}(v_1)^{(9)} - (v_0)^{(9)}]t}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}(v_1)^{(9)} - (v_0)^{(9)}]t}}, \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}$$

it follows  $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{c})^{(9)} (\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)} (\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}] t}}{1 + (\bar{c})^{(9)} e^{[-(a_{45})^{(9)} (\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}] t}}, \quad \boxed{(\bar{c})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}$$

From which we deduce  $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If  $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$  we find like in the previous case,

$$(v_1)^{(9)} \leq \frac{(v_1)^{(9)} + (c)^{(9)} (v_2)^{(9)} e^{[-(a_{45})^{(9)} ((v_1)^{(9)} - (v_2)^{(9)}) t]}}{1 + (c)^{(9)} e^{[-(a_{45})^{(9)} ((v_1)^{(9)} - (v_2)^{(9)}) t]}} \leq v^{(9)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(9)} + (\bar{c})^{(9)} (\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)} (\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}] t}}{1 + (\bar{c})^{(9)} e^{[-(a_{45})^{(9)} (\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}] t}} \leq (\bar{v}_1)^{(9)}$$

If  $0 < (v_1)^{(9)} \leq (\bar{v}_1)^{(9)} \leq \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$ , we obtain

$$(v_1)^{(9)} \leq v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{c})^{(9)} (\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)} (\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}] t}}{1 + (\bar{c})^{(9)} e^{[-(a_{45})^{(9)} (\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)}] t}} \leq (v_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(9)}(t)$  :-

$$(m_2)^{(9)} \leq v^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{v^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(9)}(t)$  :-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{44})^{(9)} = (a''_{45})^{(9)}$ , then  $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$  and in this case  $(v_1)^{(9)} = (\bar{v}_1)^{(9)}$  if in addition  $(v_0)^{(9)} = (v_1)^{(9)}$  then  $v^{(9)}(t) = (v_0)^{(9)}$  and as a consequence  $G_{44}(t) = (v_0)^{(9)} G_{45}(t)$  **this also defines  $(v_0)^{(9)}$  for the special case .**

Analogously if  $(b''_{44})^{(9)} = (b''_{45})^{(9)}$ , then  $(\tau_1)^{(9)} = (\tau_2)^{(9)}$  and then  $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$  if in addition  $(u_0)^{(9)} = (u_1)^{(9)}$  then  $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$  This is an important consequence of the relation between  $(v_1)^{(9)}$  and  $(\bar{v}_1)^{(9)}$ , **and definition of  $(u_0)^{(9)}$ .**

We can prove the following

425

**Theorem :** If  $(a'_i)^{(1)}$  and  $(b'_i)^{(1)}$  are independent on  $t$ , and the conditions with the notations

$$(a'_{13})^{(1)} (a'_{14})^{(1)} - (a_{13})^{(1)} (a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)} (a'_{14})^{(1)} - (a_{13})^{(1)} (a_{14})^{(1)} + (a_{13})^{(1)} (p_{13})^{(1)} + (a'_{14})^{(1)} (p_{14})^{(1)} + (p_{13})^{(1)} (p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0 ,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with  $(p_{13})^{(1)}, (r_{14})^{(1)}$  as defined by equation are satisfied , then the system

**Theorem :** If  $(a_i'')^{(2)}$  and  $(b_i'')^{(2)}$  are independent on  $t$  , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \quad 427$$

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0 , \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with  $(p_{16})^{(2)}, (r_{17})^{(2)}$  as defined by equation are satisfied , then the system

**Theorem :** If  $(a_i'')^{(3)}$  and  $(b_i'')^{(3)}$  are independent on  $t$  , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0 ,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with  $(p_{20})^{(3)}, (r_{21})^{(3)}$  as defined by equation are satisfied , then the system

We can prove the following 432

**Theorem :** If  $(a_i'')^{(4)}$  and  $(b_i'')^{(4)}$  are independent on  $t$  , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0 ,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with  $(p_{24})^{(4)}, (r_{25})^{(4)}$  as defined by equation are satisfied , then the system

**Theorem :** If  $(a_i'')^{(5)}$  and  $(b_i'')^{(5)}$  are independent on  $t$  , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0 ,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{28})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with  $(p_{28})^{(5)}, (r_{29})^{(5)}$  as defined by equation are satisfied , then the system

**Theorem** If  $(a_i'')^{(6)}$  and  $(b_i'')^{(6)}$  are independent on  $t$  , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{32})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with  $(p_{32})^{(6)}, (r_{33})^{(6)}$  as defined by equation are satisfied , then the system

**Theorem :** If  $(a_i'')^{(7)}$  and  $(b_i'')^{(7)}$  are independent on  $t$  , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0 ,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{36})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with  $(p_{36})^{(7)}, (r_{37})^{(7)}$  as defined by equation are satisfied , then the system

**Theorem :** If  $(a_i'')^{(8)}$  and  $(b_i'')^{(8)}$  are independent on  $t$  , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0 ,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{40})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with  $(p_{40})^{(8)}, (r_{41})^{(8)}$  as defined by equation are satisfied , then the system

**Theorem :** If  $(a_i'')^{(9)}$  and  $(b_i'')^{(9)}$  are independent on  $t$  , and the conditions (with the notations 436  
 45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with  $(p_{44})^{(9)}, (r_{45})^{(9)}$  as defined by equation 45 are satisfied, then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution, which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution, which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution, which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}))]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}))]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}))]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

**Proof:** 485

(a) Indeed the first two equations have a nontrivial solution  $G_{13}, G_{14}$  if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

**Proof:** 486

(b) Indeed the first two equations have a nontrivial solution  $G_{16}, G_{17}$  if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

**Proof:** 487

(a) Indeed the first two equations have a nontrivial solution  $G_{20}, G_{21}$  if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

**Proof:** 488

(a) Indeed the first two equations have a nontrivial solution  $G_{24}, G_{25}$  if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

**Proof:** 489

(a) Indeed the first two equations have a nontrivial solution  $G_{28}, G_{29}$  if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

**Proof:** 490

(a) Indeed the first two equations have a nontrivial solution  $G_{32}, G_{33}$  if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

**Proof:** 491

(a) Indeed the first two equations have a nontrivial solution  $G_{36}, G_{37}$  if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

**Proof:** 492

(a) Indeed the first two equations have a nontrivial solution  $G_{40}, G_{41}$  if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

**Proof:** 492

(a) Indeed the first two equations have a nontrivial solution  $G_{44}, G_{45}$  if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

**Definition and uniqueness of  $T_{14}^*$  :-** 493

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(1)}(T_{14})$  being increasing, it follows that there exists a unique  $T_{14}^*$  for which  $f(T_{14}^*) = 0$ . With this value, we obtain from the three first

equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^*)]}$$

**Definition and uniqueness of  $T_{17}^*$  :-** 494

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(2)}(T_{17})$  being increasing, it follows that there exists a unique  $T_{17}^*$  for which  $f(T_{17}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

**Definition and uniqueness of  $T_{21}^*$  :-** 496

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(1)}(T_{21})$  being increasing, it follows that there exists a unique  $T_{21}^*$  for which  $f(T_{21}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

**Definition and uniqueness of  $T_{25}^*$  :-** 497

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(4)}(T_{25})$  being increasing, it follows that there exists a unique  $T_{25}^*$  for which  $f(T_{25}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

**Definition and uniqueness of  $T_{29}^*$  :-** 498

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(5)}(T_{29})$  being increasing, it follows that there exists a unique  $T_{29}^*$  for which  $f(T_{29}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

**Definition and uniqueness of  $T_{33}^*$  :-** 499

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(6)}(T_{33})$  being increasing, it follows that there exists a unique  $T_{33}^*$  for which  $f(T_{33}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

**Definition and uniqueness of  $T_{37}^*$  :-** 500

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(7)}(T_{37})$  being increasing, it follows that

there exists a unique  $T_{37}^*$  for which  $f(T_{37}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

**Definition and uniqueness of  $T_{41}^*$  :-**

501

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(8)}(T_{41})$  being increasing, it follows that there exists a unique  $T_{41}^*$  for which  $f(T_{41}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^*)]}$$

**Definition and uniqueness of  $T_{45}^*$  :-**

501  
A

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(9)}(T_{45})$  being increasing, it follows that there exists a unique  $T_{45}^*$  for which  $f(T_{45}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions  $G_{13}, G_{14}$  if

502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in  $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{14}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{14}^*$  such that  $\varphi(G^*) = 0$

By the same argument, the equations admit solutions  $G_{16}, G_{17}$  if

503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in  $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{17}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{17}^*$  such that  $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions  $G_{20}, G_{21}$  if

505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in  $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{21}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{21}^*$  such that  $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions  $G_{24}, G_{25}$  if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in  $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{25}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{25}^*$  such that  $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions  $G_{28}, G_{29}$  if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in  $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{29}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{29}^*$  such that  $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions  $G_{32}, G_{33}$  if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in  $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{33}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{33}^*$  such that  $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions  $G_{36}, G_{37}$  if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in  $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{37}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{37}^*$  such that  $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions  $G_{40}, G_{41}$  if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$$

$$[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in  $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{41}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{41}^*$  such that  $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions  $G_{44}, G_{45}$  if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in  $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{45}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{45}^*$  such that  $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

$G_{14}^*$  given by  $\varphi(G^*) = 0, T_{14}^*$  given by  $f(T_{14}^*) = 0$  and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

$G_{17}^*$  given by  $\varphi((G_{19})^*) = 0, T_{17}^*$  given by  $f(T_{17}^*) = 0$  and

512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} , G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

513

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} , T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]}$$

514

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

515

$G_{21}^*$  given by  $\varphi((G_{23})^*) = 0, T_{21}^*$  given by  $f(T_{21}^*) = 0$  and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} , G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

516

$G_{25}^*$  given by  $\varphi(G_{27}) = 0, T_{25}^*$  given by  $f(T_{25}^*) = 0$  and

$$G_{24}^* = \frac{(a_{24})^{(4)}G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} , G_{26}^* = \frac{(a_{26})^{(4)}G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)}T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)}T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

$G_{29}^*$  given by  $\varphi((G_{31})^*) = 0$  ,  $T_{29}^*$  given by  $f(T_{29}^*) = 0$  and

$$G_{28}^* = \frac{(a_{28})^{(5)}G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)}G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)}T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)}T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \quad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

$G_{33}^*$  given by  $\varphi((G_{35})^*) = 0$  ,  $T_{33}^*$  given by  $f(T_{33}^*) = 0$  and

$$G_{32}^* = \frac{(a_{32})^{(6)}G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)}G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)}T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)}T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

$G_{37}^*$  given by  $\varphi((G_{39})^*) = 0$  ,  $T_{37}^*$  given by  $f(T_{37}^*) = 0$  and

$$G_{36}^* = \frac{(a_{36})^{(7)}G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)}G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)}T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)}T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]} \quad 523$$

Finally we obtain the unique solution 523

$G_{41}^*$  given by  $\varphi((G_{43})^*) = 0$  ,  $T_{41}^*$  given by  $f(T_{41}^*) = 0$  and

$$G_{40}^* = \frac{(a_{40})^{(8)}G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)}G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)}T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)}T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^*)]} \quad 524$$

Finally we obtain the unique solution of 89 to 99 523

$G_{45}^*$  given by  $\varphi((G_{47})^*) = 0$ ,  $T_{45}^*$  given by  $f(T_{45}^*) = 0$  and

A

$$G_{44}^* = \frac{(a_{44})^{(9)}G_{45}^*}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)}G_{45}^*}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)}T_{45}^*}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)}T_{45}^*}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^*)]}$$

**ASYMPTOTIC STABILITY ANALYSIS**

524

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a'_i)^{(1)}$  and  $(b'_i)^{(1)}$  Belong to  $C^{(1)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

**Proof:** Denote

**Definition of**  $\mathbb{G}_i, \mathbb{T}_i$  :-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a'_{14})^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial(b'_i)^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \tag{525}$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \tag{526}$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \tag{527}$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}(s_{(13)(j)})T_{13}^*\mathbb{G}_j \tag{528}$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}(s_{(14)(j)})T_{14}^*\mathbb{G}_j \tag{529}$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}(s_{(15)(j)})T_{15}^*\mathbb{G}_j \tag{530}$$

**ASYMPTOTIC STABILITY ANALYSIS**

531

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a'_i)^{(2)}$  and  $(b'_i)^{(2)}$  Belong to  $C^{(2)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

**Proof:** Denote

**Definition of**  $\mathbb{G}_i, \mathbb{T}_i$  :-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a'_{17})^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)}, \quad \frac{\partial(b''_i)^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \tag{534}$$

$$\frac{dG_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \tag{535}$$

$$\frac{dG_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \tag{536}$$

$$\frac{dT_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(16)(j)})T_{16}^*G_j \tag{537}$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)})T_{17}^*G_j \tag{538}$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)})T_{18}^*G_j \tag{539}$$

**ASYMPTOTIC STABILITY ANALYSIS** 540

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a''_i)^{(3)}$  and  $(b''_i)^{(3)}$  belong to  $C^{(3)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial(b''_i)^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \tag{541}$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \tag{542}$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \tag{543}$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)})T_{20}^*G_j \tag{544}$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)})T_{21}^*G_j \tag{545}$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)})T_{22}^*G_j \tag{546}$$

**ASYMPTOTIC STABILITY ANALYSIS** 547

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions

$(a_i'')^{(4)}$  and  $(b_i'')^{(4)}$  Belong to  $C^{(4)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :- 548

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})G_{24} + (a_{24})^{(4)}G_{25} - (q_{24})^{(4)}G_{24}^*T_{25} \quad 549$$

$$\frac{dG_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})G_{25} + (a_{25})^{(4)}G_{24} - (q_{25})^{(4)}G_{25}^*T_{25} \quad 550$$

$$\frac{dG_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})G_{26} + (a_{26})^{(4)}G_{25} - (q_{26})^{(4)}G_{26}^*T_{25} \quad 551$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)})T_{24}^*G_j \quad 552$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)})T_{25}^*G_j \quad 553$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)})T_{26}^*G_j \quad 554$$

**ASYMPTOTIC STABILITY ANALYSIS** 555

**Theorem 5:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(5)}$  and  $(b_i'')^{(5)}$  Belong to  $C^{(5)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :- 556

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)}, \quad \frac{\partial(b_i'')^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})G_{28} + (a_{28})^{(5)}G_{29} - (q_{28})^{(5)}G_{28}^*T_{29} \quad 557$$

$$\frac{dG_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})G_{29} + (a_{29})^{(5)}G_{28} - (q_{29})^{(5)}G_{29}^*T_{29} \quad 558$$

$$\frac{dG_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})G_{30} + (a_{30})^{(5)}G_{29} - (q_{30})^{(5)}G_{30}^*T_{29} \quad 559$$

$$\frac{dT_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})T_{28} + (b_{28})^{(5)}T_{29} + \sum_{j=28}^{30} (s_{(28)(j)})T_{28}^*G_j \quad 560$$

$$\frac{dT_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})T_{29} + (b_{29})^{(5)}T_{28} + \sum_{j=28}^{30} (s_{(29)(j)})T_{29}^*G_j \quad 561$$

$$\frac{dT_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})T_{30} + (b_{30})^{(5)}T_{29} + \sum_{j=28}^{30} (s_{(30)(j)})T_{30}^*G_j \quad 562$$

**ASYMPTOTIC STABILITY ANALYSIS** 563

**Theorem 6:** If the conditions of the previous theorem are satisfied and if the functions  $(a'_i)^{(6)}$  and  $(b'_i)^{(6)}$  belong to  $C^{(6)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of  $G_i, T_i$  :-** 564

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial(b'_i)^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^*T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^*T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^*T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34} (s_{(32)(j)})T_{32}^*G_j \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34} (s_{(33)(j)})T_{33}^*G_j \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34} (s_{(34)(j)})T_{34}^*G_j \quad 570$$

**ASYMPTOTIC STABILITY ANALYSIS** 571

**Theorem 7:** If the conditions of the previous theorem are satisfied and if the functions  $(a'_i)^{(7)}$  and  $(b'_i)^{(7)}$  belong to  $C^{(7)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of  $G_i, T_i$  :-** 572

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a''_{37})^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial(b'_i)^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{dG_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})G_{36} + (a_{36})^{(7)}G_{37} - (q_{36})^{(7)}G_{36}^*T_{37} \quad 573$$

$$\frac{dG_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})G_{37} + (a_{37})^{(7)}G_{36} - (q_{37})^{(7)}G_{37}^*T_{37} \quad 574$$

$$\frac{dG_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})G_{38} + (a_{38})^{(7)}G_{37} - (q_{38})^{(7)}G_{38}^*T_{37} \quad 575$$

$$\frac{dT_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})T_{36} + (b_{36})^{(7)}T_{37} + \sum_{j=36}^{38} (s_{(36)(j)})T_{36}^*G_j \quad 576$$

$$\frac{dT_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})T_{37} + (b_{37})^{(7)}T_{36} + \sum_{j=36}^{38} (s_{(37)(j)})T_{37}^*G_j \quad 578$$

$$\frac{dT_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})T_{38} + (b_{38})^{(7)}T_{37} + \sum_{j=36}^{38} (s_{(38)(j)})T_{38}^*G_j \quad 579$$

Obviously, these values represent an equilibrium solution

### ASYMPTOTIC STABILITY ANALYSIS

**Theorem 8:** If the conditions of the previous theorem are satisfied and if the functions  $(a'_i)^{(8)}$  and  $(b'_i)^{(8)}$  belong to  $C^{(8)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a'_{41})^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial (b'_{41})^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^*T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^*T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^*T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)})T_{40}^*G_j \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)})T_{41}^*G_j \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)})T_{42}^*G_j \quad 586$$

### ASYMPTOTIC STABILITY ANALYSIS

**Theorem 9:** If the conditions of the previous theorem are satisfied and if the functions  $(a'_i)^{(9)}$  and  $(b'_i)^{(9)}$  belong to  $C^{(9)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

**Proof:** Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial(a'_{45})^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial(b'_{47})^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586$$

B

$$\frac{dG_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})G_{45} + (a_{45})^{(9)}G_{44} - (q_{45})^{(9)}G_{45}^*T_{45} \quad 586$$

C

$$\frac{dG_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})G_{46} + (a_{46})^{(9)}G_{45} - (q_{46})^{(9)}G_{46}^*T_{45} \quad 586$$

D

$$\frac{dT_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})T_{44} + (b_{44})^{(9)}T_{45} + \sum_{j=44}^{46} (s_{(44)(j)})T_{44}^*G_j \quad 586$$

E

$$\frac{dT_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})T_{45} + (b_{45})^{(9)}T_{44} + \sum_{j=44}^{46} (s_{(45)(j)})T_{45}^*G_j \quad 586$$

F

$$\frac{dT_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})T_{46} + (b_{46})^{(9)}T_{45} + \sum_{j=44}^{46} (s_{(46)(j)})T_{46}^*G_j \quad 586$$

G

**The characteristic equation of this system is**

587

$$((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right]$$

$$\left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \right)$$

$$+ \left( ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \right)$$

$$\left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \right)$$

$$\left( ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right)$$

$$\left( ((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right)$$

$$+ \left( ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)}G_{15}$$

$$+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) \left( (a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^* \right)$$

$$\left. \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \right) \right\} = 0$$

+

$$\begin{aligned}
 & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\
 & [((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^*]\} \\
 & ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \\
 & + ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \\
 & ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \\
 & ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} \\
 & ((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}) (\lambda)^{(2)} \\
 & + ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}) (\lambda)^{(2)} (q_{18})^{(2)}G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}) ((a_{18})^{(2)}(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}G_{16}^*) \\
 & ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(18)}T_{17}^* + (b_{17})^{(2)}s_{(16),(18)}T_{16}^* \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)})\{((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & [((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)})(q_{21})^{(3)}G_{21}^* + (a_{21})^{(3)}(q_{20})^{(3)}G_{20}^*]\} \\
 & ((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)})s_{(21),(21)}T_{21}^* + (b_{21})^{(3)}s_{(20),(21)}T_{21}^* \\
 & + ((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)})(q_{20})^{(3)}G_{20}^* + (a_{20})^{(3)}(q_{21})^{(1)}G_{21}^* \\
 & ((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)})s_{(21),(20)}T_{21}^* + (b_{21})^{(3)}s_{(20),(20)}T_{20}^* \\
 & ((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} \\
 & ((\lambda)^{(3)})^2 + ((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)}) (\lambda)^{(3)} \\
 & + ((\lambda)^{(3)})^2 + ((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)}) (\lambda)^{(3)} (q_{22})^{(3)}G_{22} \\
 & + ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)}) ((a_{22})^{(3)}(q_{21})^{(3)}G_{21}^* + (a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G_{20}^*) \\
 & ((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)})s_{(21),(22)}T_{21}^* + (b_{21})^{(3)}s_{(20),(22)}T_{20}^* \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)})\{((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & \left[ ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)})(q_{25})^{(4)}G_{25}^* + (a_{25})^{(4)}(q_{24})^{(4)}G_{24}^* \right] \\
 & \left( ((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)})S_{(25),(25)}T_{25}^* + (b_{25})^{(4)}S_{(24),(25)}T_{25}^* \right) \\
 & + \left( ((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)})(q_{24})^{(4)}G_{24}^* + (a_{24})^{(4)}(q_{25})^{(4)}G_{25}^* \right) \\
 & \left( ((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)})S_{(25),(24)}T_{25}^* + (b_{25})^{(4)}S_{(24),(24)}T_{24}^* \right) \\
 & \left( ((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & \left( ((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & + \left( ((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) (q_{26})^{(4)}G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)}(q_{25})^{(4)}G_{25}^* + (a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G_{24}^*) \\
 & \left. \left( ((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)})S_{(25),(26)}T_{25}^* + (b_{25})^{(4)}S_{(24),(26)}T_{24}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)})\{((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[ ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)})(q_{29})^{(5)}G_{29}^* + (a_{29})^{(5)}(q_{28})^{(5)}G_{28}^* \right] \\
 & \left( ((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})S_{(29),(29)}T_{29}^* + (b_{29})^{(5)}S_{(28),(29)}T_{29}^* \right) \\
 & + \left( ((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)})(q_{28})^{(5)}G_{28}^* + (a_{28})^{(5)}(q_{29})^{(5)}G_{29}^* \right) \\
 & \left( ((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})S_{(29),(28)}T_{29}^* + (b_{29})^{(5)}S_{(28),(28)}T_{28}^* \right) \\
 & \left( ((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \left( ((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & + \left( ((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)}G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)}(q_{29})^{(5)}G_{29}^* + (a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G_{28}^*) \\
 & \left. \left( ((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})S_{(29),(30)}T_{29}^* + (b_{29})^{(5)}S_{(28),(30)}T_{28}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)})\{((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[ ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)})(q_{33})^{(6)}G_{33}^* + (a_{33})^{(6)}(q_{32})^{(6)}G_{32}^* \right] \\
 & \left( ((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})S_{(33),(33)}T_{33}^* + (b_{33})^{(6)}S_{(32),(33)}T_{33}^* \right) \\
 & + \left( ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)})(q_{32})^{(6)}G_{32}^* + (a_{32})^{(6)}(q_{33})^{(6)}G_{33}^* \right) \\
 & \left( ((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})S_{(33),(32)}T_{33}^* + (b_{33})^{(6)}S_{(32),(32)}T_{32}^* \right) \\
 & \left( ((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & \left( ((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 & + \left( ((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) (q_{34})^{(6)}G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)}(q_{33})^{(6)}G_{33}^* + (a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G_{32}^*) \\
 & \left. \left( ((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})S_{(33),(34)}T_{33}^* + (b_{33})^{(6)}S_{(32),(34)}T_{32}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)})\{((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & \left[ ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)})(q_{37})^{(7)}G_{37}^* + (a_{37})^{(7)}(q_{36})^{(7)}G_{36}^* \right] \\
 & \left( ((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})S_{(37),(37)}T_{37}^* + (b_{37})^{(7)}S_{(36),(37)}T_{37}^* \right) \\
 & + \left( ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)})(q_{36})^{(7)}G_{36}^* + (a_{36})^{(7)}(q_{37})^{(7)}G_{37}^* \right) \\
 & \left( ((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})S_{(37),(36)}T_{37}^* + (b_{37})^{(7)}S_{(36),(36)}T_{36}^* \right) \\
 & \left( ((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & \left( ((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)}) (\lambda)^{(7)} \right) \\
 & + \left( ((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)}) (\lambda)^{(7)} \right) (q_{38})^{(7)}G_{38} \\
 & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}) ((a_{38})^{(7)}(q_{37})^{(7)}G_{37}^* + (a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G_{36}^*) \\
 & \left. \left( ((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})S_{(37),(38)}T_{37}^* + (b_{37})^{(7)}S_{(36),(38)}T_{36}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)})\{((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\
 & \left[ ((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)})(q_{41})^{(8)}G_{41}^* + (a_{41})^{(8)}(q_{40})^{(8)}G_{40}^* \right] \\
 & \left( ((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)})S_{(41),(41)}T_{41}^* + (b_{41})^{(8)}S_{(40),(41)}T_{41}^* \right) \\
 & + \left( ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)})(q_{40})^{(8)}G_{40}^* + (a_{40})^{(8)}(q_{41})^{(8)}G_{41}^* \right) \\
 & \left( ((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)})S_{(41),(40)}T_{41}^* + (b_{41})^{(8)}S_{(40),(40)}T_{40}^* \right) \\
 & \left( ((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 & \left( ((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)}) (\lambda)^{(8)} \right) \\
 & + \left( ((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)}) (\lambda)^{(8)} \right) (q_{42})^{(8)}G_{42} \\
 & + ((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)}) ((a_{42})^{(8)}(q_{41})^{(8)}G_{41}^* + (a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G_{40}^*) \\
 & \left. \left( ((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)})S_{(41),(42)}T_{41}^* + (b_{41})^{(8)}S_{(40),(42)}T_{40}^* \right) \right\} = 0 \\
 & + \\
 & ((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)})\{((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 & \left[ ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)})(q_{45})^{(9)}G_{45}^* + (a_{45})^{(9)}(q_{44})^{(9)}G_{44}^* \right] \\
 & \left( ((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)})S_{(45),(45)}T_{45}^* + (b_{45})^{(9)}S_{(44),(45)}T_{45}^* \right) \\
 & + \left( ((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)})(q_{44})^{(9)}G_{44}^* + (a_{44})^{(9)}(q_{45})^{(9)}G_{45}^* \right) \\
 & \left( ((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)})S_{(45),(44)}T_{45}^* + (b_{45})^{(9)}S_{(44),(44)}T_{44}^* \right) \\
 & \left( ((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 & \left( ((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}) (\lambda)^{(9)} \right) \\
 & + \left( ((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}) (\lambda)^{(9)} \right) (q_{46})^{(9)}G_{46} \\
 & + ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}) ((a_{46})^{(9)}(q_{45})^{(9)}G_{45}^* + (a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G_{44}^*) \\
 & \left. \left( ((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)})S_{(45),(46)}T_{45}^* + (b_{45})^{(9)}S_{(44),(46)}T_{44}^* \right) \right\} = 0
 \end{aligned}$$

**And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.**

## References

- (1) ^ Wolchover, Natalie (November 29, 2012). "Supersymmetry Fails Test, Forcing Physics to Seek New Ideas". Scientific American.
- (2) ^ <http://www.bbc.co.uk/news/science-environment-23431797>
- (3) ^ a b c Gordon L. Kane, The Dawn of Physics Beyond the Standard Model, Scientific American, June 2003, page 60 and The frontiers of physics, special edition, Vol 15, #3, page 8 "Indirect evidence for supersymmetry comes from the extrapolation of interactions to high energies."
- (4) ^ Jonathan Feng: Supersymmetric Dark Matter (pdf), University of California, Irvine, 11 May 2007
- (5) ^ Torsten Bringmann: The WIMP "Miracle" (pdf) University of Hamburg
- (6) ^ <http://profmattstrassler.com/articles-and-posts/lhcposts/what-do-current-mid-august-2011-lhc-results-imply-about-supersymmetry/>
- (7) ^ a b ATLAS SUSY search documents
- (8) ^ a b CMS SUSY search documents
- (9) ^ R. Haag, J. T. Lopuszanski and M. Sohnius, "All Possible Generators Of Supersymmetries Of The S Matrix", Nucl. Phys. B 88 (1975) 257
- (10) ^ H. Miyazawa (1966). "Baryon Number Changing Currents". Prog. Theor. Phys. 36 (6): 1266–1276. Bibcode:1966PTPh..36.1266M. doi:10.1143/PTP.36.1266.
- (11) ^ H. Miyazawa (1968). "Spinor Currents and Symmetries of Baryons and Mesons". Phys. Rev. 170 (5): 1586–1590. Bibcode:1968PhRv..170.1586M. doi:10.1103/PhysRev.170.1586.
- (12) ^ Michio Kaku, Quantum Field Theory, ISBN 0-19-509158-2, pg 663.
- (13) ^ Peter Freund, Introduction to Supersymmetry, ISBN 0-521-35675-X, pages 26-27, 138.
- (14) ^ Gervais, J. -L.; Sakita, B. (1971). "Field theory interpretation of supergauges in dual models". Nuclear Physics B 34 (2): 632. doi:10.1016/0550-3213(71)90351-8.
- (15) ^ D.V. Volkov, V.P. Akulov, Pisma Zh.Eksp.Teor.Fiz. 16 (1972) 621; Phys.Lett. B46 (1973) 109; V.P. Akulov, D.V. Volkov, Teor.Mat.Fiz. 18 (1974) 39
- (16) ^ Ramond, P. (1971). "Dual Theory for Free Fermions". Physical Review D 3 (10): 2415. doi:10.1103/PhysRevD.3.2415.
- (17) ^ Wess, J.; Zumino, B. (1974). "Supergauge transformations in four dimensions". Nuclear Physics B 70: 39. doi:10.1016/0550-3213(74)90355-1.
- (18) ^ Iachello, F. (1980). "Dynamical Supersymmetries in Nuclei". Physical Review Letters