

# Yang–Mills Existence and Mass Gap (Unsolved Problem): Aufklärung La Alltagsgeschichte: Enlightenment of a Micro History

Dr. K.N.P. Kumar

Post doctoral fellow, Department of mathematics, Kuvempu University, Shimoga, Karnataka, India

**Abstract:** Yang–Mills theory is the (non-Abelian) quantum field theory underlying the Standard Model of particle physics;  $\mathbb{R}^4$  is Euclidean 4-space; the mass gap  $\Delta$  is the mass of the least massive particle predicted by the theory. Therefore, the winner must first prove that Yang–Mills theory exists and that it satisfies the standard of rigor that characterizes contemporary mathematical physics, in particular constructive quantum field theory, which is referenced in the papers 45 and 35 cited in the official problem description by Jaffe and Witten. The winner must then prove that the mass of the least massive particle of the force field predicted by the theory is strictly positive. For example, in the case of  $G=SU(3)$  - the strong nuclear interaction - the winner must prove that glueballs have a lower mass bound, and thus cannot be arbitrarily light. Biagio Lucini, Michael Teper, Urs Wenger studied Glueballs and k-strings in  $SU(N)$  gauge theories : calculations with improved operators testing a variety of blocking and smearing algorithms **for constructing** glueball and string wave-functionals, **and find** some with much improved overlaps onto the lightest states. They use these algorithms to **obtain** improved results on the tensions of k-strings in  $SU(4)$ ,  $SU(6)$ , and  $SU(8)$  gauge theories. Authors emphasise the major systematic errors that still need **to be controlled** in calculations of heavier k-strings, and perform calculations in  $SU(4)$  on an anisotropic lattice in a bid to minimise one of these. **All these results point to** the k-string tensions lying part-way between the 'MQCD' and 'Casimir Scaling' conjectures, with the power in  $1/N$  of the leading correction lying in  $[1,2]$ . (See the paper). **They also obtain some evidence** for the presence of quasi-stable strings in calculations that do not use sources, and observe some near-degeneracies between (excited) strings in different representations. We also calculate the lightest glueball masses for  $N=2\dots 8$ , and extrapolate to  $N=\infty$ , **obtaining results** compatible with earlier work. Biagio Lucini et al show that the  $N=\infty$  factorization of the Euclidean correlators that are used in such mass calculations does not make the masses any less calculable at large  $N$ . JHEP0406:012,2004DOI: 10.1088/1126-6708/2004/06/012 arXiv: hep-lat/0404008. Quantum field theory (QFT) is a theoretical framework for constructing quantum mechanical models of subatomic particles in particle physics and quasiparticles in condensed matter physics, by treating a particle as an excited state of an underlying physical field. These excited states are called field quanta. For example, quantum electrodynamics (QED) has one electron field and one photon field, quantum chromodynamics (QCD) has one field for each type of quark, and in condensed matter there is an atomic displacement field that gives rise to phonon particles. Ed Witten describes QFT as "by far" the most difficult theory in modern physics. Towards the end of consummation of solution of this long outstanding problem we make two assumptions that the statements are true or not and the properties is testified by manifested actions. This bears ample testimony, infallible observatory and impeccable demonstration of the fact that state mental propositions in either case shall testify the prediction, projection, stability analysis results by experiments to prove or disprove the theory. In essence the method is that of false princeps and reductio ad absurdum. Quintessentially it is one model. Towards the end of circumvention of repeated projection of superscripts and subscripts which is of the order 56, we give the model in two sections. Notwithstanding variables are all to be taken as different and concatenation is to be done. As said towards the end of

obtention of felicity of expression and avoiding the extensive superscriptal and subscriptal typing which might cause systemic errors, model is bifurcated in to two. Section two is only progressive of section one.

### INTRODUCTION—VARIABLES USED

Source: Wikipedia

The problem is phrased as follows:

Yang–Mills Existence and Mass Gap

- (1) For any compact simple gauge group  $G$ , a non-trivial quantum Yang–Mills theory **exists (eb) on  $\mathbb{R}^4$**  and has a mass gap  $\Delta > 0$ . Existence includes establishing axiomatic properties at least as strong as those cited in Streater & Wightman (1964), Osterwalder & Schrader (1973) and Osterwalder & Schrader (1975).
- (2) For any compact simple gauge group  $G$ , a non-trivial quantum Yang–Mills theory does not exist **(eb) on  $\mathbb{R}^4$**  and has a mass gap  $\Delta > 0$ . Existence includes establishing axiomatic properties at least as strong as those cited in Streater & Wightman (1964), Osterwalder & Schrader (1973) and Osterwalder & Schrader (1975).
- (3) In this statement, Yang–Mills theory is (=) the (non-Abelian) quantum field theory underlying the Standard Model of particle physics;  $\mathbb{R}^4$  is Euclidean 4-space
- (4) The mass gap  $\Delta$  is the mass of the least massive particle predicted by the theory.
- (5) Therefore, the winner must first prove that Yang–Mills theory exists and that **it (eb) satisfies** the standard of rigor that characterizes contemporary mathematical physics, in particular constructive quantum field theory, which is referenced in the papers 45 and 35 cited in the official problem description by Jaffe and Witten.
- (6) The winner must then prove that the mass of the least massive particle of the force field predicted by the theory **is (=) strictly positive**.
- (7) For example, in the case of  $G=\text{SU}(3)$  - the strong nuclear interaction - the winner must prove that glueballs **have (e) a lower mass bound**
- (8) Thus glueballs cannot (e) be arbitrarily light.
- (9) Yang–Mills theories are a special example of gauge theory with a non-abelian symmetry group given by the Lagrangian

$$\mathcal{L}_{\text{gf}} = -\frac{1}{2} \text{Tr}(F^2) = -\frac{1}{4} F^{a\mu\nu} F_{\mu\nu}^a$$

with the generators of the Lie algebra corresponding to the F-quantities (the curvature or field-strength form) satisfying

$$\text{Tr}(T^a T^b) = \frac{1}{2} \delta^{ab}, \quad [T^a, T^b] = i f^{abc} T^c$$

and the covariant derivative defined as

$$D_\mu = I \partial_\mu - ig T^a A_\mu^a$$

where  $I$  is the identity for the group generators,  $A_\mu^a$  is the vector potential, and  $g$  is the coupling constant. In four dimensions, the coupling constant  $g$  is a pure number and for a  $\text{SU}(N)$  group one

has  $a, b, c = 1 \dots N^2 - 1$ .

The relation

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + gf^{abc} A_\mu^b A_\nu^c$$

can be derived by the commutator

$$[D_\mu, D_\nu] = -igT^a F_{\mu\nu}^a.$$

The field has the property of being self-interacting and equations of motion that one obtains are said to be semilinear, as nonlinearities are both with and without derivatives. This means that one can manage this theory only by perturbation theory, with small nonlinearities.

Note that the transition between "upper" ("contravariant") and "lower" ("covariant") vector or tensor components is trivial for a indices (e.g.  $f^{abc} = f_{abc}$ ), whereas for  $\mu$  and  $\nu$  it is nontrivial, corresponding e.g. to the usual Lorentz signature,  $\eta_{\mu\nu} = \text{diag}(+ - - -)$ .

From the given Lagrangian one can derive the equations of motion given by

$$\partial^\mu F_{\mu\nu}^a + gf^{abc} A^{\mu b} F_{\mu\nu}^c = 0.$$

Putting  $F_{\mu\nu} = T^a F_{\mu\nu}^a$ , these can be rewritten as

$$(D^\mu F_{\mu\nu})^a = 0.$$

A Bianchi identity holds

$$(D_\mu F_{\nu\kappa})^a + (D_\kappa F_{\mu\nu})^a + (D_\nu F_{\kappa\mu})^a = 0.$$

which is equivalent to the Jacobi identity

$$[D_\mu, [D_\nu, D_\kappa]] + [D_\kappa, [D_\mu, D_\nu]] + [D_\nu, [D_\kappa, D_\mu]] = 0.$$

since  $[D_\mu, F_{\nu\kappa}^a] = -igD_\mu F_{\nu\kappa}^a$ . Define the dual strength tensor  $\tilde{F}^{\mu\nu} = \frac{1}{2}\epsilon^{\mu\nu\rho\sigma} F_{\rho\sigma}$ , then the Bianchi identity can be rewritten as

$$D_\mu \tilde{F}^{\nu\kappa} = 0.$$

A source  $J_\mu^a$  enters into **the equations of motion** as

$$\partial^\mu F_{\mu\nu}^a + gf^{abc} A^{\mu b} F_{\mu\nu}^c = -J_\nu^a.$$

Note that the currents must properly change under gauge group transformations.

We give here some comments about the physical dimensions of the coupling. We note that, in D dimensions, the field scales as  $[A] = [L^{\frac{2-D}{2}}]$  and so the coupling must scale as  $[g^2] = [L^{D-4}]$ . This implies that Yang–Mills theory is not renormalizable for dimensions greater than four. Further, we note that, for D = 4, the coupling is dimensionless and both the field and the square of the coupling have the same dimensions of the field and the coupling of a massless quartic scalar field theory. So, these theories share the scale invariance at the classical level.

## NOTATION

### Module One

For any compact simple gauge group G, a non-trivial quantum Yang–Mills theory **exists (eb) on  $\mathbb{R}^4$**  and has a mass gap  $\Delta > 0$ . Existence includes establishing axiomatic properties at least as strong as those cited in Streater & Wightman (1964), Osterwalder & Schrader (1973) and Osterwalder & Schrader (1975).

$G_{13}$  : Category one of For any compact simple gauge group G, a non-trivial quantum Yang–Mills theory

$G_{14}$  : Category two of For any compact simple gauge group G, a non-trivial quantum Yang–Mills theory. Systemic differentiation. There are various systems to which Yang Mills theory is applicable and mass gap exists. Characteristics of these systems are taken I to consideration in the consummation of the diaspora fabric of the classification doxa.

$G_{15}$  : Category three of For any compact simple gauge group G, a non-trivial quantum Yang–Mills theory

$T_{13}$  : Category one of **exists (eb) on  $\mathbb{R}^4$**  and has a mass gap  $\Delta > 0$ . Existence includes establishing axiomatic properties at least as strong as those cited in Streater & Wightman (1964), Osterwalder & Schrader (1973) and Osterwalder & Schrader (1975).

$T_{14}$  : Category two of **exists (eb) on  $\mathbb{R}^4$**  and has a mass gap  $\Delta > 0$ . Existence includes establishing axiomatic properties at least as strong as those cited in Streater & Wightman (1964), Osterwalder & Schrader (1973) and Osterwalder & Schrader (1975).

$T_{15}$  : Category three of **exists (eb) on  $\mathbb{R}^4$**  and has a mass gap  $\Delta > 0$ . Existence includes establishing axiomatic properties at least as strong as those cited in Streater & Wightman (1964), Osterwalder & Schrader (1973) and Osterwalder & Schrader (1975).

### Module Two

For any compact simple gauge group G, a non-trivial quantum Yang–Mills theory does not exist **(eb) on  $\mathbb{R}^4$**  and has a mass gap  $\Delta > 0$ . Existence includes establishing axiomatic properties at least as strong as those cited in Streater & Wightman (1964), Osterwalder & Schrader (1973) and Osterwalder & Schrader (1975)

$G_{16}$  : Category one of For any compact simple gauge group G, a non-trivial quantum Yang–Mills theory does not

$G_{17}$ : Category two of For any compact simple gauge group G, a non-trivial quantum Yang–Mills theory does not

$G_{18}$ : Category three of For any compact simple gauge group G, a non-trivial quantum Yang–Mills theory does not

$T_{16}$ : Category one of existence **(eb) on  $\mathbb{R}^4$**  and has a mass gap  $\Delta > 0$ . Existence includes establishing axiomatic properties at least as strong as those cited in Streater & Wightman (1964), Osterwalder & Schrader

(1973) and Osterwalder & Schrader (1975)

$T_{17}$  : Category two of existence (eb) on  $\mathbb{R}^4$  and has a mass gap  $\Delta > 0$ . Existence includes establishing axiomatic properties at least as strong as those cited in Streater & Wightman (1964), Osterwalder & Schrader (1973) and Osterwalder & Schrader (1975)

$T_{18}$  : Category three of existence (eb) on  $\mathbb{R}^4$  and has a mass gap  $\Delta > 0$ . Existence includes establishing axiomatic properties at least as strong as those cited in Streater & Wightman (1964), Osterwalder & Schrader (1973) and Osterwalder & Schrader (1975)

### Module three

In this statement, Yang–Mills theory is (=) the (non-Abelian) quantum field theory underlying the Standard Model of particle physics;  $\mathbb{R}^4$  is Euclidean 4-space

$G_{20}$  : Category one of (non-Abelian) quantum field theory underlying the Standard Model of particle physics;  $\mathbb{R}^4$  is Euclidean 4-space

$G_{21}$  : Category two of (non-Abelian) quantum field theory underlying the Standard Model of particle physics;  $\mathbb{R}^4$  is Euclidean 4-space

$G_{22}$  : Category three of (non-Abelian) quantum field theory underlying the Standard Model of particle physics;  $\mathbb{R}^4$  is Euclidean 4-space

$T_{20}$  : Category one of Yang–Mills theory. Systemic differentiation is undertaken for execution. There are various systems in the world that satisfy the axiomatic predications, postulation alcovishness, and phenomenological correlates of the Yang mills Theory. Some of them are under experimental observation. Characteristics of these systems so mentioned in the foregoing and which are under the investigation form the bastion for the classification scheme.

$T_{21}$ : Category two of Yang–Mills theory

$T_{22}$  : Category three of Yang–Mills theory

### Module four

The mass gap  $\Delta$  is the mass of the least massive particle predicted by the theory

$G_{24}$  : Category one of mass of the least massive particle predicted by the theory

$G_{25}$  : Category two of mass of the least massive particle predicted by the theory

$G_{26}$  : Category three of mass of the least massive particle predicted by the theory

$T_{24}$  : Category one of mass gap  $\Delta$ . Please note that the characteristics of the investigatory systems that are under consideration and has mass gap syndrome form the stylobate and sentinel , the fulcrum of the classification scheme.

$T_{25}$  : Category two of mass gap  $\Delta$

$T_{26}$  : Category three of mass gap  $\Delta$

### Module five

Therefore, the winner must first prove that Yang–Mills theory exists and that **it (eb) satisfies** the standard of rigor that characterizes contemporary mathematical physics, in particular constructive quantum field theory, which is referenced in the papers 45 and 35 cited in the official problem description by Jaffe and Witten. We assume the proposition and give the model. **Model gives prediction, projection and prognostication of the variables involved, and in the eventuality of the correctness of the statement it shall remain with the initial conditions stated in unmistakable terms in the final results in the dovetailed mathematical exposition.**

$G_{28}$  : Category one of Yang–Mills theory exists and that **it**

$G_{29}$  : Category two of Yang–Mills theory exists and that **it**

$G_{30}$  : Category three of Yang–Mills theory exists and that **it**

$T_{28}$  : Category one of standard of rigor that characterizes contemporary mathematical physics, in particular constructive quantum field theory, which is referenced in the papers 45 and 35 cited in the official problem description by Jaffe and Witten

$T_{29}$  : Category two of standard of rigor that characterizes contemporary mathematical physics, in particular constructive quantum field theory, which is referenced in the papers 45 and 35 cited in the official problem description by Jaffe and Witten

$T_{30}$  : Category three of standard of rigor that characterizes contemporary mathematical physics, in particular constructive quantum field theory, which is referenced in the papers 45 and 35 cited in the official problem description by Jaffe and Witten

### Module six

The winner must then prove that the mass of the least massive particle of the force field predicted by the theory is (=) strictly positive. **We assume the proposition and delineate and disseminate the model. Should the correctness exist then the prognostication and prediction formulas given at the end of the paper should be correct in consistent with the observation of any data or experimental observation. Lest the converse is true namely, that the force field predicted by the theory is (=) not strictly positive.**

$G_{32}$  : Category one of strictly positive

$G_{33}$  : Category two of strictly positive

$G_{34}$  : Category three of strictly positive

$T_{32}$  : Category one of mass of the least massive particle of the force field predicted by the theory. Systemic differentiation. Kindly note that whatever explanation is given of the predicational anteriorities, character constitution and phenomenological correlates must hold good for all the systems which satisfy the essence of the statement under question.

$T_{33}$  : Category two of mass of the least massive particle of the force field predicted by the theory

$T_{34}$  : Category three of mass of the least massive particle of the force field predicted by the theory

### Module seven

For example, in the case of  $G=SU(3)$  - the strong nuclear interaction - the winner must prove that glueballs **have (e) a lower mass bound. We assume the proposition and give the model. In the next**

**part we assume the inverse and give the results. One of them must hold good.**

$G_{36}$  : Category one of lower mass bound

$G_{37}$  : Category two of lower mass bound

$G_{38}$  : Category three of lower mass bound

$T_{36}$  : Category one of  $G=SU(3)$  - the strong nuclear interaction glueballs

$T_{37}$  : Category two of  $G=SU(3)$  - the strong nuclear interaction glueballs

$T_{38}$  : Category three of  $G=SU(3)$  - the strong nuclear interaction glueballs

### Module eight

Thus glueballs cannot (e) be arbitrarily light

$G_{40}$  : Category one of arbitrarily light

$G_{41}$  : Category two of arbitrarily light

$G_{42}$  : Category three of arbitrarily light

$T_{40}$  : Category one of glueballs

$T_{41}$  : Category two of glueballs

$T_{42}$  : Category three of glueballs

### Module Nine

Yang–Mills theories are a special example of gauge theory with a non-abelian symmetry group given by the Lagrangian

$$\mathcal{L}_{\text{gf}} = -\frac{1}{2} \text{Tr}(F^2) = -\frac{1}{4} F^{a\mu\nu} F_{\mu\nu}^a$$

with the generators of the Lie algebra corresponding to the F-quantities (the curvature or field-strength form) satisfying

$$\text{Tr}(T^a T^b) = \frac{1}{2} \delta^{ab}, \quad [T^a, T^b] = i f^{abc} T^c$$

and the covariant derivative defined as

$$D_\mu = I \partial_\mu - ig T^a A_\mu^a$$

where  $I$  is the identity for the group generators,  $A_\mu^a$  is the vector potential, and  $g$  is the coupling constant. In four dimensions, the coupling constant  $g$  is a pure number and for a  $SU(N)$  group one has  $a, b, c = 1 \dots N^2 - 1$ .

The relation

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + gf^{abc} A_\mu^b A_\nu^c$$

can be derived by the commutator

$$[D_\mu, D_\nu] = -igT^a F_{\mu\nu}^a.$$

The field has the property of being self-interacting and equations of motion that one obtains are said to be semilinear, as nonlinearities are both with and without derivatives. This means that one can manage this theory only by perturbation theory, with small nonlinearities.

Note that the transition between "upper" ("contravariant") and "lower" ("covariant") vector or tensor components is trivial for a indices (e.g.  $f^{abc} = f_{abc}$ ), whereas for  $\mu$  and  $\nu$  it is nontrivial, corresponding e.g. to the usual Lorentz signature,  $\eta_{\mu\nu} = \text{diag}(+ - - -)$ .

From the given Lagrangian one can derive the equations of motion given by

$$\partial^\mu F_{\mu\nu}^a + gf^{abc} A^{\mu b} F_{\mu\nu}^c = 0.$$

Putting  $F_{\mu\nu} = T^a F_{\mu\nu}^a$ , these can be rewritten as

$$(D^\mu F_{\mu\nu})^a = 0.$$

A Bianchi identity holds

$$(D_\mu F_{\nu\kappa})^a + (D_\kappa F_{\mu\nu})^a + (D_\nu F_{\kappa\mu})^a = 0.$$

which is equivalent to the Jacobi identity

$$[D_\mu, [D_\nu, D_\kappa]] + [D_\kappa, [D_\mu, D_\nu]] + [D_\nu, [D_\kappa, D_\mu]] = 0.$$

since  $[D_\mu, F_{\nu\kappa}^a] = -igD_\mu F_{\nu\kappa}^a$ . Define the dual strength tensor  $\tilde{F}^{\mu\nu} = \frac{1}{2}\epsilon^{\mu\nu\rho\sigma} F_{\rho\sigma}$ , then the Bianchi identity can be rewritten as

$$D_\mu \tilde{F}^{\nu\kappa} = 0.$$

A source  $J_\mu^a$  enters into **the equations of motion** as

$$\partial^\mu F_{\mu\nu}^a + gf^{abc} A^{\mu b} F_{\mu\nu}^c = -J_\nu^a.$$

Note that the currents must properly change under gauge group transformations.

We give here some comments about the physical dimensions of the coupling. We note that, in D dimensions, the field scales as  $[A] = [L^{\frac{2-D}{2}}]$  and so the coupling must scale as  $[g^2] = [L^{D-4}]$ . This implies

that Yang–Mills theory is not renormalizable for dimensions greater than four. Further, we note that, for  $D = 4$ , the coupling is dimensionless and both the field and the square of the coupling have the same dimensions of the field and the coupling of a massless quartic scalar field theory. So, these theories share the scale invariance at the classical level.

**Note: When we write A+B, it means that we are adding B to A until B is exhausted. There may be time lag or may not be time lag. It is almost like adding water to milk. When we write B+A it means adding water to milk until water is fully exhausted, which we are familiar. A-B implies removing B from A, with or without time lag. All these commentaries are true for all additions, subtractions, mappings and transformations. In the eventuality of multiplication, logarithms can be taken to separate the variables and hence the terms becomes separate and give results of the prediction for a time t in the model. As said, there are many systems with phenomenological correlates, differential contiguities, presuppositional resemblances and ontological consonance and primordial exactitude. Those systems which are under the scanner can be classified in to three compartments as we have done based on their characteristics. These statements hold good for the entire monograph. We shall not repeat this again. We have done this exercise term by term in earlier papers and shall not repeat the same. Kindly bear with me.**

$G_{44}$  : Category one of LHS of all the equations stated in the foregoing (Yang Mills Theory including the Lagrangian and the Hamiltonian)

$G_{45}$  : Category two of LHS of all the equations stated in the foregoing (Yang Mills Theory including the Lagrangian and the Hamiltonian)

$G_{46}$  : Category three of LHS of all the equations stated in the foregoing (Yang Mills Theory including the Lagrangian and the Hamiltonian)

$T_{44}$  : Category one of RHS of all the equations stated in the foregoing (Yang Mills Theory including the Lagrangian and the Hamiltonian)

$T_{45}$  : Category two of RHS of all the equations stated in the foregoing (Yang Mills Theory including the Lagrangian and the Hamiltonian)

$T_{46}$  : Category three of RHS of all the equations stated in the foregoing (Yang Mills Theory including the Lagrangian and the Hamiltonian)

### The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$ ;  
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$  ,  
 $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}, (a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}$  ,  
 $(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$   
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$   
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$   
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$ ,

$$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}, (a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_2, (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}, (a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}, (a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}, (a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$$

are Dissipation coefficients

**Module Numbered One**

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \end{aligned}$$

$+(a''_{13})^{(1)}(T_{14}, t) =$  First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$  First detritions factor

**Module Numbered Two**

The differential system of this model is now (Module numbered two)

$$\begin{aligned} \frac{dG_{16}}{dt} &= (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} & 7 \\ \frac{dG_{17}}{dt} &= (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} & 8 \\ \frac{dG_{18}}{dt} &= (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} & 9 \\ \frac{dT_{16}}{dt} &= (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} & 10 \\ \frac{dT_{17}}{dt} &= (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} & 11 \\ \frac{dT_{18}}{dt} &= (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} & 12 \end{aligned}$$

$+(a''_{16})^{(2)}(T_{17}, t) =$  First augmentation factor

$-(b''_{16})^{(2)}((G_{19}), t) =$  First detritions factor

**Module Numbered Three**

The differential system of this model is now (Module numbered three)

$$\begin{aligned} \frac{dG_{20}}{dt} &= (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} & 13 \\ \frac{dG_{21}}{dt} &= (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} & 14 \\ \frac{dG_{22}}{dt} &= (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} & 15 \end{aligned}$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$  First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$  First detritions factor

**Module Numbered Four**

**The differential system of this model is now (Module numbered Four)**

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ First augmentation factor

$-(b''_{24})^{(4)}((G_{27}), t) =$ First detritions factor

**Module Numbered Five:**

**The differential system of this model is now (Module number five)**

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ First augmentation factor

$-(b''_{28})^{(5)}((G_{31}), t) =$ First detritions factor

**Module Numbered Six**

**The differential system of this model is now (Module numbered Six)**

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$

**Module Numbered Seven:**

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$

**Module Numbered Eight**

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42} \quad 48$$

**Module Numbered Nine**

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45} \tag{53}$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46} \tag{54}$$

$+(a''_{44})^{(9)}(T_{45}, t) = \text{First augmentation factor}$

$-(b''_{44})^{(9)}((G_{47}), t) = \text{First detrition factor}$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[ \begin{array}{l} (a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) + (a''_{16})^{(2,2)}(T_{17}, t) + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7)}(T_{37}, t) + (a''_{40})^{(8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13} \tag{55}$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[ \begin{array}{l} (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t) + (a''_{17})^{(2,2)}(T_{17}, t) + (a''_{21})^{(3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7)}(T_{37}, t) + (a''_{41})^{(8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14} \tag{56}$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[ \begin{array}{l} (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t) + (a''_{18})^{(2,2)}(T_{17}, t) + (a''_{22})^{(3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7)}(T_{37}, t) + (a''_{42})^{(8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15} \tag{57}$$

Where  $(a''_{13})^{(1)}(T_{14}, t)$ ,  $(a''_{14})^{(1)}(T_{14}, t)$ ,  $(a''_{15})^{(1)}(T_{14}, t)$  are first augmentation coefficients for category 1, 2 and 3

$(a''_{16})^{(2,2)}(T_{17}, t)$ ,  $(a''_{17})^{(2,2)}(T_{17}, t)$ ,  $(a''_{18})^{(2,2)}(T_{17}, t)$  are second augmentation coefficient for category 1, 2 and 3

$(a''_{20})^{(3,3)}(T_{21}, t)$ ,  $(a''_{21})^{(3,3)}(T_{21}, t)$ ,  $(a''_{22})^{(3,3)}(T_{21}, t)$  are third augmentation coefficient for category 1, 2 and 3

$(a''_{24})^{(4,4,4,4)}(T_{25}, t)$ ,  $(a''_{25})^{(4,4,4,4)}(T_{25}, t)$ ,  $(a''_{26})^{(4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficient for category 1, 2 and 3

$(a''_{28})^{(5,5,5,5)}(T_{29}, t)$ ,  $(a''_{29})^{(5,5,5,5)}(T_{29}, t)$ ,  $(a''_{30})^{(5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficient for category 1, 2 and 3

$(a''_{32})^{(6,6,6,6)}(T_{33}, t)$ ,  $(a''_{33})^{(6,6,6,6)}(T_{33}, t)$ ,  $(a''_{34})^{(6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficient for category 1, 2 and 3

$(a''_{38})^{(7,7)}(T_{37}, t)$ ,  $(a''_{37})^{(7,7)}(T_{37}, t)$ ,  $(a''_{36})^{(7,7)}(T_{37}, t)$  are seventh augmentation coefficient for 1,2,3

$(a''_{40})^{(8,8)}(T_{41}, t)$ ,  $(a''_{41})^{(8,8)}(T_{41}, t)$ ,  $(a''_{42})^{(8,8)}(T_{41}, t)$  are eight augmentation coefficient for 1,2,3

$(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ ,  $(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ ,  $(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$  are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[ \begin{array}{l} (b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{16})^{(2,2)}(G_{19}, t) - (b''_{20})^{(3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7)}(G_{39}, t) - (b''_{40})^{(8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \tag{58}$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[ \begin{array}{l} (b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t) - (b''_{17})^{(2,2)}(G_{19}, t) - (b''_{21})^{(3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7)}(G_{39}, t) - (b''_{41})^{(8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14} \tag{59}$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[ \begin{array}{l} (b'_{15})^{(1)} \boxed{-(b''_{15})^{(1)}(G, t)} \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7)}(G_{39}, t)} \boxed{-(b''_{42})^{(8,8)}(G_{43}, t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15} \tag{60}$$

Where  $\boxed{-(b''_{13})^{(1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1)}(G, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2)}(G_{19}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)}$  are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)}$  are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)}$  are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{36})^{(7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7)}(G_{39}, t)}$  are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{41})^{(8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{42})^{(8,8)}(G_{43}, t)}$  are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$  are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[ \begin{array}{l} (a'_{16})^{(2)} \boxed{+(a''_{16})^{(2)}(T_{17}, t)} \boxed{+(a''_{13})^{(1,1)}(T_{14}, t)} \boxed{+(a''_{20})^{(3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37}, t)} \boxed{+(a''_{40})^{(8,8,8)}(T_{41}, t)} \boxed{+(a''_{44})^{(9,9)}(T_{45}, t)} \end{array} \right] G_{16} \tag{61}$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[ \begin{array}{l} (a'_{17})^{(2)} \boxed{+(a''_{17})^{(2)}(T_{17}, t)} \boxed{+(a''_{14})^{(1,1)}(T_{14}, t)} \boxed{+(a''_{21})^{(3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37}, t)} \boxed{+(a''_{41})^{(8,8,8)}(T_{41}, t)} \boxed{+(a''_{45})^{(9,9)}(T_{45}, t)} \end{array} \right] G_{17} \tag{62}$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[ \begin{array}{l} (a'_{18})^{(2)} \boxed{+(a''_{18})^{(2)}(T_{17}, t)} \boxed{+(a''_{15})^{(1,1)}(T_{14}, t)} \boxed{+(a''_{22})^{(3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37}, t)} \boxed{+(a''_{42})^{(8,8,8)}(T_{41}, t)} \boxed{+(a''_{46})^{(9,9)}(T_{45}, t)} \end{array} \right] G_{18} \tag{63}$$

Where  $\boxed{+(a''_{16})^{(2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2)}(T_{17}, t)}$  are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1)}(T_{14}, t)}$  are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3,3)}(T_{21}, t)}$  are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)}$  are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)}$  are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)}$  are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{37})^{(7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{38})^{(7,7,7)}(T_{37}, t)}$  are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{41})^{(8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{42})^{(8,8,8)}(T_{41}, t)}$  are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{45})^{(9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{46})^{(9,9)}(T_{45}, t)}$  are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[ \begin{array}{l} \boxed{(b'_{16})^{(2)} - \boxed{(b''_{16})^{(2)}(G_{19}, t)} - \boxed{(b''_{13})^{(1,1)}(G, t)} - \boxed{(b''_{20})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)} - \boxed{(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)} - \boxed{(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)} - \boxed{(b''_{40})^{(8,8,8)}(G_{43}, t)} - \boxed{(b''_{44})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[ \begin{array}{l} \boxed{(b'_{17})^{(2)} - \boxed{(b''_{17})^{(2)}(G_{19}, t)} - \boxed{(b''_{14})^{(1,1)}(G, t)} - \boxed{(b''_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} - \boxed{(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} - \boxed{(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)} - \boxed{(b''_{41})^{(8,8,8)}(G_{43}, t)} - \boxed{(b''_{45})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[ \begin{array}{l} \boxed{(b'_{18})^{(2)} - \boxed{(b''_{18})^{(2)}(G_{19}, t)} - \boxed{(b''_{15})^{(1,1)}(G, t)} - \boxed{(b''_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} - \boxed{(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} - \boxed{(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)} - \boxed{(b''_{42})^{(8,8,8)}(G_{43}, t)} - \boxed{(b''_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \quad 66$$

where  $\boxed{-(b''_{16})^{(2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2)}(G_{19}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1)}(G, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$  are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$  are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$  are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)}$  are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)}$  are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{46})^{(9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{45})^{(9,9)}(G_{47}, t)}$  are ninth detrition coefficients for category 1, 2 and 3

1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[ \begin{array}{l} (a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) + (a''_{16})^{(2,2,2)}(T_{17}, t) + (a''_{13})^{(1,1,1)}(T_{14}, t) \\ + (a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{20} \quad 67$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)} G_{20} - \left[ \begin{array}{l} (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) + (a''_{17})^{(2,2,2)}(T_{17}, t) + (a''_{14})^{(1,1,1)}(T_{14}, t) \\ + (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)} G_{21} - \left[ \begin{array}{l} (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) + (a''_{18})^{(2,2,2)}(T_{17}, t) + (a''_{15})^{(1,1,1)}(T_{14}, t) \\ + (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9)}(T_{45}, t) \end{array} \right] G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21}, t), +(a''_{21})^{(3)}(T_{21}, t), +(a''_{22})^{(3)}(T_{21}, t)$  are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2)}(T_{17}, t)$  are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1)}(T_{14}, t)$  are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t), +(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t), +(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7)}(T_{37}, t)$  are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8)}(T_{41}, t)$  are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9)}(T_{45}, t)$  are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)} T_{21} - \left[ \begin{array}{l} (b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{16})^{(2,2,2)}(G_{19}, t) - (b''_{13})^{(1,1,1)}(G, t) \\ - (b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)} T_{20} - \left[ \begin{array}{l} (b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t) - (b''_{17})^{(2,2,2)}(G_{19}, t) - (b''_{14})^{(1,1,1)}(G, t) \\ - (b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[ \begin{array}{l} (b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t) - (b''_{18})^{(2,2,2)}(G_{19}, t) - (b''_{15})^{(1,1,1)}(G, t) \\ - (b''_{26})^{(4,4,4,4,4)}(G_{27}, t) - (b''_{30})^{(5,5,5,5,5)}(G_{31}, t) - (b''_{34})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b''_{38})^{(7,7,7,7)}(G_{39}, t) - (b''_{42})^{(8,8,8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9)}(G_{47}, t) \end{array} \right] T_{22} \quad 72$$

$-(b''_{20})^{(3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3)}(G_{23}, t)$  are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2,2,2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2,2,2)}(G_{19}, t)$  are second detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1,1)}(G, t)$ ,  $-(b''_{15})^{(1,1,1)}(G, t)$  are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)$  are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)$  are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)$  are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{37})^{(7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{38})^{(7,7,7,7)}(G_{39}, t)$  are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{41})^{(8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{42})^{(8,8,8,8)}(G_{43}, t)$  are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47}, t)$ ,  $-(b''_{45})^{(9,9,9)}(G_{47}, t)$ ,  $-(b''_{44})^{(9,9,9)}(G_{47}, t)$  are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[ \begin{array}{l} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) + (a''_{28})^{(5,5)}(T_{29}, t) + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[ \begin{array}{l} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) + (a''_{29})^{(5,5)}(T_{29}, t) + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[ \begin{array}{l} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) + (a''_{30})^{(5,5)}(T_{29}, t) + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t)$ ,  $(a''_{25})^{(4)}(T_{25}, t)$ ,  $(a''_{26})^{(4)}(T_{25}, t)$  are first augmentation coefficients category 1, 2 3

$+(a''_{28})^{(5,5)}(T_{29}, t)$ ,  $+(a''_{29})^{(5,5)}(T_{29}, t)$ ,  $+(a''_{30})^{(5,5)}(T_{29}, t)$  are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6)}(T_{33}, t)$ ,  $+(a''_{33})^{(6,6)}(T_{33}, t)$ ,  $+(a''_{34})^{(6,6)}(T_{33}, t)$  are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1)}(T_{14}, t)$ ,  $+(a''_{14})^{(1,1,1,1)}(T_{14}, t)$ ,  $+(a''_{15})^{(1,1,1,1)}(T_{14}, t)$  are fourth augmentation coefficient

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17}, t)}$ ,  
 $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17}, t)}$  are fifth augmentation coefficients for category 1, 2 and 3  
 $\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21}, t)}$ ,  
 $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21}, t)}$  are sixth augmentation coefficients for category 1, 2 and 3  
 $\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)}$ ,  
 $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37}, t)}$  are seventh augmentation coefficients for category 1, 2 and 3  
 $\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41}, t)}$   
 are eighth augmentation coefficients for category 1, 2 and 3  
 $\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45}, t)}$  are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left. \begin{array}{l} \boxed{(b'_{24})^{(4)} - \boxed{-(b''_{24})^{(4)}(G_{27}, t)} - \boxed{-(b''_{28})^{(5,5)}(G_{31}, t)} - \boxed{-(b''_{32})^{(6,6)}(G_{35}, t)}} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G, t)} - \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19}, t)} - \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39}, t)} - \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43}, t)} - \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47}, t)} \end{array} \right\} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left. \begin{array}{l} \boxed{(b'_{25})^{(4)} - \boxed{-(b''_{25})^{(4)}(G_{27}, t)} - \boxed{-(b''_{29})^{(5,5)}(G_{31}, t)} - \boxed{-(b''_{33})^{(6,6)}(G_{35}, t)}} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G, t)} - \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19}, t)} - \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39}, t)} - \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43}, t)} - \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47}, t)} \end{array} \right\} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left. \begin{array}{l} \boxed{(b'_{26})^{(4)} - \boxed{-(b''_{26})^{(4)}(G_{27}, t)} - \boxed{-(b''_{30})^{(5,5)}(G_{31}, t)} - \boxed{-(b''_{34})^{(6,6)}(G_{35}, t)}} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G, t)} - \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19}, t)} - \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39}, t)} - \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43}, t)} - \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47}, t)} \end{array} \right\} T_{26} \quad 78$$

Where  $\boxed{-(b''_{24})^{(4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4)}(G_{27}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5)}(G_{31}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6)}(G_{35}, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1,1)}(G, t)}$ ,  
 $\boxed{-(b''_{15})^{(1,1,1,1)}(G, t)}$  are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2,2)}(G_{19}, t)}$ ,  
 $\boxed{-(b''_{18})^{(2,2,2,2)}(G_{19}, t)}$  are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3,3)}(G_{23}, t)}$  are sixth detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39}, t)}$

$\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39}, t)}$  are seventh detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43}, t)}$

are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47}, t)}$  are ninth detrition coefficients for category 1, 2 and 3

category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[ \begin{array}{l} (a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) + (a''_{24})^{(4,4)}(T_{25}, t) + (a''_{32})^{(6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[ \begin{array}{l} (a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t) + (a''_{25})^{(4,4)}(T_{25}, t) + (a''_{33})^{(6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[ \begin{array}{l} (a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t) + (a''_{26})^{(4,4)}(T_{25}, t) + (a''_{34})^{(6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{30} \quad 81$$

Where  $+(a''_{28})^{(5)}(T_{29}, t)$ ,  $+(a''_{29})^{(5)}(T_{29}, t)$ ,  $+(a''_{30})^{(5)}(T_{29}, t)$  are first augmentation coefficients for category 1, 2 and 3

And  $+(a''_{24})^{(4,4)}(T_{25}, t)$ ,  $+(a''_{25})^{(4,4)}(T_{25}, t)$ ,  $+(a''_{26})^{(4,4)}(T_{25}, t)$  are second augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6)}(T_{33}, t)$ ,  $+(a''_{33})^{(6,6,6)}(T_{33}, t)$ ,  $+(a''_{34})^{(6,6,6)}(T_{33}, t)$  are third augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)$ ,  $+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)$ ,  $+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)$  are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)$  are fifth augmentation coefficients for category 1,2, and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)$ ,  $+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)$ ,  $+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)$  are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)$ ,  $+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)$ ,  $+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)$  are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)$ ,  $+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)$ ,  $+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)$  are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)$ ,  $+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)$ ,  $+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)$  are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[ \begin{array}{l} (b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t) - (b''_{24})^{(4,4)}(G_{27}, t) - (b''_{32})^{(6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1)}(G, t) - (b''_{16})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{20})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[ \begin{array}{l} (b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t) - (b''_{25})^{(4,4)}(G_{27}, t) - (b''_{33})^{(6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1)}(G, t) - (b''_{17})^{(2,2,2,2,2)}(G_{19}, t) - (b''_{21})^{(3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[ \begin{array}{l} (b'_{30})^{(5)} \boxed{-(b''_{30})^{(5)}(G_{31}, t)} \boxed{-(b''_{26})^{(4,4)}(G_{27}, t)} \boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)} \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)} \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)} \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)} \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{30} \tag{84}$$

where  $\boxed{-(b''_{28})^{(5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5)}(G_{31}, t)}$  are first detrition coefficients for category 1,2 and 3  
 $\boxed{-(b''_{24})^{(4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4)}(G_{27}, t)}$  are second detrition coefficients for category 1,2 and 3  
 $\boxed{-(b''_{32})^{(6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)}$  are third detrition coefficients for category 1,2 and 3  
 $\boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)}$  are fourth detrition coefficients for category 1,2, and 3  
 $\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)}$  are fifth detrition coefficients for category 1,2, and 3  
 $\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)}$  are sixth detrition coefficients for category 1,2, and 3  
 $\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)}$  are seventh detrition coefficients for category 1,2, and 3  
 $\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)}$  are eighth detrition coefficients for category 1,2, and 3  
 $\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)}$  are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[ \begin{array}{l} (a'_{32})^{(6)} \boxed{+(a''_{32})^{(6)}(T_{33}, t)} \boxed{+(a''_{28})^{(5,5,5)}(T_{29}, t)} \boxed{+(a''_{24})^{(4,4,4)}(T_{25}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)} \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)} \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)} \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)} \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{32} \tag{85}$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[ \begin{array}{l} (a'_{33})^{(6)} \boxed{+(a''_{33})^{(6)}(T_{33}, t)} \boxed{+(a''_{29})^{(5,5,5)}(T_{29}, t)} \boxed{+(a''_{25})^{(4,4,4)}(T_{25}, t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)} \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)} \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)} \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)} \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{33} \tag{86}$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[ \begin{array}{l} (a'_{34})^{(6)} \boxed{+(a''_{34})^{(6)}(T_{33}, t)} \boxed{+(a''_{30})^{(5,5,5)}(T_{29}, t)} \boxed{+(a''_{26})^{(4,4,4)}(T_{25}, t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)} \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)} \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)} \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)} \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{34} \tag{87}$$

$\boxed{+(a''_{32})^{(6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6)}(T_{33}, t)}$  are first augmentation coefficients for category 1, 2 and 3  
 $\boxed{+(a''_{28})^{(5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5)}(T_{29}, t)}$  are second augmentation coefficients for category 1, 2 and 3  
 $\boxed{+(a''_{24})^{(4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4)}(T_{25}, t)}$  are third augmentation coefficients for category 1, 2 and 3  
 $\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)}$  - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)}$  - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)}$  sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t)}$ ,  
 $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)}$  seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)}$

Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)}$  ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[ \begin{array}{l} \boxed{(b'_{32})^{(6)} - \boxed{-(b''_{32})^{(6)}(G_{35}, t)} - \boxed{-(b''_{28})^{(5,5,5)}(G_{31}, t)} - \boxed{-(b''_{24})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G, t)} - \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)} - \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)} - \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)} - \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[ \begin{array}{l} \boxed{(b'_{33})^{(6)} - \boxed{-(b''_{33})^{(6)}(G_{35}, t)} - \boxed{-(b''_{29})^{(5,5,5)}(G_{31}, t)} - \boxed{-(b''_{25})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G, t)} - \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)} - \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)} - \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)} - \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[ \begin{array}{l} \boxed{(b'_{34})^{(6)} - \boxed{-(b''_{34})^{(6)}(G_{35}, t)} - \boxed{-(b''_{30})^{(5,5,5)}(G_{31}, t)} - \boxed{-(b''_{26})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G, t)} - \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)} - \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)} - \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)} - \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6)}(G_{35}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5)}(G_{31}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4)}(G_{27}, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G, t)}$  are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)}$  are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)}$  are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)}$  are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)}$

are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)}$  are ninth detrition

coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)} G_{37} \tag{91}$$

$$- \left[ \begin{array}{l} (a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) + (a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)} G_{36} \tag{92}$$

$$- \left[ \begin{array}{l} (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t) + (a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} \tag{93}$$

$$- \left[ \begin{array}{l} (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t) + (a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where  $(a''_{36})^{(7)}(T_{37}, t)$ ,  $(a''_{37})^{(7)}(T_{37}, t)$ ,  $(a''_{38})^{(7)}(T_{37}, t)$  are first augmentation coefficients for category 1, 2 and 3

$(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)$  are second augmentation coefficient for category 1, 2 and 3

$(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ ,  $(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)$ ,  $(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)$  are third augmentation coefficient for category 1, 2 and 3

$(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ ,  $(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)$ ,  $(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficient for category 1, 2 and 3

$(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ ,  $(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)$ ,  $(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficient for category 1, 2 and 3

$(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ ,  $(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)$ ,  $(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficient for category 1, 2 and 3

$(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ ,  $(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)$ ,  $(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)$  are seventh augmentation coefficient for category 1, 2 and 3

$(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ ,  $(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)$ ,  $(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)$  are eighth augmentation coefficient for 1,2,3

$(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ ,  $(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)$ ,  $(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)$  are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - \left[ \begin{array}{l} (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}, t) - (b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) - (b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1)}(G, t) - (b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13} \tag{94}$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[ \begin{array}{l} (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}, t) - (b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) - (b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1)}(G, t) - (b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[ \begin{array}{l} (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}, t) - (b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) - (b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) - (b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) - (b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1)}(G, t) - (b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) - (b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where  $-(b''_{36})^{(7)}(G_{39}, t)$ ,  $-(b''_{37})^{(7)}(G_{39}, t)$ ,  $-(b''_{38})^{(7)}(G_{39}, t)$  are first detrition coefficients for category 1, 2 and 3  
 $-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)$  are second detrition coefficients for category 1, 2 and 3  
 $-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)$  are third detrition coefficients for category 1, 2 and 3  
 $-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)$  are fourth detrition coefficients for category 1, 2 and 3  
 $-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)$  are fifth detrition coefficients for category 1, 2 and 3  
 $-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)$  are sixth detrition coefficients for category 1, 2 and 3  
 $-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)$  are seventh detrition coefficients for category 1, 2 and 3  
 $-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)$  are eighth detrition coefficients for category 1, 2 and 3  
 $-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)$  are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} \tag{95}$$

$$= (a_{40})^{(8)}G_{41} - \left[ \begin{array}{l} (a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) + (a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)}G_{40} - \left[ \begin{array}{l} (a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t) + (a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[ \begin{array}{l} (a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t) + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{15}$$

Where  $(a'_{40})^{(8)}(T_{41}, t)$ ,  $(a''_{41})^{(8)}(T_{41}, t)$ ,  $(a''_{42})^{(8)}(T_{41}, t)$  are first augmentation coefficients for category 1, 2 and 3

$(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$  are second augmentation coefficient for category 1, 2 and 3

$(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ ,  $(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$ ,  $(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t)$  are third augmentation coefficient for category 1, 2 and 3

$(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ ,  $(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$ ,  $(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficient for category 1, 2 and 3

$(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ ,  $(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$ ,  $(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficient for category 1, 2 and 3

$(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ ,  $(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$ ,  $(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficient for category 1, 2 and 3

$(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ ,  $(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$ ,  $(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t)$  are seventh augmentation coefficient for 1,2,3

$(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ ,  $(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$ ,  $(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t)$  are eighth augmentation coefficient for 1,2,3

$(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ ,  $(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$ ,  $(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)$  are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[ \begin{array}{l} (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}, t) - (b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) - (b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) - (b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) - (b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[ \begin{array}{l} (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}, t) - (b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) - (b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) - (b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) - (b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[ \begin{array}{l} (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}, t) - (b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) - (b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) - (b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) - (b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ - (b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) - (b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) - (b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{15}$$

Where  $-(b''_{36})^{(7)}(G_{39}, t)$ ,  $-(b''_{37})^{(7)}(G_{39}, t)$ ,  $-(b''_{38})^{(7)}(G_{39}, t)$  are first detrition coefficients for category 1, 2 and 3  
 $-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$  are second detrition coefficients for category 1, 2 and 3  
 $-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$  are third detrition coefficients for category 1, 2 and 3  
 $-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$  are fourth detrition coefficients for category 1, 2 and 3  
 $-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$  are fifth detrition coefficients for category 1, 2 and 3  
 $-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$  are sixth detrition coefficients for category 1, 2 and 3  
 $-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{38})^{(7,7)}(G_{39}, t)$  are seventh detrition coefficients for category 1, 2 and 3  
 $-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$  are eighth detrition coefficients for category 1, 2 and 3  
 $-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$  are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[ \begin{array}{l} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) + (a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[ \begin{array}{l} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) + (a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[ \begin{array}{l} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) + (a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where  $+(a''_{44})^{(9)}(T_{45}, t)$ ,  $+(a''_{45})^{(9)}(T_{45}, t)$ ,  $+(a''_{46})^{(9)}(T_{37}, t)$  are first augmentation coefficients for category 1, 2 and 3  
 $+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17}, t)$  are second

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)}$$
 are third

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)}$$
 are fourth

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)}$$
 are fifth

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)}$$
 are sixth

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)}$$
 are Seventh

augmentation coefficient for category 1, 2 and 3

$$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)}, \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)}$$
 are eighth

augmentation coefficient for 1,2,3

$$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)}$$
 are ninth

augmentation coefficient for 1,2,3

$$\begin{aligned} & \frac{dT_{44}}{dt} \\ &= (b_{44})^{(9)}T_{45} \\ & - \left[ \begin{array}{l} \boxed{(b'_{44})^{(9)} - \boxed{(b''_{44})^{(9)}(G_{47}, t)} - \boxed{(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19}, t)} - \boxed{(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27}, t)} - \boxed{(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31}, t)} - \boxed{(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G, t)} - \boxed{(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39}, t)} - \boxed{(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{13} \\ & \frac{dT_{45}}{dt} \\ &= (b_{45})^{(9)}T_{44} - \left[ \begin{array}{l} \boxed{(b'_{45})^{(9)} - \boxed{(b''_{45})^{(9)}(G_{47}, t)} - \boxed{(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19}, t)} - \boxed{(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27}, t)} - \boxed{(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31}, t)} - \boxed{(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G, t)} - \boxed{(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39}, t)} - \boxed{(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{14} \\ & \frac{dT_{46}}{dt} \\ &= (b_{46})^{(9)}T_{45} - \left[ \begin{array}{l} \boxed{(b'_{46})^{(9)} - \boxed{(b''_{46})^{(9)}(G_{47}, t)} - \boxed{(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19}, t)} - \boxed{(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27}, t)} - \boxed{(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31}, t)} - \boxed{(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G, t)} - \boxed{(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39}, t)} - \boxed{(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43}, t)} \end{array} \right] T_{15} \end{aligned}$$

Where  $\boxed{(b''_{44})^{(9)}(G_{47}, t)}$ ,  $\boxed{(b''_{45})^{(9)}(G_{47}, t)}$ ,  $\boxed{(b''_{46})^{(9)}(G_{47}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23}, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27}, t)}$  are fourth detrition

coefficients for category 1, 2 and 3

$$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)}$$
 are fifth detrition

coefficients for category 1, 2 and 3

$$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t)}$$
 are sixth detrition

coefficients for category 1, 2 and 3

$$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)}, \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)}$$
 are seventh detrition

coefficients for category 1, 2 and 3

$$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)}, \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)}$$
 are eighth detrition

coefficients for category 1, 2 and 3

$$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)}, \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43}, t)}$$
 are ninth

detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i'')^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, i, j = 13,14,15 \tag{97}$$

The functions  $(a_i'')^{(1)}, (b_i'')^{(1)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(1)}, (r_i)^{(1)}$ :

$$\begin{aligned} (a_i'')^{(1)}(T_{14}, t) &\leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)} \\ (b_i'')^{(1)}(G, t) &\leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)} \\ \lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) &= (p_i)^{(1)} \\ \lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) &= (r_i)^{(1)} \end{aligned} \tag{98}$$

**Definition of**  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$  :

Where  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$  are positive constants and  $i = 13,14,15$

They satisfy Lipschitz condition: 99

$$\begin{aligned} |(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| &\leq (\hat{k}_{13})^{(1)} |T'_{14} - T_{14}| e^{-(\hat{M}_{13})^{(1)}t} \\ |(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| &< (\hat{k}_{13})^{(1)} \|G - G'\| e^{-(\hat{M}_{13})^{(1)}t} \end{aligned}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(1)}(T'_{14}, t)$  and  $(a_i'')^{(1)}(T_{14}, t)$ .  $(T'_{14}, t)$  and  $(T_{14}, t)$  are points belonging to the interval  $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$ . It is to be noted that  $(a_i'')^{(1)}(T_{14}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{13})^{(1)} = 1$  then the function  $(a_i'')^{(1)}(T_{14}, t)$ , the first augmentation coefficient attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$  : 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$ , are positive constants

$$\frac{(a_i)^{(1)}}{(\widehat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}} < 1$$

**Definition of**  $(\widehat{P}_{13})^{(1)}, (\widehat{Q}_{13})^{(1)}$  : 101

There exists two constants  $(\widehat{P}_{13})^{(1)}$  and  $(\widehat{Q}_{13})^{(1)}$  which together With  $(\widehat{M}_{13})^{(1)}, (\widehat{k}_{13})^{(1)}, (\widehat{A}_{13})^{(1)}$  and  $(\widehat{B}_{13})^{(1)}$  and the constants  $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13,14,15$ , satisfy the inequalities

$$\frac{1}{(\widehat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\widehat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\widehat{B}_{13})^{(1)} + (\widehat{Q}_{13})^{(1)} (\widehat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a'_i)^{(2)}, (a''_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (b''_i)^{(2)} > 0, i, j = 16,17,18$$

The functions  $(a''_i)^{(2)}, (b''_i)^{(2)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(2)}, (r_i)^{(2)}$ :

$$(a''_i)^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\widehat{A}_{16})^{(2)} \tag{102}$$

$$(b''_i)^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b'_i)^{(2)} \leq (\widehat{B}_{16})^{(2)} \tag{103}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(2)}(T_{17}, t) = (p_i)^{(2)} \tag{104}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(2)}(G_{19}, t) = (r_i)^{(2)} \tag{105}$$

**Definition of**  $(\widehat{A}_{16})^{(2)}, (\widehat{B}_{16})^{(2)}$  : 106

Where  $(\widehat{A}_{16})^{(2)}, (\widehat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$  are positive constants and  $i = 16,17,18$

They satisfy Lipschitz condition:

$$|(a''_i)^{(2)}(T'_{17}, t) - (a''_i)^{(2)}(T_{17}, t)| \leq (\widehat{k}_{16})^{(2)} |T_{17} - T'_{17}| e^{-(\widehat{M}_{16})^{(2)}t} \tag{107}$$

$$|(b''_i)^{(2)}(G'_{19}, t) - (b''_i)^{(2)}(G_{19}, t)| < (\widehat{k}_{16})^{(2)} |(G_{19}) - (G'_{19})'| e^{-(\widehat{M}_{16})^{(2)}t} \tag{108}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(2)}(T'_{17}, t)$  and  $(a''_i)^{(2)}(T_{17}, t)$ .  $(T'_{17}, t)$  and  $(T_{17}, t)$  are points belonging to the interval  $[(\widehat{k}_{16})^{(2)}, (\widehat{M}_{16})^{(2)}]$ . It is to be noted that  $(a''_i)^{(2)}(T_{17}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\widehat{M}_{16})^{(2)} = 1$  then the function  $(a''_i)^{(2)}(T_{17}, t)$ , the first augmentation coefficient attributable to the system, would be absolutely continuous.

**Definition of**  $(\widehat{M}_{16})^{(2)}, (\widehat{k}_{16})^{(2)}$  :

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$ , are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

**Definition of**  $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$  :

There exists two constants  $(\hat{P}_{16})^{(2)}$  and  $(\hat{Q}_{16})^{(2)}$  which together with  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$  and  $(\hat{B}_{16})^{(2)}$  and the constants  $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18$ ,

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [ (a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)} ] < 1 \tag{110}$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [ (b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)} ] < 1 \tag{111}$$

Where we suppose

$$(a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, i, j = 20, 21, 22 \tag{112}$$

The functions  $(a''_i)^{(3)}, (b''_i)^{(3)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(3)}, (r_i)^{(3)}$ :

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \tag{113}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

**Definition of**  $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$  :

Where  $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$  are positive constants and  $i = 20, 21, 22$

They satisfy Lipschitz condition: 114

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T'_{21} - T_{21}| e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b''_i)^{(3)}(G'_{23}, t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} |G'_{23} - G_{23}| e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(3)}(T'_{21}, t)$  and  $(a''_i)^{(3)}(T_{21}, t)$ .  $(T'_{21}, t)$  and  $(T_{21}, t)$  are points belonging to the interval  $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$ . It is to be noted that  $(a''_i)^{(3)}(T_{21}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{20})^{(3)} = 1$  then the function  $(a''_i)^{(3)}(T_{21}, t)$ , the first augmentation coefficient attributable to the system, would

be absolutely continuous.

**Definition of**  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$  : 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$ , are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants  $(\hat{P}_{20})^{(3)}$  and  $(\hat{Q}_{20})^{(3)}$  which together with  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$  and  $(\hat{B}_{20})^{(3)}$  and the constants  $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$ , satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, i, j = 24, 25, 26 \quad 117$$

The functions  $(a''_i)^{(4)}, (b''_i)^{(4)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(4)}, (r_i)^{(4)}$ :

$$(a''_i)^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b''_i)^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b'_i)^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad 118$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

**Definition of**  $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$  :

Where  $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$  are positive constants and  $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$|(a''_i)^{(4)}(T'_{25}, t) - (a''_i)^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T'_{25} - T_{25}| e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b''_i)^{(4)}((G'_{27}), t) - (b''_i)^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} |(G'_{27}) - (G_{27})| e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(4)}(T'_{25}, t)$  and  $(a''_i)^{(4)}(T_{25}, t)$ .  $(T'_{25}, t)$  and  $(T_{25}, t)$  are points belonging to the interval  $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$ . It is to be noted that  $(a''_i)^{(4)}(T_{25}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{24})^{(4)} = 1$  then the function  $(a''_i)^{(4)}(T_{25}, t)$ , the **first augmentation coefficient** attributable to the system, would

be absolutely continuous.

**Definition of**  $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$  : 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$ , are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} , \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

**Definition of**  $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$  : 121

There exists two constants  $(\hat{P}_{24})^{(4)}$  and  $(\hat{Q}_{24})^{(4)}$  which together with  $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$  and  $(\hat{B}_{24})^{(4)}$  and the constants  $(a_i)^{(4)}, (a'_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a'_i)^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, i, j = 28, 29, 30 \quad 122$$

The functions  $(a''_i)^{(5)}, (b''_i)^{(5)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(5)}, (r_i)^{(5)}$ :

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}(G_{31}, t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)} \quad 123$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

**Definition of**  $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$  :

Where  $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$  are positive constants and  $i = 28, 29, 30$

They satisfy Lipschitz condition: 124

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T'_{29} - T_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b''_i)^{(5)}(G'_{31}, t) - (b''_i)^{(5)}(G_{31}, t)| < (\hat{k}_{28})^{(5)} |(G'_{31}) - (G_{31})| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(5)}(T'_{29}, t)$  and  $(a''_i)^{(5)}(T_{29}, t)$ .  $(T'_{29}, t)$  and  $(T_{29}, t)$  are points belonging to the interval  $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$ . It is to

be noted that  $(a_i'')^{(5)}(T_{29}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{28})^{(5)} = 1$  then the function  $(a_i'')^{(5)}(T_{29}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$  : 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$ , are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

**Definition of**  $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$  : 126

There exists two constants  $(\hat{P}_{28})^{(5)}$  and  $(\hat{Q}_{28})^{(5)}$  which together with  $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$  and  $(\hat{B}_{28})^{(5)}$  and the constants  $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, i, j = 32, 33, 34 \quad 127$$

The functions  $(a_i'')^{(6)}, (b_i'')^{(6)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(6)}, (r_i)^{(6)}$ :

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)} \quad 128$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

**Definition of**  $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$  :

Where  $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$  are positive constants and  $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T_{33}'| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} |(G_{35}) - (G_{35})'| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(6)}(T_{33}, t)$  and  $(a_i'')^{(6)}(T_{33}, t) \cdot (T_{33}, t)$  and  $(T_{33}, t)$  are points belonging to the interval  $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$ . It is to be noted that  $(a_i'')^{(6)}(T_{33}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{32})^{(6)} = 1$  then the function  $(a_i'')^{(6)}(T_{33}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$  : 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$ , are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}} , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

**Definition of**  $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$  : 130

There exists two constants  $(\hat{P}_{32})^{(6)}$  and  $(\hat{Q}_{32})^{(6)}$  which together with  $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$  and  $(\hat{B}_{32})^{(6)}$  and the constants  $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [ (a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)} ] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [ (b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)} ] < 1$$

Where we suppose

(A)  $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, i, j = 36, 37, 38$  131

(B) The functions  $(a_i'')^{(7)}, (b_i'')^{(7)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(7)}, (r_i)^{(7)}$ :

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (\hat{B}_{36})^{(7)}$$

(C)  $\lim_{T_2 \rightarrow \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

(D)

$$\lim_{G \rightarrow \infty} (b_i'')^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

132

**Definition of**  $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$  :

Where  $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}$  are positive constants and  $i = 36, 37, 38$

They satisfy Lipschitz condition:

133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37}' - T_{37}| e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} \| (G_{39})' - (G_{39}) \| e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(7)}(T_{37}', t)$  and  $(a_i'')^{(7)}(T_{37}, t)$ .  $(T_{37}', t)$  and  $(T_{37}, t)$  are points belonging to the interval  $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$ . It is to be noted that  $(a_i'')^{(7)}(T_{37}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{36})^{(7)} = 1$  then the function  $(a_i'')^{(7)}(T_{37}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$  :

134

(E)  $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$ , are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

**Definition of**  $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$  :

135

(F) There exists two constants  $(\hat{P}_{36})^{(7)}$  and  $(\hat{Q}_{36})^{(7)}$  which together with  $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$  and  $(\hat{B}_{36})^{(7)}$  and the constants  $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36, 37, 38$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [ (a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)} ] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [ (b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)} ] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, i, j = 40, 41, 42$$

136

The functions  $(a_i'')^{(8)}, (b_i'')^{(8)}$  are positive continuous increasing and bounded

**Definition of**  $(p_i)^{(8)}, (r_i)^{(8)}$  :

137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$$

138

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$$

139

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$$

140

$$\lim_{G \rightarrow \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \tag{141}$$

**Definition of**  $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$  :

Where  $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$  are positive constants and  $i = 40, 41, 42$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41}' - T_{41}| e^{-(\hat{M}_{40})^{(8)}t} \tag{142}$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} |(G_{43})' - (G_{43})| e^{-(\hat{M}_{40})^{(8)}t} \tag{143}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(8)}(T_{41}', t)$  and  $(a_i'')^{(8)}(T_{41}, t)$ .  $(T_{41}', t)$  and  $(T_{41}, t)$  are points belonging to the interval  $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$ . It is to be noted that  $(a_i'')^{(8)}(T_{41}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{40})^{(8)} = 1$  then the function  $(a_i'')^{(8)}(T_{41}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$  :

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$ , are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \tag{144}$$

**Definition of**  $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$  :

There exists two constants  $(\hat{P}_{40})^{(8)}$  and  $(\hat{Q}_{40})^{(8)}$  which together with  $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$  and the constants  $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$ , Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \tag{145}$$

$$\frac{1}{(\hat{M}_{40})^{(8)}} [(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \tag{146}$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, i, j = 44, 45, 46 \tag{146}$$

A

The functions  $(a_i'')^{(9)}, (b_i'')^{(9)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(9)}, (r_i)^{(9)}$ :

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

**Definition of**  $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$  :

Where  $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$  are positive constants and  $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45}' - T_{45}| e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} |(G_{47})' - (G_{47})| e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(9)}(T_{45}', t)$  and  $(a_i'')^{(9)}(T_{45}, t)$ .  $(T_{45}', t)$  and  $(T_{45}, t)$  are points belonging to the interval  $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$ . It is to be noted that  $(a_i'')^{(9)}(T_{45}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{44})^{(9)} = 1$  then the function  $(a_i'')^{(9)}(T_{45}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$  :

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$ , are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}} , \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

**Definition of**  $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$  :

There exists two constants  $(\hat{P}_{44})^{(9)}$  and  $(\hat{Q}_{44})^{(9)}$  which together with  $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)}$  and  $(\hat{B}_{44})^{(9)}$  and the constants  $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

**Theorem 1:** if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} , \quad T_i(0) = T_i^0 > 0$$

**Theorem 2 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

**Definition of**  $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad T_i(0) = T_i^0 > 0$$

**Theorem 3 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}, \quad T_i(0) = T_i^0 > 0$$

**Theorem 4 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 5 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 6 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 7 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}, \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 8 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t} , \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t} , \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 9:** if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t} , \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t} , \boxed{T_i(0) = T_i^0 > 0}$$

**Proof:** Consider operator  $\mathcal{A}^{(1)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy 154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[ (a_{13})^{(1)} G_{14}(s_{(13)}) - \left( (a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[ (a_{14})^{(1)} G_{13}(s_{(13)}) - \left( (a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[ (a_{15})^{(1)} G_{14}(s_{(13)}) - \left( (a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[ (b_{13})^{(1)} T_{14}(s_{(13)}) - \left( (b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[ (b_{14})^{(1)} T_{13}(s_{(13)}) - \left( (b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[ (b_{15})^{(1)} T_{14}(s_{(13)}) - \left( (b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where  $s_{(13)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

159

Consider operator  $\mathcal{A}^{(2)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By

160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[ (a_{16})^{(2)} G_{17}(s_{(16)}) - \left( (a'_{16})^{(2)} + a''_{16} \right)^{(2)} (T_{17}(s_{(16)}), s_{(16)}) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[ (a_{17})^{(2)} G_{16}(s_{(16)}) - \left( (a'_{17})^{(2)} + a''_{17} \right)^{(2)} (T_{17}(s_{(16)}), s_{(17)}) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[ (a_{18})^{(2)} G_{17}(s_{(16)}) - \left( (a'_{18})^{(2)} + a''_{18} \right)^{(2)} (T_{17}(s_{(16)}), s_{(16)}) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[ (b_{16})^{(2)} T_{17}(s_{(16)}) - \left( (b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[ (b_{17})^{(2)} T_{16}(s_{(16)}) - \left( (b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[ (b_{18})^{(2)} T_{17}(s_{(16)}) - \left( (b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where  $s_{(16)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(3)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[ (a_{20})^{(3)} G_{21}(s_{(20)}) - \left( (a'_{20})^{(3)} + a''_{20} \right)^{(3)} (T_{21}(s_{(20)}), s_{(20)}) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[ (a_{21})^{(3)} G_{20}(s_{(20)}) - \left( (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[ (a_{22})^{(3)} G_{21}(s_{(20)}) - \left( (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[ (b_{20})^{(3)} T_{21}(s_{(20)}) - \left( (b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[ (b_{21})^{(3)} T_{20}(s_{(20)}) - \left( (b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[ (b_{22})^{(3)} T_{21}(s_{(20)}) - \left( (b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where  $s_{(20)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:** Consider operator  $\mathcal{A}^{(4)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[ (a_{24})^{(4)} G_{25}(s_{(24)}) - \left( (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[ (a_{25})^{(4)} G_{24}(s_{(24)}) - \left( (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[ (a_{26})^{(4)} G_{25}(s_{(24)}) - \left( (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[ (b_{24})^{(4)} T_{25}(s_{(24)}) - \left( (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[ (b_{25})^{(4)} T_{24}(s_{(24)}) - \left( (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[ (b_{26})^{(4)} T_{25}(s_{(24)}) - \left( (b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where  $s_{(24)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:** Consider operator  $\mathcal{A}^{(5)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[ (a_{28})^{(5)} G_{29}(s_{(28)}) - \left( (a'_{28})^{(5)} + a''_{28} \right)^{(5)} (T_{29}(s_{(28)}), s_{(28)}) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[ (a_{29})^{(5)} G_{28}(s_{(28)}) - \left( (a'_{29})^{(5)} + a''_{29} \right)^{(5)} (T_{29}(s_{(28)}), s_{(28)}) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[ (a_{30})^{(5)} G_{29}(s_{(28)}) - \left( (a'_{30})^{(5)} + a''_{30} \right)^{(5)} (T_{29}(s_{(28)}), s_{(28)}) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[ (b_{28})^{(5)} T_{29}(s_{(28)}) - \left( (b'_{28})^{(5)} - b''_{28} \right)^{(5)} (G_{31}(s_{(28)}), s_{(28)}) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[ (b_{29})^{(5)} T_{28}(s_{(28)}) - \left( (b'_{29})^{(5)} - b''_{29} \right)^{(5)} (G_{31}(s_{(28)}), s_{(28)}) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[ (b_{30})^{(5)} T_{29}(s_{(28)}) - \left( (b'_{30})^{(5)} - b''_{30} \right)^{(5)} (G_{31}(s_{(28)}), s_{(28)}) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where  $s_{(28)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(6)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[ (a_{32})^{(6)} G_{33}(s_{(32)}) - \left( (a'_{32})^{(6)} + a''_{32} \right)^{(6)} (T_{33}(s_{(32)}), s_{(32)}) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[ (a_{33})^{(6)} G_{32}(s_{(32)}) - \left( (a'_{33})^{(6)} + a''_{33} \right)^{(6)} (T_{33}(s_{(32)}), s_{(32)}) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[ (a_{34})^{(6)} G_{33}(s_{(32)}) - \left( (a'_{34})^{(6)} + a''_{34} \right)^{(6)} (T_{33}(s_{(32)}), s_{(32)}) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[ (b_{32})^{(6)} T_{33}(s_{(32)}) - \left( (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[ (b_{33})^{(6)} T_{32}(s_{(32)}) - \left( (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[ (b_{34})^{(6)} T_{33}(s_{(32)}) - \left( (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where  $s_{(32)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(7)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[ (a_{36})^{(7)} G_{37}(s_{(36)}) - \left( (a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[ (a_{37})^{(7)} G_{36}(s_{(36)}) - \left( (a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[ (a_{38})^{(7)} G_{37}(s_{(36)}) - \left( (a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[ (b_{36})^{(7)} T_{37}(s_{(36)}) - \left( (b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[ (b_{37})^{(7)} T_{36}(s_{(36)}) - \left( (b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[ (b_{38})^{(7)} T_{37}(s_{(36)}) - \left( (b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where  $s_{(36)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(8)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[ (a_{40})^{(8)} G_{41}(s_{(40)}) - \left( (a'_{40})^{(8)} + a''_{40} \right)^{(8)} (T_{41}(s_{(40)}), s_{(40)}) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[ (a_{41})^{(8)} G_{40}(s_{(40)}) - \left( (a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[ (a_{42})^{(8)} G_{41}(s_{(40)}) - \left( (a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[ (b_{40})^{(8)} T_{41}(s_{(40)}) - \left( (b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[ (b_{41})^{(8)} T_{40}(s_{(40)}) - \left( (b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[ (b_{42})^{(8)} T_{41}(s_{(40)}) - \left( (b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where  $s_{(40)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

166

Consider operator  $\mathcal{A}^{(9)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[ (a_{44})^{(9)} G_{45}(s_{(44)}) - \left( (a'_{44})^{(9)} + a''_{44} \right)^{(9)} (T_{45}(s_{(44)}), s_{(44)}) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[ (a_{45})^{(9)} G_{44}(s_{(44)}) - \left( (a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[ (a_{46})^{(9)} G_{45}(s_{(44)}) - \left( (a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[ (b_{44})^{(9)} T_{45}(s_{(44)}) - \left( (b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[ (b_{45})^{(9)} T_{44}(s_{(44)}) - \left( (b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[ (b_{46})^{(9)} T_{45}(s_{(44)}) - \left( (b'_{46})^{(9)} - (b''_{46})^{(9)} (G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where  $s_{(44)}$  is the integrand that is integrated over an interval  $(0, t)$

The operator  $\mathcal{A}^{(1)}$  maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[ (a_{13})^{(1)} \left( G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}} \right) \right] ds_{(13)} =$$

$$(1 + (a_{13})^{(1)} t) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left( e^{(\hat{M}_{13})^{(1)} t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[ \left( (\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{\hat{M}_{13}^{(1)}}} + (\hat{P}_{13})^{(1)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 1

Analogous inequalities hold also for  $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator  $\mathcal{A}^{(2)}$  maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[ (a_{16})^{(2)} \left( G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} \tag{169}$$

$$= (1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left( e^{(\hat{M}_{16})^{(2)} t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[ \left( (\hat{P}_{16})^{(2)} + G_{17}^0 \right) e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{\hat{M}_{16}^{(2)}}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for  $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator  $\mathcal{A}^{(3)}$  maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[ (a_{20})^{(3)} \left( G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)}t)G_{21}^0 + \frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}(e^{(\hat{M}_{20})^{(3)}t} - 1)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0)e^{-(\hat{M}_{20})^{(3)}t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[ ((\hat{P}_{20})^{(3)} + G_{21}^0)e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for  $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator  $\mathcal{A}^{(4)}$  maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t [(a_{24})^{(4)}(G_{25}^0 + (\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}})] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)}t)G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}(e^{(\hat{M}_{24})^{(4)}t} - 1)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(\hat{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[ ((\hat{P}_{24})^{(4)} + G_{25}^0)e^{-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}} + (\hat{P}_{24})^{(4)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 4

The operator  $\mathcal{A}^{(5)}$  maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t [(a_{28})^{(5)}(G_{29}^0 + (\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}})] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}(e^{(\hat{M}_{28})^{(5)}t} - 1)$$

From which it follows that 175

$$(G_{28}(t) - G_{28}^0)e^{-(\hat{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left[ ((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}} + (\hat{P}_{28})^{(5)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 5

The operator  $\mathcal{A}^{(6)}$  maps the space of functions satisfying Equations into itself. Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t [(a_{32})^{(6)}(G_{33}^0 + (\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}})] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}(e^{(\hat{M}_{32})^{(6)}t} - 1)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(\hat{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[ ((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}} + (\hat{P}_{32})^{(6)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 6

Analogous inequalities hold also for  $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(a) The operator  $\mathcal{A}^{(7)}$  maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t [(a_{36})^{(7)} (G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}s_{(36)}})] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}(e^{(\hat{M}_{36})^{(7)}t} - 1)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[ ((\hat{P}_{36})^{(7)} + G_{37}^0)e^{-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}} + (\hat{P}_{36})^{(7)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 7

The operator  $\mathcal{A}^{(8)}$  maps the space of functions satisfying Equations into itself .Indeed it is obvious that

180

$$G_{40}(t) \leq G_{40}^0 + \int_0^t [(a_{40})^{(8)} (G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}})] ds_{(40)} =$$

$$(1 + (a_{40})^{(8)}t)G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}(e^{(\hat{M}_{40})^{(8)}t} - 1)$$

From which it follows that

181

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[ ((\hat{P}_{40})^{(8)} + G_{41}^0)e^{-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}} + (\hat{P}_{40})^{(8)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 8

Analogous inequalities hold also for  $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator  $\mathcal{A}^{(9)}$  maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t [(a_{44})^{(9)} (G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}s_{(44)}})] ds_{(44)} =$$

$$(1 + (a_{44})^{(9)}t)G_{45}^0 + \frac{(a_{44})^{(9)}(\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}(e^{(\hat{M}_{44})^{(9)}t} - 1)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0)e^{-(\hat{M}_{44})^{(9)}t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[ ((\hat{P}_{44})^{(9)} + G_{45}^0)e^{-\left(\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 9

Analogous inequalities hold also for  $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take  $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$  and to choose 182

$(\hat{P}_{13})^{(1)}$  and  $(\hat{Q}_{13})^{(1)}$  large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[ (\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0)e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{13})^{(1)} \tag{183}$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[ ((\hat{Q}_{13})^{(1)} + T_j^0)e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)} \tag{184}$$

In order that the operator  $\mathcal{A}^{(1)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(1)}$  is a contraction with respect to the metric 185

$$d((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{13})^{(1)}t} \}$$

Indeed if we denote

**Definition of**  $\tilde{G}, \tilde{T} : (\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\begin{aligned}
 |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\
 &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{-(\bar{M}_{13})^{(1)}s_{(13)}} + \\
 (a''_{13})^{(1)}(T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}} + \\
 G_{13}^{(2)} |(a'_{13})^{(1)}(T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)}(T_{14}^{(2)}, s_{(13)})| e^{-(\bar{M}_{13})^{(1)}s_{(13)}} e^{(\bar{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)}
 \end{aligned}$$

Where  $s_{(13)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned}
 |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{13})^{(1)}t} & \leq \frac{1}{(\bar{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\bar{A}_{13})^{(1)}) \\
 & + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)} d((G^{(1)}, T^{(1)}); G^{(2)}, T^{(2)})
 \end{aligned}$$
186

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 1:** The fact that we supposed  $(a''_{13})^{(1)}$  and  $(b''_{13})^{(1)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\hat{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$  and  $(\hat{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(1)}$  and  $(b''_i)^{(1)}$ ,  $i = 13, 14, 15$  depend only on  $T_{14}$  and respectively on  $G$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 19 to 24 it results

$$\begin{aligned}
 G_i(t) &\geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(1)} - (a''_i)^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}]} \geq 0 \\
 T_i(t) &\geq T_i^0 e^{-(b'_i)^{(1)}t} > 0 \text{ for } t > 0
 \end{aligned}$$

**Definition of**  $((\bar{M}_{13})^{(1)})_1, ((\bar{M}_{13})^{(1)})_2$  and  $((\bar{M}_{13})^{(1)})_3$  : 187

**Remark 3:** if  $G_{13}$  is bounded, the same property have also  $G_{14}$  and  $G_{15}$ . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$  it follows  $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)}G_{14}$  and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If  $G_{14}$  or  $G_{15}$  is bounded, the same property follows for  $G_{13}$  ,  $G_{15}$  and  $G_{13}$  ,  $G_{14}$  respectively.

**Remark 4:** If  $G_{13}$  is bounded, from below, the same property holds for  $G_{14}$  and  $G_{15}$  . The proof is analogous with the preceding one. An analogous property is true if  $G_{14}$  is bounded from below. 188

**Remark 5:** If  $T_{13}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(1)}(G(t), t)) = (b'_{14})^{(1)}$  then  $T_{14} \rightarrow \infty$ . 189

**Definition of**  $(m)^{(1)}$  and  $\varepsilon_1$  :

Indeed let  $t_1$  be so that for  $t > t_1$

$$(b_{14})^{(1)} - (b''_i)^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then  $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$  which leads to

$$T_{14} \geq \left( \frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left( \frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for  $T_{15}$  if  $\lim_{t \rightarrow \infty} ((b''_{15})^{(1)}(G(t), t)) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take  $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$  and to choose 190

$(\widehat{P}_{16})^{(2)}$  and  $(\widehat{Q}_{16})^{(2)}$  large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} \left[ (\widehat{P}_{16})^{(2)} + ((\widehat{P}_{16})^{(2)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} \left[ ((\widehat{Q}_{16})^{(2)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{16})^{(2)} \right] \leq (\widehat{Q}_{16})^{(2)} \quad 192$$

In order that the operator  $\mathcal{A}^{(2)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself 193

The operator  $\mathcal{A}^{(2)}$  is a contraction with respect to the metric 194

$$d \left( ((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote 195

**Definition of**  $\widetilde{G}_{19}, \widetilde{T}_{19} : (\widetilde{G}_{19}, \widetilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\begin{aligned} |\widetilde{G}_{16}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)}(T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} | (a''_{16})^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)}(T_{17}^{(2)}, s_{(16)}) | e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} \} ds_{(16)} \end{aligned}$$

Where  $s_{(16)}$  represents integrand that is integrated over the interval  $[0, t]$  197

From the hypotheses it follows

$$\begin{aligned} |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} \\ \leq \frac{1}{(\bar{M}_{16})^{(2)}} ((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)}) \\ + (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)} d((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows 198

**Remark 6:** The fact that we supposed  $(a''_{16})^{(2)}$  and  $(b''_{16})^{(2)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$  and  $(\widehat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)}t}$  respectively of  $\mathbb{R}_+$ . 199

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(2)}$  and  $(b''_i)^{(2)}$ ,  $i = 16, 17, 18$  depend only on  $T_{17}$  and respectively on  $(G_{19})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 7:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  200

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{ (a'_i)^{(2)} - (a''_i)^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \} ds_{(16)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(2)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{16})^{(2)})_1, ((\widehat{M}_{16})^{(2)})_2$  and  $((\widehat{M}_{16})^{(2)})_3$  : 201

**Remark 8:** if  $G_{16}$  is bounded, the same property have also  $G_{17}$  and  $G_{18}$  . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$  it follows  $\frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a'_{17})^{(2)}G_{17}$  and by integrating

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a'_{17})^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a'_{18})^{(2)}$$

If  $G_{17}$  or  $G_{18}$  is bounded, the same property follows for  $G_{16}$  ,  $G_{18}$  and  $G_{16}$  ,  $G_{17}$  respectively.

**Remark 9:** If  $G_{16}$  is bounded, from below, the same property holds for  $G_{17}$  and  $G_{18}$  . The proof is analogous with the preceding one. An analogous property is true if  $G_{17}$  is bounded from below. 202

**Remark 10:** If  $T_{16}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(2)}((G_{19})(t), t)) = (b'_{17})^{(2)}$  then  $T_{17} \rightarrow \infty$ . 203

**Definition of**  $(m)^{(2)}$  and  $\varepsilon_2$  :

Indeed let  $t_2$  be so that for  $t > t_2$

$$(b_{17})^{(2)} - (b''_i)^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then  $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$  which leads to 204

$$T_{17} \geq \left( \frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$$

If we take  $t$  such that  $e^{-\varepsilon_2 t} = \frac{1}{2}$  it results

$$T_{17} \geq \left( \frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right), t = \log \frac{2}{\varepsilon_2}$$

By taking now  $\varepsilon_2$  sufficiently small one sees that  $T_{17}$  is unbounded. 205

The same property holds for  $T_{18}$  if  $\lim_{t \rightarrow \infty} (b''_{18})^{(2)}((G_{19})(t), t) = (b'_{18})^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take  $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$  and to choose 207

$(\widehat{P}_{20})^{(3)}$  and  $(\widehat{Q}_{20})^{(3)}$  large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} \left[ (\widehat{P}_{20})^{(3)} + ((\widehat{P}_{20})^{(3)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{20})^{(3)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{20})^{(3)}$$

208

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} \left[ ((\widehat{Q}_{20})^{(3)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{20})^{(3)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{20})^{(3)} \right] \leq (\widehat{Q}_{20})^{(3)}$$

209

In order that the operator  $\mathcal{A}^{(3)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself 210

The operator  $\mathcal{A}^{(3)}$  is a contraction with respect to the metric 211

$$d\left(\left((G_{23})^{(1)}, (T_{23})^{(1)}\right), \left((G_{23})^{(2)}, (T_{23})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \right\}$$

Indeed if we denote 212

**Definition of**  $\widetilde{G}_{23}, \widetilde{T}_{23} : ((\widetilde{G}_{23}), (\widetilde{T}_{23})) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$\begin{aligned} |\widetilde{G}_{20}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} | (a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)}) | e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned}$$

Where  $s_{(20)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \frac{1}{(\bar{M}_{20})^{(3)}} \left( (a_{20})^{(3)} + (a'_{20})^{(3)} + (\bar{A}_{20})^{(3)} \right) \\ &+ (\bar{P}_{20})^{(3)} (\bar{k}_{20})^{(3)} d\left(\left((G_{23})^{(1)}, (T_{23})^{(1)}\right); (G_{23})^{(2)}, (T_{23})^{(2)}\right) \end{aligned} \tag{214}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 11:** The fact that we supposed  $(a''_{20})^{(3)}$  and  $(b''_{20})^{(3)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\bar{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$  and  $(\bar{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$  respectively of  $\mathbb{R}_+$ . 215

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(3)}$  and  $(b''_i)^{(3)}$ ,  $i = 20, 21, 22$  depend only on  $T_{21}$  and respectively on  $(G_{23})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 12:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  216

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(3)} - (a''_i)^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \} ds_{(20)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{20})^{(3)})_1, ((\widehat{M}_{20})^{(3)})_2$  and  $((\widehat{M}_{20})^{(3)})_3$  : 217

**Remark 13:** if  $G_{20}$  is bounded, the same property have also  $G_{21}$  and  $G_{22}$  . indeed if

$G_{20} < ((\widehat{M}_{20})^{(3)})$  it follows  $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a_{21}')^{(3)}G_{21}$  and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a_{22}')^{(3)}$$

If  $G_{21}$  or  $G_{22}$  is bounded, the same property follows for  $G_{20}$  ,  $G_{22}$  and  $G_{20}$  ,  $G_{21}$  respectively.

**Remark 14:** If  $G_{20}$  is bounded, from below, the same property holds for  $G_{21}$  and  $G_{22}$  . The proof is analogous with the preceding one. An analogous property is true if  $G_{21}$  is bounded from below. 218

**Remark 15:** If  $T_{20}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$  then  $T_{21} \rightarrow \infty$ . 219

**Definition of**  $(m)^{(3)}$  and  $\varepsilon_3$  :

Indeed let  $t_3$  be so that for  $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then  $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$  which leads to 220

$$T_{21} \geq \left( \frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left( \frac{(a_{21})^{(3)}(m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for  $T_{22}$  if  $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take  $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$  and to choose 221

$(\widehat{P}_{24})^{(4)}$  and  $(\widehat{Q}_{24})^{(4)}$  large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \left[ (\widehat{P}_{24})^{(4)} + ((\widehat{P}_{24})^{(4)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \left[ ((\widehat{Q}_{24})^{(4)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{24})^{(4)} \right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator  $\mathcal{A}^{(4)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself 224

The operator  $\mathcal{A}^{(4)}$  is a contraction with respect to the metric 225

$$d\left(\left((G_{27})^{(1)}, (T_{27})^{(1)}\right), \left((G_{27})^{(2)}, (T_{27})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \right\}$$

Indeed if we denote

**Definition of**  $(\widetilde{G}_{27}), (\widetilde{T}_{27})$ :  $((\widetilde{G}_{27}), (\widetilde{T}_{27})) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\widetilde{G}_{24}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} + \\ &\int_0^t \{ (a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} + \\ &G_{24}^{(2)} | (a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)}) | e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} \} ds_{(24)} \end{aligned}$$

Where  $s_{(24)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} &\leq \frac{1}{(\bar{M}_{24})^{(4)}} \left( (a_{24})^{(4)} + (a'_{24})^{(4)} + (\bar{A}_{24})^{(4)} \right) \\ &+ (\bar{P}_{24})^{(4)} (\bar{k}_{24})^{(4)} d\left(\left((G_{27})^{(1)}, (T_{27})^{(1)}\right); \left((G_{27})^{(2)}, (T_{27})^{(2)}\right)\right) \end{aligned} \quad 226$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 16:** The fact that we supposed  $(a''_{24})^{(4)}$  and  $(b''_{24})^{(4)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\bar{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$  and  $(\bar{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$  respectively of  $\mathbb{R}_+$ . 227

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(4)}$  and  $(b''_i)^{(4)}$ ,  $i = 24, 25, 26$  depend only on  $T_{25}$  and respectively on  $(G_{27})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 17:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  228

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$  and  $((\widehat{M}_{24})^{(4)})_3$  : 229

**Remark 18:** if  $G_{24}$  is bounded, the same property have also  $G_{25}$  and  $G_{26}$ . indeed if

$G_{24} < ((\widehat{M}_{24})^{(4)})$  it follows  $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a_{25}')^{(4)}G_{25}$  and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25}')^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a_{25}')^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26}')^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a_{26}')^{(4)}$$

If  $G_{25}$  or  $G_{26}$  is bounded, the same property follows for  $G_{24}$ ,  $G_{26}$  and  $G_{24}$ ,  $G_{25}$  respectively.

**Remark 19:** If  $G_{24}$  is bounded, from below, the same property holds for  $G_{25}$  and  $G_{26}$ . The proof is 230  
 analogous with the preceding one. An analogous property is true if  $G_{25}$  is bounded from below.

**Remark 20:** If  $T_{24}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$  then  $T_{25} \rightarrow \infty$ . 231

**Definition of**  $(m)^{(4)}$  and  $\varepsilon_4$  :

Indeed let  $t_4$  be so that for  $t > t_4$

$$(b_{25}')^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then  $\frac{dT_{25}}{dt} \geq (a_{25}')^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$  which leads to 232

$$T_{25} \geq \left( \frac{(a_{25}')^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$$T_{25} \geq \left( \frac{(a_{25}')^{(4)}(m)^{(4)}}{2} \right), t = \log \frac{2}{\varepsilon_4} \text{ By taking now } \varepsilon_4 \text{ sufficiently small one sees that } T_{25} \text{ is unbounded.}$$

The same property holds for  $T_{26}$  if  $\lim_{t \rightarrow \infty} (b_{26}'')^{(4)}((G_{27})(t), t) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for  $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take  $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$  and to choose 233

$(\widehat{P}_{28})^{(5)}$  and  $(\widehat{Q}_{28})^{(5)}$  large to have

$$\frac{(a_i)^{(5)}}{(\bar{M}_{28})^{(5)}} \left[ (\hat{P}_{28})^{(5)} + ((\hat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\bar{M}_{28})^{(5)}} \left[ ((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator  $\mathcal{A}^{(5)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(5)}$  is a contraction with respect to the metric 236

$$d \left( ((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{28})^{(5)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{28})^{(5)}t} \right\}$$

Indeed if we denote

**Definition of**  $(\widetilde{G}_{31}), (\widetilde{T}_{31})$ :  $((\widetilde{G}_{31}), (\widetilde{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\tilde{G}_{28}^{(1)} - \tilde{G}_{28}^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}} ds_{(28)} + \\ &\int_0^t \{ (a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{-(\bar{M}_{28})^{(5)}s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}} + \\ &G_{28}^{(2)} | (a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)}) | e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}} \} ds_{(28)} \end{aligned}$$

Where  $s_{(28)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\bar{M}_{28})^{(5)}t} &\tag{237} \\ &\leq \frac{1}{(\bar{M}_{28})^{(5)}} \left( (a_{28})^{(5)} + (a'_{28})^{(5)} + (\hat{A}_{28})^{(5)} \right) \\ &+ (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)} d \left( ((G_{31})^{(1)}, (T_{31})^{(1)}); (G_{31})^{(2)}, (T_{31})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 21:** The fact that we supposed  $(a''_{28})^{(5)}$  and  $(b''_{28})^{(5)}$  depending also on can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 238

$(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$  and  $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)} t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a_i'')^{(5)}$  and  $(b_i'')^{(5)}$ ,  $i = 28, 29, 30$  depend only on  $T_{29}$  and respectively on  $(G_{31})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 22:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  239

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(5)} t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$  and  $((\widehat{M}_{28})^{(5)})_3$  : 240

**Remark 23:** if  $G_{28}$  is bounded, the same property have also  $G_{29}$  and  $G_{30}$ . indeed if

$G_{28} < ((\widehat{M}_{28})^{(5)})$  it follows  $\frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a_{29}')^{(5)} G_{29}$  and by integrating

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)} ((\widehat{M}_{28})^{(5)})_1 / (a_{29}')^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)} ((\widehat{M}_{28})^{(5)})_2 / (a_{30}')^{(5)}$$

If  $G_{29}$  or  $G_{30}$  is bounded, the same property follows for  $G_{28}$ ,  $G_{30}$  and  $G_{28}$ ,  $G_{29}$  respectively.

**Remark 24:** If  $G_{28}$  is bounded, from below, the same property holds for  $G_{29}$  and  $G_{30}$ . The proof is 241  
 analogous with the preceding one. An analogous property is true if  $G_{29}$  is bounded from below.

**Remark 25:** If  $T_{28}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(5)} ((G_{31})(t), t)) = (b_{29}')^{(5)}$  then  $T_{29} \rightarrow \infty$ . 242

**Definition of**  $(m)^{(5)}$  and  $\varepsilon_5$  :

Indeed let  $t_5$  be so that for  $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)} ((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then  $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)} (m)^{(5)} - \varepsilon_5 T_{29}$  which leads to 243

$$T_{29} \geq \left( \frac{(a_{29})^{(5)} (m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left( \frac{(a_{29})^{(5)} (m)^{(5)}}{2} \right), t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for  $T_{30}$  if  $\lim_{t \rightarrow \infty} ((b_{30}'')^{(5)} ((G_{31})(t), t)) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for  $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take  $\frac{(a_i)^{(6)}}{(\bar{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\bar{M}_{32})^{(6)}} < 1$  and to choose 244

$(\hat{P}_{32})^{(6)}$  and  $(\hat{Q}_{32})^{(6)}$  large to have

$$\frac{(a_i)^{(6)}}{(\bar{M}_{32})^{(6)}} \left[ (\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\bar{M}_{32})^{(6)}} \left[ ((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator  $\mathcal{A}^{(6)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(6)}$  is a contraction with respect to the metric 247

$$d \left( ((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{32})^{(6)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{32})^{(6)}t} \}$$

Indeed if we denote

**Definition of**  $(\widetilde{G_{35}}, \widetilde{T_{35}})$  :  $(\widetilde{G_{35}}, \widetilde{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_{32}^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}} ds_{(32)} + \\ &\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{-(\bar{M}_{32})^{(6)}s_{(32)}} + \\ & (a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}} + \\ & G_{32}^{(2)} | (a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)}) | e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}} \} ds_{(32)} \end{aligned}$$

Where  $s_{(32)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned}
 & |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)}t} & 248 \\
 & \leq \frac{1}{(\widehat{M}_{32})^{(6)}} \left( (a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} \right) \\
 & + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)} d \left( ((G_{35})^{(1)}, (T_{35})^{(1)}); (G_{35})^{(2)}, (T_{35})^{(2)} \right)
 \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 26:** The fact that we supposed  $(a''_{32})^{(6)}$  and  $(b''_{32})^{(6)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$  and  $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$  respectively of  $\mathbb{R}_+$ . 249

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(6)}$  and  $(b''_i)^{(6)}$ ,  $i = 32, 33, 34$  depend only on  $T_{33}$  and respectively on  $(G_{35})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 27:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  250

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(6)} - (a''_i)^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)}t} > 0 \text{ for } t > 0$$

**Definition of**  $(\widehat{M}_{32})^{(6)}_1, (\widehat{M}_{32})^{(6)}_2$  and  $(\widehat{M}_{32})^{(6)}_3$  : 251

**Remark 28:** if  $G_{32}$  is bounded, the same property have also  $G_{33}$  and  $G_{34}$ . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$  it follows  $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)}_1 - (a'_{33})^{(6)})G_{33}$  and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)}_2) = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)}_1) / (a'_{33})^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)}_3) = G_{34}^0 + 2(a_{34})^{(6)} ((\widehat{M}_{32})^{(6)}_2) / (a'_{34})^{(6)}$$

If  $G_{33}$  or  $G_{34}$  is bounded, the same property follows for  $G_{32}$ ,  $G_{34}$  and  $G_{32}$ ,  $G_{33}$  respectively.

**Remark 29:** If  $G_{32}$  is bounded, from below, the same property holds for  $G_{33}$  and  $G_{34}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{33}$  is bounded from below. 252

**Remark 30:** If  $T_{32}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$  then  $T_{33} \rightarrow \infty$ . 253

**Definition of**  $(m)^{(6)}$  and  $\varepsilon_6$  :

Indeed let  $t_6$  be so that for  $t > t_6$

$$(b_{33})^{(6)} - (b''_i)^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then  $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$  which leads to 254

$$T_{33} \geq \left( \frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$$

If we take  $t$  such that  $e^{-\varepsilon_6 t} = \frac{1}{2}$  it results

$$T_{33} \geq \left( \frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6}$$

By taking now  $\varepsilon_6$  sufficiently small one sees that  $T_{33}$  is unbounded.

The same property holds for  $T_{34}$  if  $\lim_{t \rightarrow \infty} (b_{34})''^{(6)}((G_{35})(t), t(t), t) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for  $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$  255

It is now sufficient to take  $\frac{(a_i)^{(7)}}{(\bar{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\bar{M}_{36})^{(7)}} < 1$  and to choose  $(\hat{P}_{36})^{(7)}$  and  $(\hat{Q}_{36})^{(7)}$  large to have

$$\frac{(a_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[ (\hat{P}_{36})^{(7)} + ((\hat{P}_{36})^{(7)} + G_j^0) e^{-\left( \frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{36})^{(7)}$$
256

$$\frac{(b_i)^{(7)}}{(\bar{M}_{36})^{(7)}} \left[ ((\hat{Q}_{36})^{(7)} + T_j^0) e^{-\left( \frac{(\hat{Q}_{36})^{(7)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{36})^{(7)} \right] \leq (\hat{Q}_{36})^{(7)}$$
257

In order that the operator  $\mathcal{A}^{(7)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(7)}$  is a contraction with respect to the metric 258

$$d \left( ((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

**Definition of**  $(\widetilde{G}_{39}), (\widetilde{T}_{39}) : ((\widetilde{G}_{39}), (\widetilde{T}_{39})) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned} |\tilde{G}_{36}^{(1)} - \tilde{G}_{36}^{(2)}| &\leq \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\ &\int_0^t \{ (a_{36}')^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{-(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &(a_{36}'')^{(7)} (T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} + \\ &G_{36}^{(2)} | (a_{36}'')^{(7)} (T_{37}^{(1)}, s_{(36)}) - (a_{36}'')^{(7)} (T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} \} ds_{(36)} \end{aligned}$$

Where  $s_{(36)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} & |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\widehat{M}_{36})^{(7)}t} \\ & \leq \frac{1}{(\widehat{M}_{36})^{(7)}} ((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)}) \\ & + (\widehat{P}_{36})^{(7)} (\widehat{k}_{36})^{(7)} d \left( ((G_{39})^{(1)}, (T_{39})^{(1)}); (G_{39})^{(2)}, (T_{39})^{(2)} \right) \end{aligned} \tag{259}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 31:** The fact that we supposed  $(a''_{36})^{(7)}$  and  $(b''_{36})^{(7)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)}t}$  and  $(\widehat{Q}_{36})^{(7)} e^{(\widehat{M}_{36})^{(7)}t}$  respectively of  $\mathbb{R}_+$ . 260

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(7)}$  and  $(b''_i)^{(7)}$ ,  $i = 36, 37, 38$  depend only on  $T_{37}$  and respectively on  $(G_{39})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 32:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  261

it results

$$G_i(t) \geq G_i^0 e^{\left[ - \int_0^t \{ (a'_i)^{(7)} - (a''_i)^{(7)} \} (T_{37}(s_{(36)}), s_{(36)}) ds_{(36)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{36})^{(7)})_1, ((\widehat{M}_{36})^{(7)})_2$  and  $((\widehat{M}_{36})^{(7)})_3$  : 262

**Remark 33:** if  $G_{36}$  is bounded, the same property have also  $G_{37}$  and  $G_{38}$ . indeed if

$G_{36} < ((\widehat{M}_{36})^{(7)})$  it follows  $\frac{dG_{37}}{dt} \leq ((\widehat{M}_{36})^{(7)})_1 - (a'_{37})^{(7)}G_{37}$  and by integrating

$$G_{37} \leq ((\widehat{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)} ((\widehat{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way, one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)} ((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If  $G_{37}$  or  $G_{38}$  is bounded, the same property follows for  $G_{36}$ ,  $G_{38}$  and  $G_{36}$ ,  $G_{37}$  respectively.

**Remark 34:** If  $G_{36}$  is bounded, from below, the same property holds for  $G_{37}$  and  $G_{38}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{37}$  is bounded from below. 263

**Remark 35:** If  $T_{36}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$  then  $T_{37} \rightarrow \infty$ . 264

**Definition of**  $(m)^{(7)}$  and  $\varepsilon_7$  :

Indeed let  $t_7$  be so that for  $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then  $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$  which leads to 265

$$T_{37} \geq \left( \frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$T_{37} \geq \left( \frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right)$ ,  $t = \log \frac{2}{\varepsilon_7}$  By taking now  $\varepsilon_7$  sufficiently small one sees that  $T_{37}$  is unbounded.

The same property holds for  $T_{38}$  if  $\lim_{t \rightarrow \infty} (b_{38}'')^{(7)}((G_{39})(t), t) = (b_{38}')^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take  $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$ ,  $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$  and to choose 266

$(\hat{P}_{40})^{(8)}$  and  $(\hat{Q}_{40})^{(8)}$  large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}} \left[ (\hat{P}_{40})^{(8)} + ((\hat{P}_{40})^{(8)} + G_j^0) e^{-\left( \frac{(\hat{P}_{40})^{(8)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} \left[ ((\hat{Q}_{40})^{(8)} + T_j^0) e^{-\left( \frac{(\hat{Q}_{40})^{(8)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{40})^{(8)} \right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator  $\mathcal{A}^{(8)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(8)}$  is a contraction with respect to the metric

$$d \left( ((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\hat{M}_{40})^{(8)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\hat{M}_{40})^{(8)} t} \right\} \quad 269$$

Indeed if we denote 270

**Definition of**  $(\widetilde{G}_{43}), (\widetilde{T}_{43})$  :  $((\widetilde{G}_{43}), (\widetilde{T}_{43})) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned}
 |\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} e^{(\widehat{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\
 &\int_0^t \{(a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} + \\
 (a''_{40})^{(8)}(T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} e^{(\widehat{M}_{40})^{(8)}s_{(40)}} + \\
 G_{40}^{(2)} |(a''_{40})^{(8)}(T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)}(T_{41}^{(2)}, s_{(40)})| e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\} ds_{(40)}
 \end{aligned}$$

Where  $s_{(40)}$  represents integrand that is integrated over the interval  $[0, t]$  272

From the hypotheses it follows

$$\begin{aligned}
 |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\widehat{M}_{40})^{(8)}t} & \tag{273} \\
 \leq \frac{1}{(\widehat{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} & \\
 + (\widehat{P}_{40})^{(8)} (\widehat{k}_{40})^{(8)}) d((G_{43})^{(1)}, (T_{43})^{(1)}; (G_{43})^{(2)}, (T_{43})^{(2)}) &
 \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 36:** The fact that we supposed  $(a''_{40})^{(8)}$  and  $(b''_{40})^{(8)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)}t}$  and  $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)}t}$  respectively of  $\mathbb{R}_+$ . 274

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(8)}$  and  $(b''_i)^{(8)}$ ,  $i = 40, 41, 42$  depend only on  $T_{41}$  and respectively on  $(G_{43})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 37** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  275

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{40})^{(8)})_1, ((\widehat{M}_{40})^{(8)})_2$  and  $((\widehat{M}_{40})^{(8)})_3$  : 276

**Remark 38:** if  $G_{40}$  is bounded, the same property have also  $G_{41}$  and  $G_{42}$ . indeed if

$G_{40} < ((\widehat{M}_{40})^{(8)})$  it follows  $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$  and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If  $G_{41}$  or  $G_{42}$  is bounded, the same property follows for  $G_{40}$  ,  $G_{42}$  and  $G_{40}$  ,  $G_{41}$  respectively.

**Remark 39:** If  $G_{40}$  is bounded, from below, the same property holds for  $G_{41}$  and  $G_{42}$  . The proof is analogous with the preceding one. An analogous property is true if  $G_{41}$  is bounded from below. 277

**Remark 40:** If  $T_{40}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$  then  $T_{41} \rightarrow \infty$ . 278

**Definition of**  $(m)^{(8)}$  and  $\varepsilon_8$  :

Indeed let  $t_8$  be so that for  $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then  $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$  which leads to 279

$$T_{41} \geq \left( \frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$$

If we take  $t$  such that  $e^{-\varepsilon_8 t} = \frac{1}{2}$  it results

$$T_{41} \geq \left( \frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8}$$

By taking now  $\varepsilon_8$  sufficiently small one sees that  $T_{41}$  is unbounded.

The same property holds for  $T_{42}$  if  $\lim_{t \rightarrow \infty} ((b_{42}'')^{(8)}((G_{43})(t), t(t), t)) = (b'_{42})^{(8)}$

It is now sufficient to take  $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$  and to choose  $(\widehat{P}_{44})^{(9)}$  and  $(\widehat{Q}_{44})^{(9)}$  large to have 279

A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[ (\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{44})^{(9)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[ ((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{44})^{(9)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator  $\mathcal{A}^{(9)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying 39,35,36 into itself

The operator  $\mathcal{A}^{(9)}$  is a contraction with respect to the metric

$$d \left( ((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(M_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(M_{44})^{(9)}t} \}$$

Indeed if we denote

**Definition of**  $(\widehat{G}_{47}), (\widehat{T}_{47}) : ((\widehat{G}_{47}), (\widehat{T}_{47})) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} e^{(\widehat{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \int_0^t \{(a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} + (a''_{44})^{(9)}(T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} e^{(\widehat{M}_{44})^{(9)}s_{(44)}} + G_{44}^{(2)} |(a''_{44})^{(9)}(T_{45}^{(1)}, s_{(44)}) - (a''_{44})^{(9)}(T_{45}^{(2)}, s_{(44)})| e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\} ds_{(44)}$$

Where  $s_{(44)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$|(G_{47})^{(1)} - G^{(2)}| e^{-(\widehat{M}_{44})^{(9)}t} \leq \frac{1}{(\widehat{M}_{44})^{(9)}} ((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)}) + (\widehat{P}_{44})^{(9)} (\widehat{k}_{44})^{(9)} d((G_{47})^{(1)}, (T_{47})^{(1)}; (G_{47})^{(2)}, (T_{47})^{(2)})$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis (39,35,36) the result follows

**Remark 41:** The fact that we supposed  $(a''_{44})^{(9)}$  and  $(b''_{44})^{(9)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{44})^{(9)} e^{(\widehat{M}_{44})^{(9)}t}$  and  $(\widehat{Q}_{44})^{(9)} e^{(\widehat{M}_{44})^{(9)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(9)}$  and  $(b''_i)^{(9)}$ ,  $i = 44, 45, 46$  depend only on  $T_{45}$  and respectively on  $(G_{47})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 42:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(9)} - (a''_i)^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{44})^{(9)})_1, ((\widehat{M}_{44})^{(9)})_2$  and  $((\widehat{M}_{44})^{(9)})_3$  :

**Remark 43:** if  $G_{44}$  is bounded, the same property have also  $G_{45}$  and  $G_{46}$ . indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$  it follows  $\frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)}G_{45}$  and by integrating

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way, one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If  $G_{45}$  or  $G_{46}$  is bounded, the same property follows for  $G_{44}$ ,  $G_{46}$  and  $G_{44}$ ,  $G_{45}$  respectively.

**Remark 44:** If  $G_{44}$  is bounded, from below, the same property holds for  $G_{45}$  and  $G_{46}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{45}$  is bounded from below.

**Remark 45:** If  $T_{44}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$  then  $T_{45} \rightarrow \infty$ .

**Definition of**  $(m)^{(9)}$  and  $\varepsilon_9$  :

Indeed let  $t_9$  be so that for  $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then  $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$  which leads to

$$T_{45} \geq \left( \frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_9 t} = \frac{1}{2} \text{ it results}$$

$$T_{45} \geq \left( \frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_9} \text{ By taking now } \varepsilon_9 \text{ sufficiently small one sees that } T_{45} \text{ is unbounded.}$$

The same property holds for  $T_{46}$  if  $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

### **Behavior of the solutions of equation**

280

**Theorem** If we denote and define

**Definition of**  $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$  :

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$  four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

**Definition of**  $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$  :

281

By  $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$  and respectively  $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$  the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

**Definition of**  $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$  :

282

By  $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$  and respectively  $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$  the roots of the equations

$$(a_{14})^{(1)}(\bar{v}^{(1)})^2 + (\sigma_2)^{(1)}\bar{v}^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(\bar{u}^{(1)})^2 + (\tau_2)^{(1)}\bar{u}^{(1)} - (b_{13})^{(1)} = 0$$

**Definition of**  $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$  :-

283

If we define  $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$  by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \quad \text{if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

and  $\boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \quad \text{if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously

284

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \quad \text{if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

and  $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)}$$

are defined

Then the solution of global equations satisfies the inequalities

285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where  $(p_i)^{(1)}$  is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\begin{aligned} & \left( \frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[ e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \right) \leq G_{15}(t) \\ & \leq \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[ e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t} \end{aligned} \quad 286$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)} ((R_1)^{(1)} - (b'_{15})^{(1)})} \left[ e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)} ((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[ e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

**Definition of**  $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$  :- 290

Where  $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

**Behavior of the solutions of equation** 291

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$  : 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$  four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)} \quad 293$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 294$$

**Definition of**  $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$  : 295

By  $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$  and respectively  $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$  the roots 296

of the equations  $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$  297

and  $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$  and 298

**Definition of**  $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$  : 299

By  $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$  and respectively  $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$  the 300

roots of the equations  $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$  301

and  $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$  302

**Definition of**  $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$  :- 303

If we define  $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$  by 304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \quad \text{if } (v_0)^{(2)} < (v_1)^{(2)} \quad 305$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \quad \text{if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 306$$

and  $(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \quad \text{if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 307$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \quad \mathbf{if} (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \mathbf{if} (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and  $(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \quad \mathbf{if} (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 309$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{(S_1)^{(2)} - (p_{16})^{(2)} t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$  is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} - (p_{16})^{(2)} t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left( \frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \right) [e^{((S_1)^{(2)} - (p_{16})^{(2)}) t} - e^{-(S_2)^{(2)} t}] + G_{18}^0 e^{-(S_2)^{(2)} t} \leq G_{18}(t) \quad 312$$

$$\leq \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} [e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}] + G_{18}^0 e^{-(a'_{18})^{(2)} t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \leq T_{16}(t) \leq T_{16}^0 e^{(R_1)^{(2)} + (r_{16})^{(2)} t} \quad 313}$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} + (r_{16})^{(2)} t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} [e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \leq T_{18}(t) \leq \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} [e^{((R_1)^{(2)} + (r_{16})^{(2)}) t} - e^{-(R_2)^{(2)} t}] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

**Definition of**  $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$ :- 316

Where  $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$  317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)} (\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

**Behavior of the solutions** 319

**Theorem 3:** If we denote and define

**Definition of**  $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$  :

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$  four constants satisfying

$$\begin{aligned}
 -(\sigma_2)^{(3)} &\leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)} \\
 -(\tau_2)^{(3)} &\leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}
 \end{aligned}$$

**Definition of**  $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$  : 320

By  $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$  and respectively  $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$  the roots of the equations

$$\begin{aligned}
 (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} &= 0 \\
 \text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} &= 0 \text{ and}
 \end{aligned}$$

By  $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$  and respectively  $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$  the

roots of the equations  $(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and  $(b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

**Definition of**  $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$  :- 321

If we define  $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$  by

$$\begin{aligned}
 (m_2)^{(3)} &= (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \quad \text{if } (v_0)^{(3)} < (v_1)^{(3)} \\
 (m_2)^{(3)} &= (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \quad \text{if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},
 \end{aligned}$$

and 
$$(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \quad \text{if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously 322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \quad \text{if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \quad \text{if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \quad \text{and } (u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \quad \text{if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$  is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\left( \frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[ e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \leq G_{22}(t) \right. \tag{324}$$

$$\left. \leq \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})} \left[ e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \right.$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \tag{325}$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \tag{326}$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})} \left[ e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \tag{327}$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[ e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

**Definition of**  $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$ :- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

**Behavior of the solutions of equation**

**Theorem:** If we denote and define

**Definition of**  $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$  :

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$  four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

**Definition of**  $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$  : 329

By  $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$  and respectively  $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$  the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$  : 330

By  $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$  and respectively  $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$  the

$$\text{roots of the equations } (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

**Definition of**  $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$  :-

If we define  $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$  by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \quad \mathbf{if} (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \quad \mathbf{if} (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

and  $(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \quad \mathbf{if} (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \quad \mathbf{if} (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \quad \mathbf{if} (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

and  $(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \quad \mathbf{if} (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where  $(p_i)^{(4)}$  is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left( \frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[ e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \right) \leq G_{26}(t) \\ \leq \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[ e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[ e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[ e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

**Definition of**  $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$  :-

Where  $(S_1)^{(4)} = (a_{24})^{(4)} (m_2)^{(4)} - (a'_{24})^{(4)}$

331

332

333

334

335

336

337

$$\begin{aligned} (S_2)^{(4)} &= (a_{26})^{(4)} - (p_{26})^{(4)} \\ (R_1)^{(4)} &= (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)} \\ (R_2)^{(4)} &= (b'_{26})^{(4)} - (r_{26})^{(4)} \end{aligned}$$

**Behavior of the solutions of equation**

338

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$  :

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$  four constants satisfying

$$\begin{aligned} -(\sigma_2)^{(5)} &\leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)} \\ -(\tau_2)^{(5)} &\leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)} \end{aligned}$$

**Definition of**  $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$  :

339

By  $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$  and respectively  $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$  the roots of the equations  $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$  and  $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$  and

**Definition of**  $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$  :

340

By  $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$  and respectively  $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$  the roots of the equations  $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$  and  $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

**Definition of**  $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$  :-

If we define  $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$  by

$$\begin{aligned} (m_2)^{(5)} &= (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \quad \text{if } (v_0)^{(5)} < (v_1)^{(5)} \\ (m_2)^{(5)} &= (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \quad \text{if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)}, \\ \text{and } (v_0)^{(5)} &= \frac{G_{28}^0}{G_{29}^0} \end{aligned}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \quad \text{if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously

341

$$\begin{aligned} (\mu_2)^{(5)} &= (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \quad \text{if } (u_0)^{(5)} < (u_1)^{(5)} \\ (\mu_2)^{(5)} &= (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \quad \text{if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}, \\ \text{and } (u_0)^{(5)} &= \frac{T_{28}^0}{T_{29}^0} \end{aligned}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \quad \text{if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where  $(p_i)^{(5)}$  is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\begin{aligned} & \left( \frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \left[ e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \right. \\ & \left. \leq \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[ e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t} \right) \end{aligned} \quad 344$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[ e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[ e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

**Definition of**  $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$ :- 348

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

**Behavior of the solutions of equation** 349

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$  :

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$  four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

**Definition of**  $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$  : 350

By  $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$  and respectively  $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$  the roots of the equations

$$(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$$

$$\text{and } (b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$  : 351

By  $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$  and respectively  $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$  the roots of the equations  $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$  and  $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

**Definition of**  $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$  :-

If we define  $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$  by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \quad \text{if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$\text{and } (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \quad \text{if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously 352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \quad \text{if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \text{ if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where  $(p_i)^{(6)}$  is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left( \frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[ e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \right. \\ \left. \leq \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[ e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t} \right) \quad 355$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})} \left[ e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \tag{358}$$

$$\frac{(a_{34})^{(6)}T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[ e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

**Definition of**  $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$  :- 359

Where  $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

**Behavior of the solutions of equation**

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$  :

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$  four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

**Definition of**  $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$  : 361

By  $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$  and respectively  $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$  the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

and  $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$  and

**Definition of**  $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$  : 362

By  $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$  and respectively  $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$  the

roots of the equations  $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and  $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

**Definition of**  $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$  :-

If we define  $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$  by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \quad \text{if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \quad \text{if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

$$\text{and } \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \quad \text{if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \quad \text{if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \quad \text{if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

$$\text{and } \boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \quad \text{if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where  $(p_i)^{(7)}$  is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\begin{aligned} & \left( \frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) \left[ e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t} \right] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t) \quad 366 \\ & \leq \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} \left[ e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t} \right] + G_{38}^0 e^{-(a'_{38})^{(7)}t} \end{aligned}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)} ((R_1)^{(7)} - (b'_{38})^{(7)})} \left[ e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)} ((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[ e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

**Definition of**  $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$ :-

370

Where  $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

**Behavior of the solutions of equation**

371

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$  :

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$  four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

**Definition of**  $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$  :

372

By  $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$  and respectively  $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$  the roots of the equations

$$(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$  :

By  $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$  and respectively  $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$  the

$$\text{roots of the equations } (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(u^{(8)})^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$$

**Definition of**  $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$  :-

If we define  $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$  by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \quad \text{if } (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \quad \text{if } (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \quad \text{if } (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \quad \text{if } (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \quad \text{if } (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } (u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \quad \text{if } (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where  $(p_i)^{(8)}$  is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\begin{aligned} & \left( \frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \right) \left[ e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \leq G_{42}(t) \quad 377 \\ & \leq \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)} ((S_1)^{(8)} - (a'_{42})^{(8)})} \left[ e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t} \end{aligned}$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)} ((R_1)^{(8)} - (b'_{42})^{(8)})} \left[ e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)} ((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[ e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

**Definition of**  $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$ :- 381

Where  $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

**Behavior of the solutions of equation 37 to 92** 382

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$  :

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$  four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

**Definition of**  $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$  :

By  $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$  and respectively  $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$  the roots of the equations  $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$  and  $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$  and

**Definition of**  $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$  :

By  $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$  and respectively  $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$  the roots of the equations  $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$  and  $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

**Definition of**  $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$  :-

If we define  $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$  by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \quad \text{if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \quad \text{if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

and  $(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \quad \text{if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \quad \text{if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \quad \text{if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

and  $(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$ , **if**  $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$  where  $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$  are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((s_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(s_1)^{(9)}t}$$

where  $(p_i)^{(9)}$  is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[ e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a_{46})^{(9)})} \left[ e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[ e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[ e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

**Definition of**  $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$  :-

$$\text{Where } (S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

**Proof:** From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left( (a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

**Definition of**  $v^{(1)}$  :-  $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$- \left( (a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left( (a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$  :-

$$\text{For } 0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}(v_1)^{(1)} - (v_0)^{(1)}]t}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}(v_1)^{(1)} - (v_0)^{(1)}]t}}, \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

it follows  $(v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}}, \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}$$

From which we deduce  $(v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$

If  $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$  we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)} e^{[-(a_{14})^{(1)}(v_1)^{(1)} - (v_2)^{(1)}]t}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)}(v_1)^{(1)} - (v_2)^{(1)}]t}} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}} \leq (\bar{v}_1)^{(1)}$$

If  $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$ , we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)}(\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}]t}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of  $v^{(1)}(t)$  :-**

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

**Definition of  $u^{(1)}(t)$  :-**

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{13})^{(1)} = (a''_{14})^{(1)}$ , then  $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$  and in this case  $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$  if in addition  $(v_0)^{(1)} = (v_1)^{(1)}$  then  $v^{(1)}(t) = (v_0)^{(1)}$  and as a consequence  $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$  this also defines  $(v_0)^{(1)}$  for

the special case

Analogously if  $(b''_{13})^{(1)} = (b''_{14})^{(1)}$ , then  $(\tau_1)^{(1)} = (\tau_2)^{(1)}$  and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$  if in addition  $(u_0)^{(1)} = (u_1)^{(1)}$  then  $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$  This is an important consequence of the relation between  $(v_1)^{(1)}$  and  $(\bar{v}_1)^{(1)}$ , and definition of  $(u_0)^{(1)}$ .

**Proof:** From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left( (a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

**Definition of**  $v^{(2)}$  :- 388

$$v^{(2)} = \frac{G_{16}}{G_{17}}$$

It follows 389

$$-\left( (a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq -\left( (a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

**Definition of**  $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$  :-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

$$\text{it follows } (v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad (\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce  $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$  392

If  $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$  we find like in the previous case, 393

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}$$

If  $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$ , we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(2)}(t)$  :- 395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

**Definition of**  $u^{(2)}(t)$  :-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :** 397

If  $(a''_{16})^{(2)} = (a''_{17})^{(2)}$ , then  $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$  and in this case  $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$  if in addition  $(v_0)^{(2)} = (v_1)^{(2)}$  then  $v^{(2)}(t) = (v_0)^{(2)}$  and as a consequence  $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if  $(b''_{16})^{(2)} = (b''_{17})^{(2)}$ , then  $(\tau_1)^{(2)} = (\tau_2)^{(2)}$  and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$  if in addition  $(u_0)^{(2)} = (u_1)^{(2)}$  then  $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$  This is an important consequence of the relation between  $(v_1)^{(2)}$  and  $(\bar{v}_1)^{(2)}$

**Proof :** From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left( (a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

**Definition of**  $v^{(3)}$  :- 399

$$\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$$

It follows

$$-\left( (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq -\left( (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

400

From which one obtains

$$\text{For } 0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}} , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

it follows  $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get

401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

**Definition of  $(\bar{v}_1)^{(3)}$  :-**

From which we deduce  $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If  $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$  we find like in the previous case,

402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If  $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$  , we obtain

403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of  $v^{(3)}(t)$  :-**

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)} , \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

**Definition of  $u^{(3)}(t)$  :-**

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)} , \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{20})^{(3)} = (a''_{21})^{(3)}$  , then  $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$  and in this case  $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$  if in addition  $(v_0)^{(3)} =$

$(v_1)^{(3)}$  then  $v^{(3)}(t) = (v_0)^{(3)}$  and as a consequence  $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if  $(b_{20}'' )^{(3)} = (b_{21}'' )^{(3)}$ , then  $(\tau_1)^{(3)} = (\tau_2)^{(3)}$  and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$  if in addition  $(u_0)^{(3)} = (u_1)^{(3)}$  then  $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$  This is an important consequence of the relation between  $(v_1)^{(3)}$  and  $(\bar{v}_1)^{(3)}$

**Proof:** From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left( (a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

**Definition of**  $v^{(4)}$  :- 
$$v^{(4)} = \frac{G_{24}}{G_{25}}$$

It follows

$$- \left( (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left( (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$  :-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}$$

$$\text{it follows } (v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}$$

From which we deduce  $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If  $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$  we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}}{1 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If  $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$ , we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)} e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(4)}(t)$  :-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(4)}(t)$  :-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a_{24}'' )^{(4)} = (a_{25}'' )^{(4)}$ , then  $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$  and in this case  $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$  if in addition  $(v_0)^{(4)} = (v_1)^{(4)}$  then  $v^{(4)}(t) = (v_0)^{(4)}$  and as a consequence  $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$  **this also defines**  $(v_0)^{(4)}$  **for the special case .**

Analogously if  $(b_{24}'' )^{(4)} = (b_{25}'' )^{(4)}$ , then  $(\tau_1)^{(4)} = (\tau_2)^{(4)}$  and then  $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$  if in addition  $(u_0)^{(4)} = (u_1)^{(4)}$  then  $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$  This is an important consequence of the relation between  $(v_1)^{(4)}$  and  $(\bar{v}_1)^{(4)}$ , **and definition of**  $(u_0)^{(4)}$ .

408

**Proof:** From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left( (a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) \right) - (a''_{29})^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

**Definition of**  $v^{(5)}$  :-  $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left( (a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left( (a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$  :-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_0)^{(5)}]t}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}(v_1)^{(5)} - (v_0)^{(5)}]t}}, \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}$$

$$\text{it follows } (v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$$

In the same manner, we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} (\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}] t}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} (\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}] t}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}$$

From which we deduce  $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If  $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$  we find like in the previous case, 410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)} (v_2)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (v_2)^{(5)}) t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)} ((v_1)^{(5)} - (v_2)^{(5)}) t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} \leq (\bar{v}_1)^{(5)}$$

If  $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$ , we obtain 411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)} (\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)} ((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)} ((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)}) t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c), we have

**Definition of  $v^{(5)}(t)$  :-**

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

**Definition of  $u^{(5)}(t)$  :-**

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a_{28}'' )^{(5)} = (a_{29}'' )^{(5)}$ , then  $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$  and in this case  $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$  if in addition  $(v_0)^{(5)} = (v_5)^{(5)}$  then  $v^{(5)}(t) = (v_0)^{(5)}$  and as a consequence  $G_{28}(t) = (v_0)^{(5)} G_{29}(t)$  **this also defines  $(v_0)^{(5)}$  for the special case .**

Analogously if  $(b_{28}'' )^{(5)} = (b_{29}'' )^{(5)}$ , then  $(\tau_1)^{(5)} = (\tau_2)^{(5)}$  and then  $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$  if in addition  $(u_0)^{(5)} = (u_1)^{(5)}$  then  $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$  This is an important consequence of the relation between  $(v_1)^{(5)}$  and  $(\bar{v}_1)^{(5)}$ , **and definition of  $(u_0)^{(5)}$ .**

**Proof :** From global equations we obtain 412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left( (a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

**Definition of  $v^{(6)}$  :-**  $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)}\right) \leq \frac{dv^{(6)}}{dt} \leq -\left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$  :-

$$\text{For } 0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)}e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)}e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

$$\text{it follows } (v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$$

In the same manner , we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)}e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce  $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If  $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$  we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)}e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)}e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)}e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

If  $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$  , we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)}e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)}e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(6)}(t)$  :-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(6)}(t)$  :-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{32})^{(6)} = (a''_{33})^{(6)}$ , then  $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$  and in this case  $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$  if in addition  $(v_0)^{(6)} = (v_1)^{(6)}$  then  $v^{(6)}(t) = (v_0)^{(6)}$  and as a consequence  $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$  **this also defines  $(v_0)^{(6)}$  for the special case.**

Analogously if  $(b''_{32})^{(6)} = (b''_{33})^{(6)}$ , then  $(\tau_1)^{(6)} = (\tau_2)^{(6)}$  and then  $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$  if in addition  $(u_0)^{(6)} = (u_1)^{(6)}$  then  $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$  This is an important consequence of the relation between  $(v_1)^{(6)}$  and  $(\bar{v}_1)^{(6)}$ , **and definition of  $(u_0)^{(6)}$ .**

416

**Proof :** From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left( (a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

**Definition of  $v^{(7)}$  :-** 
$$v^{(7)} = \frac{G_{36}}{G_{37}}$$

It follows

$$- \left( (a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left( (a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

**Definition of  $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$  :-**

For  $0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_0)^{(7)})t]}} , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

it follows  $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce  $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If  $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$  we find like in the previous case,

418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}((v_1)^{(7)} - (v_2)^{(7)})t]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (\bar{v}_1)^{(7)}$$

If  $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$ , we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)})t]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(7)}(t)$  :-

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(7)}(t)$  :- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{36})^{(7)} = (a''_{37})^{(7)}$ , then  $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$  and in this case  $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$  if in addition  $(v_0)^{(7)} = (v_1)^{(7)}$  then  $v^{(7)}(t) = (v_0)^{(7)}$  and as a consequence  $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$  **this also defines  $(v_0)^{(7)}$  for the special case .**

Analogously if  $(b''_{36})^{(7)} = (b''_{37})^{(7)}$ , then  $(\tau_1)^{(7)} = (\tau_2)^{(7)}$  and then  $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$  if in addition  $(u_0)^{(7)} = (u_1)^{(7)}$  then  $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$  This is an important consequence of the relation between  $(v_1)^{(7)}$  and  $(\bar{v}_1)^{(7)}$ , **and definition of  $(u_0)^{(7)}$ .**

**Proof:** From global equations we obtain 421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left( (a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

**Definition of**  $v^{(8)}$  :-  $\boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left( (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq -\left( (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$  :-

$$\text{For } 0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_0)^{(8)})t]}} , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}$$

it follows  $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}$$

From which we deduce  $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If  $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$  we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}((v_1)^{(8)} - (v_2)^{(8)})t]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (\bar{v}_1)^{(8)}$$

If  $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$  , we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)})t]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(8)}(t)$  :-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)} , \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(8)}(t)$  :-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)} , \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{40})^{(8)} = (a''_{41})^{(8)}$ , then  $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$  and in this case  $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$  if in addition  $(v_0)^{(8)} = (v_1)^{(8)}$  then  $v^{(8)}(t) = (v_0)^{(8)}$  and as a consequence  $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$  **this also defines  $(v_0)^{(8)}$  for the special case .**

Analogously if  $(b''_{40})^{(8)} = (b''_{41})^{(8)}$ , then  $(\tau_1)^{(8)} = (\tau_2)^{(8)}$  and then  $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$  if in addition  $(u_0)^{(8)} = (u_1)^{(8)}$  then  $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$  This is an important consequence of the relation between  $(v_1)^{(8)}$  and  $(\bar{v}_1)^{(8)}$ , **and definition of  $(u_0)^{(8)}$ .**

**Proof :** From 99,20,44,22,23,44 we obtain

424  
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left( (a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

**Definition of  $v^{(9)}$  :-** 
$$v^{(9)} = \frac{G_{44}}{G_{45}}$$

It follows

$$- \left( (a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left( (a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

**Definition of  $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$  :-**

For  $0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)}e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)}e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}$$

it follows  $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_0)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)}e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)}e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}$$

From which we deduce  $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If  $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$  we find like in the previous case,

$$(v_1)^{(9)} \leq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)}e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_2)^{(9)})t]}}{1 + (C)^{(9)}e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_2)^{(9)})t]}} \leq v^{(9)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)}e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)}e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} \leq (\bar{v}_1)^{(9)}$$

If  $0 < (v_1)^{(9)} \leq (\bar{v}_1)^{(9)} \leq \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$ , we obtain

$$(v_1)^{(9)} \leq v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} \leq (v_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

**Definition of  $v^{(9)}(t)$  :-**

$$(m_2)^{(9)} \leq v^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{v^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

**Definition of  $u^{(9)}(t)$  :-**

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

**Particular case :**

If  $(a_{44})''^{(9)} = (a_{45})''^{(9)}$ , then  $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$  and in this case  $(v_1)^{(9)} = (\bar{v}_1)^{(9)}$  if in addition  $(v_0)^{(9)} = (v_1)^{(9)}$  then  $v^{(9)}(t) = (v_0)^{(9)}$  and as a consequence  $G_{44}(t) = (v_0)^{(9)}G_{45}(t)$  **this also defines  $(v_0)^{(9)}$  for the special case .**

Analogously if  $(b_{44})''^{(9)} = (b_{45})''^{(9)}$ , then  $(\tau_1)^{(9)} = (\tau_2)^{(9)}$  and then  $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$  if in addition  $(u_0)^{(9)} = (u_1)^{(9)}$  then  $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$  This is an important consequence of the relation between  $(v_1)^{(9)}$  and  $(\bar{v}_1)^{(9)}$ , **and definition of  $(u_0)^{(9)}$ .**

We can prove the following

425

**Theorem :** If  $(a_i'')^{(1)}$  and  $(b_i'')^{(1)}$  are independent on  $t$ , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with  $(p_{13})^{(1)}, (r_{14})^{(1)}$  as defined by equation are satisfied, then the system

**Theorem :** If  $(a_i'')^{(2)}$  and  $(b_i'')^{(2)}$  are independent on  $t$ , and the conditions with the notations

426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

427

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$$

428

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0,$$

429

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

430

with  $(p_{16})^{(2)}, (r_{17})^{(2)}$  as defined by equation are satisfied, then the system

**Theorem :** If  $(a_i'')^{(3)}$  and  $(b_i'')^{(3)}$  are independent on  $t$ , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with  $(p_{20})^{(3)}, (r_{21})^{(3)}$  as defined by equation are satisfied, then the system

We can prove the following 432

**Theorem :** If  $(a_i'')^{(4)}$  and  $(b_i'')^{(4)}$  are independent on  $t$ , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with  $(p_{24})^{(4)}, (r_{25})^{(4)}$  as defined by equation are satisfied, then the system

**Theorem :** If  $(a_i'')^{(5)}$  and  $(b_i'')^{(5)}$  are independent on  $t$ , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with  $(p_{28})^{(5)}, (r_{29})^{(5)}$  as defined by equation are satisfied, then the system

**Theorem** If  $(a_i'')^{(6)}$  and  $(b_i'')^{(6)}$  are independent on  $t$ , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with  $(p_{32})^{(6)}, (r_{33})^{(6)}$  as defined by equation are satisfied, then the system

**Theorem :** If  $(a_i'')^{(7)}$  and  $(b_i'')^{(7)}$  are independent on  $t$ , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with  $(p_{36})^{(7)}, (r_{37})^{(7)}$  as defined by equation are satisfied, then the system

**Theorem :** If  $(a_i'')^{(8)}$  and  $(b_i'')^{(8)}$  are independent on  $t$ , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with  $(p_{40})^{(8)}, (r_{41})^{(8)}$  as defined by equation are satisfied, then the system

**Theorem :** If  $(a_i'')^{(9)}$  and  $(b_i'')^{(9)}$  are independent on  $t$ , and the conditions (with the notations 436  
 45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with  $(p_{44})^{(9)}, (r_{45})^{(9)}$  as defined by equation 45 are satisfied, then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27})]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27})]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27})]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

**Proof:** 485

(a) Indeed the first two equations have a nontrivial solution  $G_{13}, G_{14}$  if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) \\ + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

**Proof:** 486

(a) Indeed the first two equations have a nontrivial solution  $G_{16}, G_{17}$  if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) \\ + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

**Proof:** 487

(a) Indeed the first two equations have a nontrivial solution  $G_{20}, G_{21}$  if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) \\ + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

**Proof:** 488

(a) Indeed the first two equations have a nontrivial solution  $G_{24}, G_{25}$  if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) \\ + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

**Proof:** 489

(a) Indeed the first two equations have a nontrivial solution  $G_{28}, G_{29}$  if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

**Proof:**

490

(a) Indeed the first two equations have a nontrivial solution  $G_{32}, G_{33}$  if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

**Proof:**

491

(a) Indeed the first two equations have a nontrivial solution  $G_{36}, G_{37}$  if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

**Proof:**

492

(a) Indeed the first two equations have a nontrivial solution  $G_{40}, G_{41}$  if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

**Proof:**

492  
A

(a) Indeed the first two equations have a nontrivial solution  $G_{44}, G_{45}$  if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

**Definition and uniqueness of  $\Gamma_{14}^*$  :-**

493

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(1)}(T_{14})$  being increasing, it follows that there exists a unique  $T_{14}^*$  for which  $f(T_{14}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

**Definition and uniqueness of  $\Gamma_{17}^*$  :-**

494

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(2)}(T_{17})$  being increasing, it follows that there exists a unique  $T_{17}^*$  for which  $f(T_{17}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 495$$

**Definition and uniqueness of  $\Gamma_{21}^*$  :-**

496

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(1)}(T_{21})$  being increasing, it follows that there exists a unique  $T_{21}^*$  for which  $f(T_{21}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

**Definition and uniqueness of  $\Gamma_{25}^*$**  :-

497

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(4)}(T_{25})$  being increasing, it follows that there exists a unique  $T_{25}^*$  for which  $f(T_{25}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

**Definition and uniqueness of  $\Gamma_{29}^*$**  :-

498

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(5)}(T_{29})$  being increasing, it follows that there exists a unique  $T_{29}^*$  for which  $f(T_{29}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

**Definition and uniqueness of  $\Gamma_{33}^*$**  :-

499

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(6)}(T_{33})$  being increasing, it follows that there exists a unique  $T_{33}^*$  for which  $f(T_{33}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

**Definition and uniqueness of  $\Gamma_{37}^*$**  :-

500

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(7)}(T_{37})$  being increasing, it follows that there exists a unique  $T_{37}^*$  for which  $f(T_{37}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

**Definition and uniqueness of  $\Gamma_{41}^*$**  :-

501

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(8)}(T_{41})$  being increasing, it follows that there exists a unique  $T_{41}^*$  for which  $f(T_{41}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^*)]}$$

**Definition and uniqueness of  $\Gamma_{45}^*$  :-**

501  
A

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(9)}(T_{45})$  being increasing, it follows that there exists a unique  $T_{45}^*$  for which  $f(T_{45}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a_{44}')^{(9)} + (a_{44}'')^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a_{46}')^{(9)} + (a_{46}'')^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions  $G_{13}, G_{14}$  if

502

$$\begin{aligned} \varphi(G) &= (b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - \\ &[(b_{13}')^{(1)}(b_{14}'')^{(1)}(G) + (b_{14}')^{(1)}(b_{13}'')^{(1)}(G)] + (b_{13}'')^{(1)}(G)(b_{14}'')^{(1)}(G) = 0 \end{aligned}$$

Where in  $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{14}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{14}^*$  such that  $\varphi(G^*) = 0$

By the same argument, the equations admit solutions  $G_{16}, G_{17}$  if

503

$$\begin{aligned} \varphi(G_{19}) &= (b_{16}')^{(2)}(b_{17}')^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - \\ &[(b_{16}')^{(2)}(b_{17}'')^{(2)}(G_{19}) + (b_{17}')^{(2)}(b_{16}'')^{(2)}(G_{19})] + (b_{16}'')^{(2)}(G_{19})(b_{17}'')^{(2)}(G_{19}) = 0 \end{aligned}$$

Where in  $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{17}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{17}^*$  such that  $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions  $G_{20}, G_{21}$  if

505

$$\begin{aligned} \varphi(G_{23}) &= (b_{20}')^{(3)}(b_{21}')^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - \\ &[(b_{20}')^{(3)}(b_{21}'')^{(3)}(G_{23}) + (b_{21}')^{(3)}(b_{20}'')^{(3)}(G_{23})] + (b_{20}'')^{(3)}(G_{23})(b_{21}'')^{(3)}(G_{23}) = 0 \end{aligned}$$

Where in  $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{21}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{21}^*$  such that  $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions  $G_{24}, G_{25}$  if

506

$$\begin{aligned} \varphi(G_{27}) &= (b_{24}')^{(4)}(b_{25}')^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - \\ &[(b_{24}')^{(4)}(b_{25}'')^{(4)}(G_{27}) + (b_{25}')^{(4)}(b_{24}'')^{(4)}(G_{27})] + (b_{24}'')^{(4)}(G_{27})(b_{25}'')^{(4)}(G_{27}) = 0 \end{aligned}$$

Where in  $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{25}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{25}^*$  such that  $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions  $G_{28}, G_{29}$  if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in  $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{29}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{29}^*$  such that  $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions  $G_{32}, G_{33}$  if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in  $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{33}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{33}^*$  such that  $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions  $G_{36}, G_{37}$  if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in  $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{37}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{37}^*$  such that  $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions  $G_{40}, G_{41}$  if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$$

$$[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in  $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{41}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{41}^*$  such that  $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions  $G_{44}, G_{45}$  if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in  $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{45}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{45}^*$  such that  $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

$G_{14}^*$  given by  $\varphi(G^*) = 0, T_{14}^*$  given by  $f(T_{14}^*) = 0$  and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

$$G_{17}^* \text{ given by } \varphi((G_{19})^*) = 0 , T_{17}^* \text{ given by } f(T_{17}^*) = 0 \text{ and} \tag{512}$$

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} , G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \tag{513}$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} , T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \tag{514}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

$$G_{21}^* \text{ given by } \varphi((G_{23})^*) = 0 , T_{21}^* \text{ given by } f(T_{21}^*) = 0 \text{ and}$$

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} , G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

$$G_{25}^* \text{ given by } \varphi(G_{27}) = 0 , T_{25}^* \text{ given by } f(T_{25}^*) = 0 \text{ and}$$

$$G_{24}^* = \frac{(a_{24})^{(4)}G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} , G_{26}^* = \frac{(a_{26})^{(4)}G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)}T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} , T_{26}^* = \frac{(b_{26})^{(4)}T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \tag{517}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

$$G_{29}^* \text{ given by } \varphi((G_{31})^*) = 0 , T_{29}^* \text{ given by } f(T_{29}^*) = 0 \text{ and}$$

$$G_{28}^* = \frac{(a_{28})^{(5)}G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} , G_{30}^* = \frac{(a_{30})^{(5)}G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)}T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} , T_{30}^* = \frac{(b_{30})^{(5)}T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \tag{519}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

$G_{33}^*$  given by  $\varphi((G_{35})^*) = 0$ ,  $T_{33}^*$  given by  $f(T_{33}^*) = 0$  and

$$G_{32}^* = \frac{(a_{32})^{(6)}G_{33}^*}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} , G_{34}^* = \frac{(a_{34})^{(6)}G_{33}^*}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)}T_{33}^*}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^*)]} , T_{34}^* = \frac{(b_{34})^{(6)}T_{33}^*}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^*)]} \quad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

$G_{37}^*$  given by  $\varphi((G_{39})^*) = 0$ ,  $T_{37}^*$  given by  $f(T_{37}^*) = 0$  and

$$G_{36}^* = \frac{(a_{36})^{(7)}G_{37}^*}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} , G_{38}^* = \frac{(a_{38})^{(7)}G_{37}^*}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)}T_{37}^*}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^*)]} , T_{38}^* = \frac{(b_{38})^{(7)}T_{37}^*}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^*)]} \quad 523$$

Finally we obtain the unique solution 523

$G_{41}^*$  given by  $\varphi((G_{43})^*) = 0$ ,  $T_{41}^*$  given by  $f(T_{41}^*) = 0$  and

$$G_{40}^* = \frac{(a_{40})^{(8)}G_{41}^*}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^*)]} , G_{42}^* = \frac{(a_{42})^{(8)}G_{41}^*}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)}T_{41}^*}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^*)]} , T_{42}^* = \frac{(b_{42})^{(8)}T_{41}^*}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^*)]} \quad 524$$

Finally we obtain the unique solution of 89 to 99 523  
A

$G_{45}^*$  given by  $\varphi((G_{47})^*) = 0$ ,  $T_{45}^*$  given by  $f(T_{45}^*) = 0$  and

$$G_{44}^* = \frac{(a_{44})^{(9)}G_{45}^*}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^*)]} , G_{46}^* = \frac{(a_{46})^{(9)}G_{45}^*}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)}T_{45}^*}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^*)]} , T_{46}^* = \frac{(b_{46})^{(9)}T_{45}^*}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^*)]} \quad 524$$

### ASYMPTOTIC STABILITY ANALYSIS 524

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(1)}$  and  $(b_i'')^{(1)}$  Belong to  $C^{(1)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

**Proof:**Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}} (T_{14}^*) = (q_{14})^{(1)} \quad , \quad \frac{\partial (b_i'')^{(1)}}{\partial G_j} (G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})G_{13} + (a_{13})^{(1)}G_{14} - (q_{13})^{(1)}G_{13}^*T_{14} \tag{525}$$

$$\frac{dG_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})G_{14} + (a_{14})^{(1)}G_{13} - (q_{14})^{(1)}G_{14}^*T_{14} \tag{526}$$

$$\frac{dG_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})G_{15} + (a_{15})^{(1)}G_{14} - (q_{15})^{(1)}G_{15}^*T_{14} \tag{527}$$

$$\frac{dT_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})T_{13} + (b_{13})^{(1)}T_{14} + \sum_{j=13}^{15} (s_{(13)(j)})T_{13}^*G_j \tag{528}$$

$$\frac{dT_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})T_{14} + (b_{14})^{(1)}T_{13} + \sum_{j=13}^{15} (s_{(14)(j)})T_{14}^*G_j \tag{529}$$

$$\frac{dT_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})T_{15} + (b_{15})^{(1)}T_{14} + \sum_{j=13}^{15} (s_{(15)(j)})T_{15}^*G_j \tag{530}$$

**ASYMPTOTIC STABILITY ANALYSIS** 531

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(2)}$  and  $(b_i'')^{(2)}$  belong to  $C^2(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

Proof: Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial (a_{17}'')^{(2)}}{\partial T_{17}} (T_{17}^*) = (q_{17})^{(2)} \quad , \quad \frac{\partial (b_i'')^{(2)}}{\partial G_j} ((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})G_{16} + (a_{16})^{(2)}G_{17} - (q_{16})^{(2)}G_{16}^*T_{17} \tag{534}$$

$$\frac{dG_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})G_{17} + (a_{17})^{(2)}G_{16} - (q_{17})^{(2)}G_{17}^*T_{17} \tag{535}$$

$$\frac{dG_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})G_{18} + (a_{18})^{(2)}G_{17} - (q_{18})^{(2)}G_{18}^*T_{17} \tag{536}$$

$$\frac{dT_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})T_{16} + (b_{16})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(16)(j)})T_{16}^*G_j \quad 537$$

$$\frac{dT_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})T_{17} + (b_{17})^{(2)}T_{16} + \sum_{j=16}^{18} (s_{(17)(j)})T_{17}^*G_j \quad 538$$

$$\frac{dT_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})T_{18} + (b_{18})^{(2)}T_{17} + \sum_{j=16}^{18} (s_{(18)(j)})T_{18}^*G_j \quad 539$$

**ASYMPTOTIC STABILITY ANALYSIS** 540

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a''_i)^{(3)}$  and  $(b''_i)^{(3)}$  belong to  $C^{(3)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a''_{21})^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)}, \quad \frac{\partial (b''_i)^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})G_{20} + (a_{20})^{(3)}G_{21} - (q_{20})^{(3)}G_{20}^*T_{21} \quad 541$$

$$\frac{dG_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})G_{21} + (a_{21})^{(3)}G_{20} - (q_{21})^{(3)}G_{21}^*T_{21} \quad 542$$

$$\frac{dG_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})G_{22} + (a_{22})^{(3)}G_{21} - (q_{22})^{(3)}G_{22}^*T_{21} \quad 543$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)})T_{20}^*G_j \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)})T_{21}^*G_j \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)})T_{22}^*G_j \quad 546$$

**ASYMPTOTIC STABILITY ANALYSIS** 547

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a''_i)^{(4)}$  and  $(b''_i)^{(4)}$  belong to  $C^{(4)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $\mathbb{G}_i, \mathbb{T}_i$  :- 548

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{25}''^{(4)})}{\partial T_{25}} (T_{25}^*) = (q_{25})^{(4)} \quad , \quad \frac{\partial (b_i''^{(4)})}{\partial G_j} ((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^*\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)})T_{24}^*\mathbb{G}_j \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)})T_{25}^*\mathbb{G}_j \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)})T_{26}^*\mathbb{G}_j \quad 554$$

**ASYMPTOTIC STABILITY ANALYSIS** 555

**Theorem 5:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(5)}$  and  $(b_i'')^{(5)}$  belong to  $C^{(5)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $\mathbb{G}_i, \mathbb{T}_i$  :- 556

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}''^{(5)})}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)} \quad , \quad \frac{\partial (b_i''^{(5)})}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{dG_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})G_{30} + (a_{30})^{(5)}G_{29} - (q_{30})^{(5)}G_{30}^* T_{29} \quad 559$$

$$\frac{dT_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})T_{28} + (b_{28})^{(5)}T_{29} + \sum_{j=28}^{30} (s_{(28)(j)})T_{28}^* G_j \quad 560$$

$$\frac{dT_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})T_{29} + (b_{29})^{(5)}T_{28} + \sum_{j=28}^{30} (s_{(29)(j)})T_{29}^* G_j \quad 561$$

$$\frac{dT_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})T_{30} + (b_{30})^{(5)}T_{29} + \sum_{j=28}^{30} (s_{(30)(j)})T_{30}^* G_j \quad 562$$

**ASYMPTOTIC STABILITY ANALYSIS** 563

**Theorem 6:** If the conditions of the previous theorem are satisfied and if the functions  $(a''_i)^{(6)}$  and  $(b''_i)^{(6)}$  belong to  $C^{(6)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :- 564

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a''_{33})^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)}, \quad \frac{\partial (b''_i)^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})G_{32} + (a_{32})^{(6)}G_{33} - (q_{32})^{(6)}G_{32}^* T_{33} \quad 565$$

$$\frac{dG_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})G_{33} + (a_{33})^{(6)}G_{32} - (q_{33})^{(6)}G_{33}^* T_{33} \quad 566$$

$$\frac{dG_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})G_{34} + (a_{34})^{(6)}G_{33} - (q_{34})^{(6)}G_{34}^* T_{33} \quad 567$$

$$\frac{dT_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})T_{32} + (b_{32})^{(6)}T_{33} + \sum_{j=32}^{34} (s_{(32)(j)})T_{32}^* G_j \quad 568$$

$$\frac{dT_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})T_{33} + (b_{33})^{(6)}T_{32} + \sum_{j=32}^{34} (s_{(33)(j)})T_{33}^* G_j \quad 569$$

$$\frac{dT_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})T_{34} + (b_{34})^{(6)}T_{33} + \sum_{j=32}^{34} (s_{(34)(j)})T_{34}^* G_j \quad 570$$

**ASYMPTOTIC STABILITY ANALYSIS** 571

**Theorem 7:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(7)}$  and  $(b_i'')^{(7)}$  belong to  $C^{(7)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :- 572

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{37}'')^{(7)}}{\partial T_{37}} (T_{37}^*) = (q_{37})^{(7)}, \quad \frac{\partial (b_i'')^{(7)}}{\partial G_j} ((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^* \mathbb{T}_{37} \tag{573}$$

$$\frac{d\mathbb{G}_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^* \mathbb{T}_{37} \tag{574}$$

$$\frac{d\mathbb{G}_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^* \mathbb{T}_{37} \tag{575}$$

$$\frac{d\mathbb{T}_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38} (s_{(36)(j)})T_{36}^* \mathbb{G}_j \tag{576}$$

$$\frac{d\mathbb{T}_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38} (s_{(37)(j)})T_{37}^* \mathbb{G}_j \tag{578}$$

$$\frac{d\mathbb{T}_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38} (s_{(38)(j)})T_{38}^* \mathbb{G}_j \tag{579}$$

Obviously, these values represent an equilibrium solution

**ASYMPTOTIC STABILITY ANALYSIS**

**Theorem 8:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(8)}$  and  $(b_i'')^{(8)}$  belong to  $C^{(8)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}} (T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial (b_i'')^{(8)}}{\partial G_j} ((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})G_{40} + (a_{40})^{(8)}G_{41} - (q_{40})^{(8)}G_{40}^* T_{41} \quad 581$$

$$\frac{dG_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})G_{41} + (a_{41})^{(8)}G_{40} - (q_{41})^{(8)}G_{41}^* T_{41} \quad 582$$

$$\frac{dG_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})G_{42} + (a_{42})^{(8)}G_{41} - (q_{42})^{(8)}G_{42}^* T_{41} \quad 583$$

$$\frac{dT_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})T_{40} + (b_{40})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(40)(j)})T_{40}^* G_j \quad 584$$

$$\frac{dT_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})T_{41} + (b_{41})^{(8)}T_{40} + \sum_{j=40}^{42} (s_{(41)(j)})T_{41}^* G_j \quad 585$$

$$\frac{dT_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})T_{42} + (b_{42})^{(8)}T_{41} + \sum_{j=40}^{42} (s_{(42)(j)})T_{42}^* G_j \quad 586$$

### ASYMPTOTIC STABILITY ANALYSIS

**Theorem 9:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(9)}$  and  $(b_i'')^{(9)}$  belong to  $C^{(9)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

**Proof:** Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^* T_{45} \quad 586$$

B

$$\frac{dG_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})G_{45} + (a_{45})^{(9)}G_{44} - (q_{45})^{(9)}G_{45}^* T_{45} \quad 586$$

C

$$\frac{dG_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})G_{46} + (a_{46})^{(9)}G_{45} - (q_{46})^{(9)}G_{46}^* T_{45} \quad 586$$

D

$$\frac{dT_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})T_{44} + (b_{44})^{(9)}T_{45} + \sum_{j=44}^{46} (s_{(44)(j)})T_{44}^* G_j \quad 586$$

E

$$\frac{dT_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})T_{45} + (b_{45})^{(9)}T_{44} + \sum_{j=44}^{46} (s_{(45)(j)}T_{45}^*G_j)$$

586  
F

$$\frac{dT_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})T_{46} + (b_{46})^{(9)}T_{45} + \sum_{j=44}^{46} (s_{(46)(j)}T_{46}^*G_j)$$

586  
G

**The characteristic equation of this system is**

587

$$\begin{aligned}
 & ((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\
 & \left[ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\
 & \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \right) \\
 & + \left( ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \right) \\
 & \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \right) \\
 & \left( ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)})(\lambda)^{(1)} \right) \\
 & \left( ((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)})(\lambda)^{(1)} \right) \\
 & + \left( ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)})(\lambda)^{(1)} \right)(q_{15})^{(1)}G_{15} \\
 & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^* \\
 & \left. \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}) \\
 & \left[ ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(q_{16})^{(2)}G_{16}^* \right] \\
 & \left( ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(17)}T_{17}^* + (b_{17})^{(2)}s_{(16),(17)}T_{17}^* \right) \\
 & + \left( ((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)})(q_{16})^{(2)}G_{16}^* + (a_{16})^{(2)}(q_{17})^{(2)}G_{17}^* \right) \\
 & \left( ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(16)}T_{17}^* + (b_{17})^{(2)}s_{(16),(16)}T_{16}^* \right) \\
 & \left( ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)})(\lambda)^{(2)} \right) \\
 & \left( ((\lambda)^{(2)})^2 + ((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)})(\lambda)^{(2)} \right) \\
 & + \left( ((\lambda)^{(2)})^2 + ((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)})(\lambda)^{(2)} \right)(q_{18})^{(2)}G_{18} \\
 & + ((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)})(a_{18})^{(2)}(q_{17})^{(2)}G_{17}^* + (a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}G_{16}^* \\
 & \left. \left( ((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)})s_{(17),(18)}T_{17}^* + (b_{17})^{(2)}s_{(16),(18)}T_{16}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)})\{((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)}) \\
 & \left[ ((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)})(q_{21})^{(3)}G_{21}^* + (a_{21})^{(3)}(q_{20})^{(3)}G_{20}^* \right] \\
 & \left( ((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)})s_{(21),(21)}T_{21}^* + (b_{21})^{(3)}s_{(20),(21)}T_{21}^* \right) \\
 & + \left( ((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)})(q_{20})^{(3)}G_{20}^* + (a_{20})^{(3)}(q_{21})^{(3)}G_{21}^* \right)
 \end{aligned}$$

$$\begin{aligned}
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)})\{((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & [((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)})(q_{25})^{(4)}G_{25}^* + (a_{25})^{(4)}(q_{24})^{(4)}G_{24}^*] \\
 & ((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)})s_{(25),(25)}T_{25}^* + (b_{25})^{(4)}s_{(24),(25)}T_{25}^* \\
 & + ((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)})(q_{24})^{(4)}G_{24}^* + (a_{24})^{(4)}(q_{25})^{(4)}G_{25}^* \\
 & ((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)})s_{(25),(24)}T_{25}^* + (b_{25})^{(4)}s_{(24),(24)}T_{24}^* \\
 & (((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)})(\lambda)^{(4)}) \\
 & (((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)})(\lambda)^{(4)}) \\
 & + (((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)})(\lambda)^{(4)})(q_{26})^{(4)}G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)})(a_{26})^{(4)}(q_{25})^{(4)}G_{25}^* + (a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G_{24}^* \\
 & ((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)})s_{(25),(26)}T_{25}^* + (b_{25})^{(4)}s_{(24),(26)}T_{24}^* \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)})\{((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & [((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)})(q_{29})^{(5)}G_{29}^* + (a_{29})^{(5)}(q_{28})^{(5)}G_{28}^*] \\
 & ((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})s_{(29),(29)}T_{29}^* + (b_{29})^{(5)}s_{(28),(29)}T_{29}^* \\
 & + ((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)})(q_{28})^{(5)}G_{28}^* + (a_{28})^{(5)}(q_{29})^{(5)}G_{29}^* \\
 & ((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})s_{(29),(28)}T_{29}^* + (b_{29})^{(5)}s_{(28),(28)}T_{28}^* \\
 & (((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)})(\lambda)^{(5)}) \\
 & (((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)})(\lambda)^{(5)}) \\
 & + (((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)})(\lambda)^{(5)})(q_{30})^{(5)}G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)})(a_{30})^{(5)}(q_{29})^{(5)}G_{29}^* + (a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G_{28}^* \\
 & ((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})s_{(29),(30)}T_{29}^* + (b_{29})^{(5)}s_{(28),(30)}T_{28}^* \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)})\{((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & [((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)})(q_{33})^{(6)}G_{33}^* + (a_{33})^{(6)}(q_{32})^{(6)}G_{32}^*] \\
 & ((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(33)}T_{33}^* + (b_{33})^{(6)}s_{(32),(33)}T_{33}^* \\
 & + ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)})(q_{32})^{(6)}G_{32}^* + (a_{32})^{(6)}(q_{33})^{(6)}G_{33}^* \\
 & ((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(32)}T_{33}^* + (b_{33})^{(6)}s_{(32),(32)}T_{32}^* \\
 & (((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)})(\lambda)^{(6)}) \\
 & (((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)})(\lambda)^{(6)}) \\
 & + (((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)})(\lambda)^{(6)})(q_{34})^{(6)}G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)})(a_{34})^{(6)}(q_{33})^{(6)}G_{33}^* + (a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G_{32}^* \\
 & ((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(34)}T_{33}^* + (b_{33})^{(6)}s_{(32),(34)}T_{32}^* \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)})\{((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & [((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)})(q_{37})^{(7)}G_{37}^* + (a_{37})^{(7)}(q_{36})^{(7)}G_{36}^*] \\
 & ((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(37)}T_{37}^* + (b_{37})^{(7)}s_{(36),(37)}T_{37}^* \\
 & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)})(q_{36})^{(7)}G_{36}^* + (a_{36})^{(7)}(q_{37})^{(7)}G_{37}^* \\
 & ((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(36)}T_{37}^* + (b_{37})^{(7)}s_{(36),(36)}T_{36}^* \\
 & (((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)})(\lambda)^{(7)}) \\
 & (((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)})(\lambda)^{(7)}) \\
 & + (((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)})(\lambda)^{(7)})(q_{38})^{(7)}G_{38} \\
 & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)})(a_{38})^{(7)}(q_{37})^{(7)}G_{37}^* + (a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G_{36}^* \\
 & ((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(38)}T_{37}^* + (b_{37})^{(7)}s_{(36),(38)}T_{36}^* \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)})\{((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\
 & [((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)})(q_{41})^{(8)}G_{41}^* + (a_{41})^{(8)}(q_{40})^{(8)}G_{40}^*] \\
 & ((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)})s_{(41),(41)}T_{41}^* + (b_{41})^{(8)}s_{(40),(41)}T_{41}^* \\
 & + ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)})(q_{40})^{(8)}G_{40}^* + (a_{40})^{(8)}(q_{41})^{(8)}G_{41}^* \\
 & ((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)})s_{(41),(40)}T_{41}^* + (b_{41})^{(8)}s_{(40),(40)}T_{40}^* \\
 & (((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)})(\lambda)^{(8)}) \\
 & (((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)})(\lambda)^{(8)}) \\
 & + (((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)})(\lambda)^{(8)})(q_{42})^{(8)}G_{42} \\
 & + ((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)})(a_{42})^{(8)}(q_{41})^{(8)}G_{41}^* + (a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G_{40}^* \\
 & ((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)})s_{(41),(42)}T_{41}^* + (b_{41})^{(8)}s_{(40),(42)}T_{40}^* \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)})\{((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 & [((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)})(q_{45})^{(9)}G_{45}^* + (a_{45})^{(9)}(q_{44})^{(9)}G_{44}^*] \\
 & ((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)})s_{(45),(45)}T_{45}^* + (b_{45})^{(9)}s_{(44),(45)}T_{45}^* \\
 & + ((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)})(q_{44})^{(9)}G_{44}^* + (a_{44})^{(9)}(q_{45})^{(9)}G_{45}^* \\
 & ((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)})s_{(45),(44)}T_{45}^* + (b_{45})^{(9)}s_{(44),(44)}T_{44}^* \\
 & (((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)})(\lambda)^{(9)}) \\
 & (((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)})(\lambda)^{(9)}) \\
 & + (((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)})(\lambda)^{(9)})(q_{46})^{(9)}G_{46} \\
 & + ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)})(a_{46})^{(9)}(q_{45})^{(9)}G_{45}^* + (a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G_{44}^* \\
 & ((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)})s_{(45),(46)}T_{45}^* + (b_{45})^{(9)}s_{(44),(46)}T_{44}^* \} = 0
 \end{aligned}$$

**And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.**

## SECTION TWO

### Mass Gap: Freiwirtschaft; Natural Order

#### INTRODUCTION—VARIABLES USED

##### Mass gap (Wikipedia)

- (1) In quantum field theory, the mass gap is (=) the difference in energy between the vacuum and the next lowest energy state.
- (2) The energy of the vacuum is zero by definition, and assuming that all energy states can be thought of as particles in plane-waves, the mass gap is (=) the mass of the lightest particle.
- (3) Since exact energy eigenstates are infinitely spread out and are therefore usually excluded from a formal mathematical description, a stronger definition is that the mass gap is (=) the greatest lower bound of the energy of any state which is orthogonal to the vacuum.

##### Mathematical definitions

- (4) For a given real field  $\phi(x)$ , we can say that the theory has a mass gap if the two-point function has the property

$$\langle \phi(0, t) \phi(0, 0) \rangle \sim \sum_n A_n \exp(-\Delta_n t)$$

with  $\Delta_0 > 0$  being the lowest energy value in the spectrum of the Hamiltonian and thus the mass gap. This quantity, easy to generalize to other fields, is what is generally measured in lattice computations. It was proved in this way that Yang-Mills theory develops a mass gap. The corresponding time-ordered value, the propagator, will have the property

$$\lim_{p \rightarrow 0} \Delta(p) = \text{constant}$$

with the constant being finite. A typical example is offered by a free massive particle and, in this case, the constant has the value  $1/m^2$ . In the same limit, the propagator for a massless particle is singular.

##### (5) Examples from classical theories

An example of mass gap arising for massless theories, already at the classical level, can be seen in spontaneous breaking of symmetry or Higgs mechanism. In the former case, one has to cope with the appearance of massless excitations, Goldstone bosons, which are removed in the latter case due to gauge freedom. Quantization preserves this property.

A quartic massless scalar field theory develops a mass gap already at classical level. Let us consider the

equation

$$\square\phi + \lambda\phi^3 = 0.$$

**This equation has the exact solution**

$$\phi(x) = \mu \left(\frac{2}{\lambda}\right)^{\frac{1}{4}} \operatorname{sn}(p \cdot x + \theta, -1)$$

-- where  $\mu$  and  $\theta$  are integration constants, and sn is a Jacobi elliptic function -- provided

$$p^2 = \mu^2 \sqrt{\frac{\lambda}{2}}.$$

At the classical level, a mass gap appears while, at quantum level, one has a tower of excitations and this property of the theory is preserved after quantization in the limit of momenta going to zero.

While lattice computations have suggested that Yang-Mills theory indeed has a mass gap and a tower of excitations, a theoretical proof is still missing. This is one of the Clay Institute Millennium problems and it remains an open problem. Such states for Yang-Mills theory should be physical states, named glueballs, and should be observable in the laboratory.

### (6) Källén-Lehmann representation

If Källén-Lehmann spectral representation holds, at this stage we exclude gauge theories, the spectral density function can take a very simple form with a discrete spectrum starting with a mass gap

$$\rho(\mu^2) = \sum_{n=1}^N Z_n \delta(\mu^2 - m_n^2) + \rho_c(\mu^2)$$

being  $\rho_c(\mu^2)$  the contribution from multi-particle part of the spectrum. In this case, the propagator will take the simple form

$$\Delta(p) = \sum_{n=1}^N \frac{Z_n}{p^2 - m_n^2 + i\epsilon} + \int_{4m_N^2}^{\infty} d\mu^2 \rho_c(\mu^2) \frac{1}{p^2 - \mu^2 + i\epsilon}$$

being  $4m_N^2$  approximatively the starting point of the multi-particle sector. Now, using the fact that

$$\int_0^{\infty} d\mu^2 \rho(\mu^2) = 1$$

we arrive at the following conclusion for the constants in the spectral density

$$1 = \sum_{n=1}^N Z_n + \int_0^{\infty} d\mu^2 \rho_c(\mu^2)$$

This could not be true in a gauge theory. Rather it must be proved that a Källén-Lehmann representation for the propagator holds also for this case. Absence of multi-particle contributions implies that the theory is trivial, as no bound states appear in the theory and so there is no interaction, even if the theory has a mass gap. In this case we have immediately the propagator just setting  $\rho_c(\mu^2) = 0$  in the formulas above.

### (7) Yang–Mills theory

Yang–Mills theory is a gauge theory **based on** the SU (N) group, or more generally any compact, semi-simple Lie group.

Yang–Mills theory seeks to describe the behavior of elementary particles **using these** non-Abelian Lie groups and is at the core of the unification of the Weak and Electromagnetic force (i.e.  $U(1) \times SU(2)$ ) as well as Quantum Chromodynamics, the theory of the Strong force (based on  $SU(3)$ ). Thus it forms the basis of our current understanding of particle physics, the Standard Model.

In a private correspondence, Wolfgang Pauli formulated in 1953 a six-dimensional theory of Einstein's field equations of general relativity, extending the five-dimensional theory of Kaluza, Klein, Fock and others **to (eb) a** higher dimensional internal space.

However, there is no evidence that Pauli developed the Lagrangian of a gauge field or the quantization of it. Because Pauli found that his theory "**leads to** some rather unphysical shadow particles", he refrained from publishing his results formally.

Recent research shows that an extended Kaluza–Klein theory is in general **not equivalent** to Yang–Mills theory, as the former contains additional terms.

## NOTATION

### Module One

quantum field theory states that the mass gap is (=) the difference in energy between the vacuum and the next lowest energy state

$G_{13}$  : Category one of difference in energy between the vacuum and the next lowest energy state

$G_{14}$  : Category two of difference in energy between the vacuum and the next lowest energy state

$G_{15}$  : Category three of difference in energy between the vacuum and the next lowest energy state

$T_{13}$  : Category one of quantum field theory states that the mass gap

$T_{14}$  : Category two of quantum field theory states that the mass gap

$T_{15}$  : Category three of quantum field theory states that the mass gap

### Module Two

The energy of the vacuum is zero by definition, and assuming that all energy states can be thought of as particles in plane-waves, the mass gap is (=) the mass of the lightest particle

$G_{16}$  : Category one of mass of the lightest particle

$G_{17}$  : Category two of mass of the lightest particle

$G_{18}$  : Category three of mass of the lightest particle

$T_{16}$ : Category one of energy of the vacuum is zero by definition, and assuming that all energy states can be thought of as particles in plane-waves, the mass gap

$T_{17}$ : Category two of energy of the vacuum is zero by definition, and assuming that all energy states can be thought of as particles in plane-waves, the mass gap

$T_{18}$ : Category three of energy of the vacuum is zero by definition, and assuming that all energy states can be thought of as particles in plane-waves, the mass gap

### Module three

Since exact energy eigenstates are infinitely spread out and are therefore usually excluded from a formal mathematical description, a stronger definition is that the mass gap is (=) the greatest lower bound of the energy of any state which is orthogonal to the vacuum

$G_{20}$ : Category one of greatest lower bound of the energy of any state which is orthogonal to the vacuum

$G_{21}$ : Category two of greatest lower bound of the energy of any state which is orthogonal to the vacuum

$G_{22}$ : Category three of greatest lower bound of the energy of any state which is orthogonal to the vacuum

$T_{20}$ : Category one of exact energy eigenstates are infinitely spread out and are therefore usually excluded from a formal mathematical description, a stronger definition is that the mass gap

$T_{21}$ : Category two of exact energy eigenstates are infinitely spread out and are therefore usually excluded from a formal mathematical description, a stronger definition is that the mass gap

$T_{22}$ : Category three of exact energy eigenstates are infinitely spread out and are therefore usually excluded from a formal mathematical description, a stronger definition is that the mass gap

### Module four

For a given real field  $\phi(x)$ , we can say that the theory has a mass gap if the two-point function has the property

$$\langle \phi(0, t) \phi(0, 0) \rangle \sim \sum_n A_n \exp(-\Delta_n t)$$

with  $\Delta_0 > 0$  being the lowest energy value in the spectrum of the Hamiltonian and thus the mass gap. This quantity, easy to generalize to other fields, is what is generally measured in lattice computations. It was proved in this way that Yang-Mills theory develops a mass gap. The corresponding time-ordered value, the propagator, will have the property

$$\lim_{p \rightarrow 0} \Delta(p) = \text{constant}$$

with the constant being finite. A typical example is offered by a free massive particle and, in this case, the constant has the value  $1/m^2$ . In the same limit, the propagator for a massless particle is singular

$G_{24}$ : Category one of LHS of the equation constitutive of two-point function has the property

$G_{25}$ : Category two of LHS of the equation constitutive of two-point function has the property

$G_{26}$ : Category three of LHS of the equation constitutive of two-point function has the property

$T_{24}$  : Category one of RHS of the equation constitutive of two-point function has the property

$T_{25}$  : Category two of RHS of the equation constitutive of two-point function has the property

$T_{26}$  : Category three of RHS of the equation constitutive of two-point function has the property

### Module five

#### Examples from classical theories

An example of mass gap arising for massless theories, already at the classical level, can be seen in spontaneous breaking of symmetry or Higgs mechanism. In the former case, one has to cope with the appearance of massless excitations, Goldstone bosons, which are removed in the latter case due to gauge freedom. Quantization preserves this property.

A quartic massless scalar field theory develops a mass gap already at classical level. Let us consider the equation

$$\square\phi + \lambda\phi^3 = 0.$$

**This equation has the exact solution**

$$\phi(x) = \mu \left(\frac{2}{\lambda}\right)^{\frac{1}{4}} \operatorname{sn}(p \cdot x + \theta, -1)$$

-- where  $\mu$  and  $\theta$  are integration constants, and sn is a Jacobi elliptic function -- provided

$$p^2 = \mu^2 \sqrt{\frac{\lambda}{2}}.$$

At the classical level, a mass gap appears while, at quantum level, one has a tower of excitations and this property of the theory is preserved after quantization in the limit of momenta going to zero.

While lattice computations have suggested that Yang-Mills theory indeed has a mass gap and a tower of excitations, a theoretical proof is still missing. This is one of the Clay Institute Millennium problems and it remains an open problem. Such states for Yang-Mills theory should be physical states, named glueballs, and should be observable in the laboratory.

$G_{28}$  : Category one of LHS of the equation paradigmatic and epitome of Examples from classical theories

$G_{29}$  : Category two of LHS of the equation paradigmatic and epitome of Examples from classical theories

$G_{30}$  : Category three of LHS of the equation paradigmatic and epitome of Examples from classical theories

$T_{28}$  : Category one of RHS of the equation paradigmatic and epitome of examples from classical theories

$T_{29}$  : Category two of RHS of the equation paradigmatic and epitome of examples from classical theories

$T_{30}$  : Category three of RHS of the equation paradigmatic and epitome of examples from classical theories

### Module six

#### Källén-Lehmann representation

If Källén-Lehmann spectral representation holds, at this stage we exclude gauge theories, the spectral density function can take a very simple form with a discrete spectrum starting with a mass gap

$$\rho(\mu^2) = \sum_{n=1}^N Z_n \delta(\mu^2 - m_n^2) + \rho_c(\mu^2)$$

being  $\rho_c(\mu^2)$  the contribution from multi-particle part of the spectrum. In this case, the propagator will take the simple form

$$\Delta(p) = \sum_{n=1}^N \frac{Z_n}{p^2 - m_n^2 + i\epsilon} + \int_{4m_N^2}^{\infty} d\mu^2 \rho_c(\mu^2) \frac{1}{p^2 - \mu^2 + i\epsilon}$$

being  $4m_N^2$  approximately the starting point of the multi-particle sector. Now, using the fact that

$$\int_0^{\infty} d\mu^2 \rho(\mu^2) = 1$$

we arrive at the following conclusion for the constants in the spectral density

$$1 = \sum_{n=1}^N Z_n + \int_0^{\infty} d\mu^2 \rho_c(\mu^2)$$

This could not be true in a gauge theory. Rather it must be proved that a Källén-Lehmann representation for the propagator holds also for this case. Absence of multi-particle contributions implies that the theory is trivial, as no bound states appear in the theory and so there is no interaction, even if the theory has a mass gap. In this case we have immediately the propagator just setting  $\rho_c(\mu^2) = 0$  in the formulas above

$G_{32}$  : Category one of LHS of equations under the head and appellation Källén-Lehmann representation

$G_{33}$  : Category two of LHS of equations under the head and appellation Källén-Lehmann representation

$G_{34}$  : Category three of LHS of equations under the head and appellation Källén-Lehmann representation

$T_{32}$  : Category one of RHS of equations under the head and appellation Källén-Lehmann representation

$T_{33}$  : Category two of RHS of equations under the head and appellation Källén-Lehmann representation

$T_{34}$  : Category three of RHS of equations under the head and appellation Källén-Lehmann representation

### Module seven

Yang–Mills theory is a gauge theory **based on** the SU (N) group, or more generally any compact, semi-simple Lie group

$G_{36}$  : Category one of SU (N) group, or more generally any compact, semi-simple Lie group

$G_{37}$  : Category two of SU (N) group, or more generally any compact, semi-simple Lie group

$G_{38}$ : Category three ofSU (N) group, or more generally any compact, semi-simple Lie group

$T_{36}$  : Category one ofYang–Mills theory is a gauge theory

$T_{37}$  : Category two ofYang–Mills theory is a gauge theory

$T_{38}$  : Category three ofYang–Mills theory is a gauge theory

### Module eight

Yang–Mills theory seeks to describe the behavior of elementary particles **using these** non-Abelian Lie groups and is at the core of the unification of the Weak and Electromagnetic force (i.e.  $U(1) \times SU(2)$ ) as well as Quantum Chromodynamics, the theory of the Strong force (based on  $SU(3)$ ). Thus it forms the basis of our current understanding of particle physics, the Standard Model.

$G_{40}$  : Category one ofnon-Abelian Lie groups and is at the core of the unification of the Weak and Electromagnetic force (i.e.  $U(1) \times SU(2)$ ) as well as Quantum Chromodynamics, the theory of the Strong force (based on  $SU(3)$ ). Thus it forms the basis of our current understanding of particle physics, the Standard Model.

$G_{41}$  : Category two ofnon-Abelian Lie groups and is at the core of the unification of the Weak and Electromagnetic force (i.e.  $U(1) \times SU(2)$ ) as well as Quantum Chromodynamics, the theory of the Strong force (based on  $SU(3)$ ). Thus it forms the basis of our current understanding of particle physics, the Standard Model.

$G_{42}$  : Category three ofnon-Abelian Lie groups and is at the core of the unification of the Weak and Electromagnetic force (i.e.  $U(1) \times SU(2)$ ) as well as Quantum Chromodynamics, the theory of the Strong force (based on  $SU(3)$ ). Thus it forms the basis of our current understanding of particle physics, the Standard Model.

$T_{40}$  : Category one ofYang–Mills theory seeks to describe the behavior of elementary particles

$T_{41}$  : Category two ofYang–Mills theory seeks to describe the behavior of elementary particles

$T_{42}$  : Category three ofYang–Mills theory seeks to describe the behavior of elementary particles

### Module Nine

Recent research shows that an extended Kaluza–Klein theory is in general **not equivalent** to Yang–Mills theory, as the former contains additional terms

$G_{44}$  : Category one ofKaluza–Klein theory is in general; Yang–Mills theory, as the former contains additional terms

$G_{45}$  : Category two ofKaluza–Klein theory is in general; Yang–Mills theory, as the former contains additional terms

$G_{46}$  : Category three ofKaluza–Klein theory is in general; Yang–Mills theory, as the former contains

additional terms

$T_{44}$  : Category one of Yang–Mills theory, as the former contains additional terms;Kaluza–Klein theory is in general

$T_{45}$  : Category two of Yang–Mills theory, as the former contains additional terms ;Kaluza–Klein theory is in general

$T_{46}$  : Category three of Yang–Mills theory, as the former contains additional terms;Kaluza–Klein theory is in general

**The Coefficients:**

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}, (b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$ ;  
 $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$  ,  
 $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}, (a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)}$ ;  
 $(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$   
 $(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$   
 $(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$   
 $(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)}$ ;  
 $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}, (a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{28})^{(5)}$ ;  
 $(a'_{29})^{(5)}, (a'_{30})^{(5)}, (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$   
 $(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)}$ ;  
 $(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)}$ ;  
 $(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)}$ ;

are Dissipation coefficients

**Module Numbered One**

The differential system of this model is now (Module Numbered one)

$$\begin{aligned} \frac{dG_{13}}{dt} &= (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} & 1 \\ \frac{dG_{14}}{dt} &= (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} & 2 \\ \frac{dG_{15}}{dt} &= (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} & 3 \\ \frac{dT_{13}}{dt} &= (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} & 4 \\ \frac{dT_{14}}{dt} &= (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} & 5 \\ \frac{dT_{15}}{dt} &= (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} & 6 \end{aligned}$$

$+(a''_{13})^{(1)}(T_{14}, t) =$  First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$  First detritions factor

**Module Numbered Two**

**The differential system of this model is now ( Module numbered two)**

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$  First augmentation factor

$-(b''_{16})^{(2)}((G_{19}), t) =$  First detritions factor

**Module Numbered Three**

**The differential system of this model is now (Module numbered three)**

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$  First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$  First detritions factor

**Module Numbered Four**

**The differential system of this model is now (Module numbered Four)**

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$  First augmentation factor

$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor}$

**Module Numbered Five:**

**The differential system of this model is now (Module number five)**

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor}$

$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor}$

**Module Numbered Six**

**The differential system of this model is now (Module numbered Six)**

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor}$

**Module Numbered Seven:**

**The differential system of this model is now (Seventh Module)**

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) = \text{First augmentation factor}$

### Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)]T_{42} \quad 48$$

### Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45}, t)$  = **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}), t)$  = **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[ \begin{array}{|c|c|c|c|} \hline (a'_{13})^{(1)} & + (a''_{13})^{(1)}(T_{14}, t) & + (a''_{16})^{(2,2)}(T_{17}, t) & + (a''_{20})^{(3,3)}(T_{21}, t) \\ \hline \end{array} \right] G_{13} \quad 55$$

$$\left[ \begin{array}{|c|c|c|} \hline + (a''_{24})^{(4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6)}(T_{33}, t) \\ \hline \end{array} \right] G_{13}$$

$$\left[ \begin{array}{|c|c|c|} \hline + (a''_{36})^{(7,7)}(T_{37}, t) & + (a''_{40})^{(8,8)}(T_{41}, t) & + (a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \\ \hline \end{array} \right] G_{13}$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[ \begin{array}{|c|c|c|c|} \hline (a'_{14})^{(1)} & + (a''_{14})^{(1)}(T_{14}, t) & + (a''_{17})^{(2,2)}(T_{17}, t) & + (a''_{21})^{(3,3)}(T_{21}, t) \\ \hline \end{array} \right] G_{14} \quad 56$$

$$\left[ \begin{array}{|c|c|c|} \hline + (a''_{25})^{(4,4,4,4)}(T_{25}, t) & + (a''_{29})^{(5,5,5,5)}(T_{29}, t) & + (a''_{33})^{(6,6,6,6)}(T_{33}, t) \\ \hline \end{array} \right] G_{14}$$

$$\left[ \begin{array}{|c|c|c|} \hline + (a''_{37})^{(7,7)}(T_{37}, t) & + (a''_{41})^{(8,8)}(T_{41}, t) & + (a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \\ \hline \end{array} \right] G_{14}$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[ \begin{array}{|c|c|c|c|} \hline (a'_{15})^{(1)} & + (a''_{15})^{(1)}(T_{14}, t) & + (a''_{18})^{(2,2)}(T_{17}, t) & + (a''_{22})^{(3,3)}(T_{21}, t) \\ \hline \end{array} \right] G_{15} \quad 57$$

$$\left[ \begin{array}{|c|c|c|} \hline + (a''_{26})^{(4,4,4,4)}(T_{25}, t) & + (a''_{30})^{(5,5,5,5)}(T_{29}, t) & + (a''_{34})^{(6,6,6,6)}(T_{33}, t) \\ \hline \end{array} \right] G_{15}$$

$$\left[ \begin{array}{|c|c|c|} \hline + (a''_{38})^{(7,7)}(T_{37}, t) & + (a''_{42})^{(8,8)}(T_{41}, t) & + (a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t) \\ \hline \end{array} \right] G_{15}$$

Where  $\boxed{(a''_{13})^{(1)}(T_{14}, t)}$ ,  $\boxed{(a''_{14})^{(1)}(T_{14}, t)}$ ,  $\boxed{(a''_{15})^{(1)}(T_{14}, t)}$  are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2)}(T_{17}, t)}$  are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3)}(T_{21}, t)}$  are third augmentation coefficient for

category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)}$  are fourth augmentation

coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)}$  are fifth augmentation coefficient

for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)}$  are sixth augmentation coefficient

for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{37})^{(7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{36})^{(7,7)}(T_{37}, t)}$  are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{41})^{(8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{42})^{(8,8)}(T_{41}, t)}$  are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45}, t)}$  are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[ \begin{array}{l} \boxed{(b'_{13})^{(1)} \boxed{-(b''_{13})^{(1)}(G, t)} \boxed{-(b''_{16})^{(2,2)}(G_{19}, t)} \boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}} \\ \boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7)}(G_{39}, t)} \boxed{-(b''_{40})^{(8,8)}(G_{43}, t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[ \begin{array}{l} \boxed{(b'_{14})^{(1)} \boxed{-(b''_{14})^{(1)}(G, t)} \boxed{-(b''_{17})^{(2,2)}(G_{19}, t)} \boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}} \\ \boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7)}(G_{39}, t)} \boxed{-(b''_{41})^{(8,8)}(G_{43}, t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[ \begin{array}{l} \boxed{(b'_{15})^{(1)} \boxed{-(b''_{15})^{(1)}(G, t)} \boxed{-(b''_{18})^{(2,2)}(G_{19}, t)} \boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}} \\ \boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7)}(G_{39}, t)} \boxed{-(b''_{42})^{(8,8)}(G_{43}, t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15} \quad 60$$

Where  $\boxed{-(b''_{13})^{(1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1)}(G, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2)}(G_{19}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)}$  are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)}$  are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)}$  are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7)}(G_{39}, t)}$  are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{41})^{(8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{42})^{(8,8)}(G_{43}, t)}$  are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)}$  are ninth detrition

coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)} G_{17} - \left[ \begin{array}{c} (a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) + (a''_{13})^{(1,1)}(T_{14}, t) + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{36})^{(7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9)}(T_{45}, t) \end{array} \right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)} G_{16} - \left[ \begin{array}{c} (a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t) + (a''_{14})^{(1,1)}(T_{14}, t) + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{37})^{(7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9)}(T_{45}, t) \end{array} \right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[ \begin{array}{c} (a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t) + (a''_{15})^{(1,1)}(T_{14}, t) + (a''_{22})^{(3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6)}(T_{33}, t) \\ + (a''_{38})^{(7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9)}(T_{45}, t) \end{array} \right] G_{18} \quad 63$$

Where  $(a'_{16})^{(2)}$ ,  $(a'_{17})^{(2)}$ ,  $(a'_{18})^{(2)}$  are first augmentation coefficients for category 1, 2 and 3

$(a''_{13})^{(1,1)}$ ,  $(a''_{14})^{(1,1)}$ ,  $(a''_{15})^{(1,1)}$  are second augmentation coefficient for category 1, 2 and 3

$(a''_{20})^{(3,3,3)}$ ,  $(a''_{21})^{(3,3,3)}$ ,  $(a''_{22})^{(3,3,3)}$  are third augmentation coefficient for category 1, 2 and 3

$(a''_{24})^{(4,4,4,4,4)}$ ,  $(a''_{25})^{(4,4,4,4,4)}$ ,  $(a''_{26})^{(4,4,4,4,4)}$  are fourth augmentation coefficient for category 1, 2 and 3

$(a''_{28})^{(5,5,5,5,5)}$ ,  $(a''_{29})^{(5,5,5,5,5)}$ ,  $(a''_{30})^{(5,5,5,5,5)}$  are fifth augmentation coefficient for category 1, 2 and 3

$(a''_{32})^{(6,6,6,6,6)}$ ,  $(a''_{33})^{(6,6,6,6,6)}$ ,  $(a''_{34})^{(6,6,6,6,6)}$  are sixth augmentation coefficient for category 1, 2 and 3

$(a''_{36})^{(7,7,7)}$ ,  $(a''_{37})^{(7,7,7)}$ ,  $(a''_{38})^{(7,7,7)}$  are seventh augmentation coefficient for category 1, 2 and 3

$(a''_{40})^{(8,8,8)}$ ,  $(a''_{41})^{(8,8,8)}$ ,  $(a''_{42})^{(8,8,8)}$  are eight augmentation coefficient for category 1, 2 and 3

$(a''_{44})^{(9,9)}$ ,  $(a''_{45})^{(9,9)}$ ,  $(a''_{46})^{(9,9)}$  are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[ \begin{array}{c} (b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}, t) - (b''_{13})^{(1,1)}(G, t) - (b''_{20})^{(3,3,3)}(G_{23}, t) \\ - (b''_{24})^{(4,4,4,4,4)}(G_{27}, t) - (b''_{28})^{(5,5,5,5,5)}(G_{31}, t) - (b''_{32})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b''_{36})^{(7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9)}(G_{47}, t) \end{array} \right] T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[ \begin{array}{c} (b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}, t) - (b''_{14})^{(1,1)}(G, t) - (b''_{21})^{(3,3,3)}(G_{23}, t) \\ - (b''_{25})^{(4,4,4,4,4)}(G_{27}, t) - (b''_{29})^{(5,5,5,5,5)}(G_{31}, t) - (b''_{33})^{(6,6,6,6,6)}(G_{35}, t) \\ - (b''_{37})^{(7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9)}(G_{47}, t) \end{array} \right] T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[ \begin{array}{l} (b'_{18})^{(2)} \boxed{-(b''_{18})^{(2)}(G_{19}, t)} \boxed{-(b''_{15})^{(1,1)}(G, t)} \boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)} \boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)} \boxed{-(b''_{46})^{(9,9)}(G_{47}, t)} \end{array} \right] T_{18} \tag{66}$$

where  $\boxed{-(b''_{16})^{(2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2)}(G_{19}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1)}(G, t)}$  are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)}$  are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$  are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$  are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$  are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7,7)}(G_{39}, t)}$  are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{41})^{(8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{42})^{(8,8,8)}(G_{43}, t)}$  are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{46})^{(9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{45})^{(9,9)}(G_{47}, t)}$  are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[ \begin{array}{l} (a'_{20})^{(3)} \boxed{+(a''_{20})^{(3)}(T_{21}, t)} \boxed{+(a''_{16})^{(2,2,2)}(T_{17}, t)} \boxed{+(a''_{13})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)} \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41}, t)} \boxed{+(a''_{44})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{20} \tag{67}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[ \begin{array}{l} (a'_{21})^{(3)} \boxed{+(a''_{21})^{(3)}(T_{21}, t)} \boxed{+(a''_{17})^{(2,2,2)}(T_{17}, t)} \boxed{+(a''_{14})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)} \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41}, t)} \boxed{+(a''_{45})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{21} \tag{68}$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[ \begin{array}{l} (a'_{22})^{(3)} \boxed{+(a''_{22})^{(3)}(T_{21}, t)} \boxed{+(a''_{18})^{(2,2,2)}(T_{17}, t)} \boxed{+(a''_{15})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37}, t)} \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41}, t)} \boxed{+(a''_{46})^{(9,9,9)}(T_{45}, t)} \end{array} \right] G_{22} \tag{69}$$

$\boxed{+(a''_{20})^{(3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3)}(T_{21}, t)}$  are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2)}(T_{17}, t)}$  are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1)}(T_{14}, t)}$  are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)}$  are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)}$  are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)}$  are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37}, t)}$  are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41}, t)}$  are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{45})^{(9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{46})^{(9,9,9)}(T_{45}, t)}$  are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[ \begin{array}{l} \boxed{(b'_{20})^{(3)} \boxed{-(b''_{20})^{(3)}(G_{23}, t)} \boxed{-(b''_{16})^{(2,2,2)}(G_{19}, t)} \boxed{-(b''_{13})^{(1,1,1)}(G, t)}} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39}, t)} \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43}, t)} \boxed{-(b''_{44})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{20} \quad 70$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[ \begin{array}{l} \boxed{(b'_{21})^{(3)} \boxed{-(b''_{21})^{(3)}(G_{23}, t)} \boxed{-(b''_{17})^{(2,2,2)}(G_{19}, t)} \boxed{-(b''_{14})^{(1,1,1)}(G, t)}} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39}, t)} \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43}, t)} \boxed{-(b''_{45})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{21} \quad 71$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[ \begin{array}{l} \boxed{(b'_{22})^{(3)} \boxed{-(b''_{22})^{(3)}(G_{23}, t)} \boxed{-(b''_{18})^{(2,2,2)}(G_{19}, t)} \boxed{-(b''_{15})^{(1,1,1)}(G, t)}} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39}, t)} \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43}, t)} \boxed{-(b''_{46})^{(9,9,9)}(G_{47}, t)} \end{array} \right] T_{22} \quad 72$$

$\boxed{-(b''_{20})^{(3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3)}(G_{23}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2)}(G_{19}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1)}(G, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)}$  are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)}$  are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)}$  are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39}, t)}$  are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43}, t)}$  are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{45})^{(9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{44})^{(9,9,9)}(G_{47}, t)}$  are ninth detrition coefficients for

category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[ \begin{array}{l} (a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) + (a''_{28})^{(5,5)}(T_{29}, t) + (a''_{32})^{(6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[ \begin{array}{l} (a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t) + (a''_{29})^{(5,5)}(T_{29}, t) + (a''_{33})^{(6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[ \begin{array}{l} (a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t) + (a''_{30})^{(5,5)}(T_{29}, t) + (a''_{34})^{(6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9)}(T_{45}, t) \end{array} \right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25}, t), (a''_{25})^{(4)}(T_{25}, t), (a''_{26})^{(4)}(T_{25}, t)$  are first augmentation coefficients

category 1, 2 3

$(a''_{28})^{(5,5)}(T_{29}, t), (a''_{29})^{(5,5)}(T_{29}, t), (a''_{30})^{(5,5)}(T_{29}, t)$  are second augmentation coefficient for category 1, 2 and 3

$(a''_{32})^{(6,6)}(T_{33}, t), (a''_{33})^{(6,6)}(T_{33}, t), (a''_{34})^{(6,6)}(T_{33}, t)$  are third augmentation coefficient for category 1, 2 and 3

$(a''_{13})^{(1,1,1,1)}(T_{14}, t), (a''_{14})^{(1,1,1,1)}(T_{14}, t), (a''_{15})^{(1,1,1,1)}(T_{14}, t)$  are fourth augmentation coefficient

$(a''_{16})^{(2,2,2,2)}(T_{17}, t),$

$(a''_{17})^{(2,2,2,2)}(T_{17}, t), (a''_{18})^{(2,2,2,2)}(T_{17}, t)$  are fifth augmentation coefficients for category 1, 2 and 3

$(a''_{20})^{(3,3,3,3)}(T_{21}, t), (a''_{21})^{(3,3,3,3)}(T_{21}, t),$

$(a''_{22})^{(3,3,3,3)}(T_{21}, t)$  are sixth augmentation coefficients for category 1, 2 and 3

$(a''_{36})^{(7,7,7,7)}(T_{37}, t), (a''_{37})^{(7,7,7,7)}(T_{37}, t),$

$(a''_{38})^{(7,7,7,7)}(T_{37}, t)$  are seventh augmentation coefficients for category 1, 2 and 3

$(a''_{40})^{(8,8,8,8)}(T_{41}, t), (a''_{41})^{(8,8,8,8)}(T_{41}, t), (a''_{42})^{(8,8,8,8)}(T_{41}, t)$

are eighth augmentation coefficients for category 1, 2 and 3

$(a''_{46})^{(9,9,9,9)}(T_{45}, t), (a''_{45})^{(9,9,9,9)}(T_{45}, t), (a''_{44})^{(9,9,9,9)}(T_{45}, t)$  are ninth detrition coefficients for

category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[ \begin{array}{l} (b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}, t) - (b''_{28})^{(5,5)}(G_{31}, t) - (b''_{32})^{(6,6)}(G_{35}, t) \\ - (b''_{13})^{(1,1,1,1)}(G, t) - (b''_{16})^{(2,2,2,2)}(G_{19}, t) - (b''_{20})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[ \begin{array}{l} (b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}, t) - (b''_{29})^{(5,5)}(G_{31}, t) - (b''_{33})^{(6,6)}(G_{35}, t) \\ - (b''_{14})^{(1,1,1,1)}(G, t) - (b''_{17})^{(2,2,2,2)}(G_{19}, t) - (b''_{21})^{(3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9)}(G_{47}, t) \end{array} \right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[ \begin{array}{l} (b'_{26})^{(4)} \boxed{-(b''_{26})^{(4)}(G_{27}, t)} \boxed{-(b''_{30})^{(5,5)}(G_{31}, t)} \boxed{-(b''_{34})^{(6,6)}(G_{35}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G, t)} \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19}, t)} \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39}, t)} \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43}, t)} \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{26} \tag{78}$$

Where  $\boxed{-(b''_{24})^{(4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4)}(G_{27}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5)}(G_{31}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6)}(G_{35}, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1,1)}(G, t)}$  are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2,2)}(G_{19}, t)}$  are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3,3)}(G_{23}, t)}$  are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39}, t)}$  are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43}, t)}$  are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47}, t)}$  are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[ \begin{array}{l} (a'_{28})^{(5)} \boxed{+(a''_{28})^{(5)}(T_{29}, t)} \boxed{+(a''_{24})^{(4,4)}(T_{25}, t)} \boxed{+(a''_{32})^{(6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)} \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)} \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)} \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)} \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{28} \tag{79}$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[ \begin{array}{l} (a'_{29})^{(5)} \boxed{+(a''_{29})^{(5)}(T_{29}, t)} \boxed{+(a''_{25})^{(4,4)}(T_{25}, t)} \boxed{+(a''_{33})^{(6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)} \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)} \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)} \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)} \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{29} \tag{80}$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[ \begin{array}{l} (a'_{30})^{(5)} \boxed{+(a''_{30})^{(5)}(T_{29}, t)} \boxed{+(a''_{26})^{(4,4)}(T_{25}, t)} \boxed{+(a''_{34})^{(6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)} \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)} \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)} \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)} \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{30} \tag{81}$$

Where  $\boxed{+(a''_{28})^{(5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5)}(T_{29}, t)}$  are first augmentation coefficients for category 1, 2 and 3

And  $\boxed{+(a''_{24})^{(4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4)}(T_{25}, t)}$  are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6)}(T_{33}, t)}$  are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14}, t)}$  are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17}, t)}$  are fifth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21}, t)}$  are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37}, t)}$  are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41}, t)}$  are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45}, t)}$  are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[ \begin{array}{l} \boxed{(b'_{28})^{(5)} - \boxed{(b''_{28})^{(5)}(G_{31}, t)} - \boxed{(b''_{24})^{(4,4)}(G_{27}, t)} - \boxed{(b''_{32})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)} - \boxed{(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)} - \boxed{(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)} - \boxed{(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)} - \boxed{(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[ \begin{array}{l} \boxed{(b'_{29})^{(5)} - \boxed{(b''_{29})^{(5)}(G_{31}, t)} - \boxed{(b''_{25})^{(4,4)}(G_{27}, t)} - \boxed{(b''_{33})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)} - \boxed{(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)} - \boxed{(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)} - \boxed{(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)} - \boxed{(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[ \begin{array}{l} \boxed{(b'_{30})^{(5)} - \boxed{(b''_{30})^{(5)}(G_{31}, t)} - \boxed{(b''_{26})^{(4,4)}(G_{27}, t)} - \boxed{(b''_{34})^{(6,6,6)}(G_{35}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)} - \boxed{(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)} - \boxed{(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)} - \boxed{(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)} - \boxed{(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{30} \quad 84$$

where  $\boxed{-(b''_{28})^{(5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5)}(G_{31}, t)}$  are first detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4)}(G_{27}, t)}$  are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6)}(G_{35}, t)}$  are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1,1,1)}(G, t)}$  are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19}, t)}$  are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23}, t)}$  are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39}, t)}$  are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43}, t)}$  are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47}, t)}$  are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - \left[ \begin{array}{l} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) + (a''_{28})^{(5,5,5)}(T_{29}, t) + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{32} \quad 85$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - \left[ \begin{array}{l} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) + (a''_{29})^{(5,5,5)}(T_{29}, t) + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{33} \quad 86$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - \left[ \begin{array}{l} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) + (a''_{30})^{(5,5,5)}(T_{29}, t) + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t) + (a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t) \end{array} \right] G_{34} \quad 87$$

$+(a''_{32})^{(6)}(T_{33}, t), +(a''_{33})^{(6)}(T_{33}, t), +(a''_{34})^{(6)}(T_{33}, t)$  are first augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5)}(T_{29}, t), +(a''_{29})^{(5,5,5)}(T_{29}, t), +(a''_{30})^{(5,5,5)}(T_{29}, t)$  are second augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4)}(T_{25}, t), +(a''_{25})^{(4,4,4)}(T_{25}, t), +(a''_{26})^{(4,4,4)}(T_{25}, t)$  are third augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), +(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$  - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), +(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$  - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), +(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$  sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{37})^{(7,7,7,7,7,7)}(T_{37}, t), +(a''_{38})^{(7,7,7,7,7,7)}(T_{37}, t)$  seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t), +(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)$

Eighth augmentation coefficients

$+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t), +(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)$  ninth augmentation coefficients

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - \left[ \begin{array}{l} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) - (b''_{28})^{(5,5,5)}(G_{31}, t) - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{32} \quad 88$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - \left[ \begin{array}{l} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) - (b''_{29})^{(5,5,5)}(G_{31}, t) - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \\ - (b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t) - (b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t) - (b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t) \end{array} \right] T_{33} \quad 89$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[ \begin{array}{l} (b'_{34})^{(6)} \boxed{-(b''_{34})^{(6)}(G_{35}, t)} \boxed{-(b''_{30})^{(5,5,5)}(G_{31}, t)} \boxed{-(b''_{26})^{(4,4,4)}(G_{27}, t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G, t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)} \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{34} \quad 90$$

$\boxed{-(b''_{32})^{(6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6)}(G_{35}, t)}$  are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5)}(G_{31}, t)}$  are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4)}(G_{27}, t)}$  are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G, t)}$ ,  $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G, t)}$  are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)}$  are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)}$  are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39}, t)}$  are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43}, t)}$ ,  $\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43}, t)}$  are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47}, t)}$ ,  $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47}, t)}$  are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} \quad 91$$

$$- \left[ \begin{array}{l} (a'_{36})^{(7)} \boxed{+(a''_{36})^{(7)}(T_{37}, t)} \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t)} \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t)} \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41}, t)} \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{13}$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} \quad 92$$

$$- \left[ \begin{array}{l} (a'_{37})^{(7)} \boxed{+(a''_{37})^{(7)}(T_{37}, t)} \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t)} \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t)} \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41}, t)} \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{14}$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} \quad 93$$

$$- \left[ \begin{array}{l} (a'_{38})^{(7)} \boxed{+(a''_{38})^{(7)}(T_{37}, t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)} \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41}, t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{15}$$

Where  $\boxed{(a''_{36})^{(7)}(T_{37}, t)}$ ,  $\boxed{(a''_{37})^{(7)}(T_{37}, t)}$ ,  $\boxed{(a''_{38})^{(7)}(T_{37}, t)}$  are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$  are second augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$  are third augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$  are fourth augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$  are fifth augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$  are sixth augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$  are seventh augmentation coefficient for category 1, 2 and 3  
 $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41}, t)}$  are eighth augmentation coefficient for 1,2,3  
 $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$  are ninth augmentation coefficient for 1,2,3

$$\begin{aligned}
 \frac{dT_{36}}{dt} &= (b_{36})^{(7)}T_{37} - \left[ \begin{array}{l} \boxed{(b'_{36})^{(7)} - \boxed{(b''_{36})^{(7)}(G_{39}, t) - \boxed{(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t) - \boxed{(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t) - \boxed{(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t) - \boxed{(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{(b''_{13})^{(1,1,1,1,1,1,1)}(G, t) - \boxed{(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t) - \boxed{(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{13} \\
 \frac{dT_{37}}{dt} &= (b_{37})^{(7)}T_{36} - \left[ \begin{array}{l} \boxed{(b'_{37})^{(7)} - \boxed{(b''_{37})^{(7)}(G_{39}, t) - \boxed{(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t) - \boxed{(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t) - \boxed{(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t) - \boxed{(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{(b''_{14})^{(1,1,1,1,1,1,1)}(G, t) - \boxed{(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t) - \boxed{(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{14} \\
 \frac{dT_{38}}{dt} &= (b_{38})^{(7)}T_{37} - \left[ \begin{array}{l} \boxed{(b'_{38})^{(7)} - \boxed{(b''_{38})^{(7)}(G_{39}, t) - \boxed{(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t) - \boxed{(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)} \\ \boxed{(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t) - \boxed{(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t) - \boxed{(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)} \\ \boxed{(b''_{15})^{(1,1,1,1,1,1,1)}(G, t) - \boxed{(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t) - \boxed{(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)} \end{array} \right] T_{15}
 \end{aligned}$$

Where  $\boxed{-(b''_{36})^{(7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{37})^{(7)}(G_{39}, t)}$ ,  $\boxed{-(b''_{38})^{(7)}(G_{39}, t)}$  are first detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$ ,  $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19}, t)}$  are second detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$ ,  $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23}, t)}$  are third detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$ ,  $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27}, t)}$  are fourth detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$ ,  $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31}, t)}$  are fifth detrition coefficients for category 1, 2 and 3  
 $\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$ ,  $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35}, t)}$  are sixth detrition

coefficients for category 1, 2 and 3

$$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G, t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G, t)}, \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G, t)}$$

are seventh detrition coefficients for category 1, 2 and 3

$$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43}, t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43}, t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43}, t)}$$

coefficients for category 1, 2 and 3

$$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47}, t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47}, t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47}, t)}$$

coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

95

$$= (a_{40})^{(8)} G_{41} - \left[ \begin{array}{l} \boxed{(a'_{40})^{(8)}} + \boxed{(a''_{40})^{(8)}(T_{41}, t)} + \boxed{(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)} + \boxed{(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)} + \boxed{(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)} + \boxed{(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} + \boxed{(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t)} + \boxed{(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{13}$$

$$\frac{dG_{41}}{dt}$$

$$= (a_{41})^{(8)} G_{40} - \left[ \begin{array}{l} \boxed{(a'_{41})^{(8)}} + \boxed{(a''_{41})^{(8)}(T_{41}, t)} + \boxed{(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)} + \boxed{(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)} + \boxed{(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)} + \boxed{(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)} + \boxed{(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t)} + \boxed{(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{14}$$

$$\frac{dG_{42}}{dt}$$

$$= (a_{42})^{(8)} G_{41} - \left[ \begin{array}{l} \boxed{(a'_{42})^{(8)}} + \boxed{(a''_{42})^{(8)}(T_{41}, t)} + \boxed{(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)} + \boxed{(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)} \\ \boxed{(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)} + \boxed{(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)} + \boxed{(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)} \\ \boxed{(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)} + \boxed{(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t)} + \boxed{(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)} \end{array} \right] G_{15}$$

Where  $\boxed{+(a'_{40})^{(8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{41})^{(8)}(T_{41}, t)}$ ,  $\boxed{+(a''_{42})^{(8)}(T_{41}, t)}$  are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$ ,  $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17}, t)}$  are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$ ,  $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21}, t)}$  are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$ ,  $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25}, t)}$  are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$ ,  $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29}, t)}$  are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$ ,  $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33}, t)}$  are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$ ,  $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14}, t)}$  are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37}, t)}$ ,  $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37}, t)}$  are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$ ,  $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45}, t)}$  are ninth

augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[ \begin{array}{|c|c|c|} \hline (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43}, t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ \hline \hline -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ \hline \hline -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \\ \hline \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[ \begin{array}{|c|c|c|} \hline (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43}, t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ \hline \hline -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ \hline \hline -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \\ \hline \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[ \begin{array}{|c|c|c|} \hline (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43}, t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t) \\ \hline \hline -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35}, t) \\ \hline \hline -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t) \\ \hline \end{array} \right] T_{15}$$

Where  $-(b''_{36})^{(7)}(G_{39}, t)$ ,  $-(b''_{37})^{(7)}(G_{39}, t)$ ,  $-(b''_{38})^{(7)}(G_{39}, t)$  are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19}, t)$  are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23}, t)$  are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27}, t)$  are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31}, t)$  are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G, t)$  are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{38})^{(7,7)}(G_{39}, t)$  are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39}, t)$  are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$ ,  $-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47}, t)$  are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[ \begin{array}{l} (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) + (a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[ \begin{array}{l} (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t) + (a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[ \begin{array}{l} (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{37}, t) + (a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t) + (a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t) \\ + (a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t) + (a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t) + (a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t) \\ + (a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t) + (a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t) + (a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t) \end{array} \right] G_{15}$$

Where  $(a'_{44})^{(9)}(T_{45}, t)$ ,  $(a'_{45})^{(9)}(T_{45}, t)$ ,  $(a'_{46})^{(9)}(T_{37}, t)$  are first augmentation coefficients for category 1, 2 and 3

$(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$ ,  $(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17}, t)$  are second augmentation coefficient for category 1, 2 and 3

$(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$ ,  $(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$ ,  $(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21}, t)$  are third augmentation coefficient for category 1, 2 and 3

$(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$ ,  $(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$ ,  $(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25}, t)$  are fourth augmentation coefficient for category 1, 2 and 3

$(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$ ,  $(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$ ,  $(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29}, t)$  are fifth augmentation coefficient for category 1, 2 and 3

$(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$ ,  $(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$ ,  $(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33}, t)$  are sixth augmentation coefficient for category 1, 2 and 3

$(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$ ,  $(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$ ,  $(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14}, t)$  are Seventh augmentation coefficient for category 1, 2 and 3

$(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$ ,  $(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$ ,  $(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37}, t)$  are eighth augmentation coefficient for 1,2,3

$(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$ ,  $(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$ ,  $(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41}, t)$  are ninth augmentation coefficient for 1,2,3

$$\begin{aligned} & \frac{dT_{44}}{dt} \\ &= (b_{44})^{(9)}T_{45} \\ &= \left[ \begin{array}{l} (b'_{44})^{(9)} \left[ \begin{array}{l} -(b''_{44})^{(9)}(G_{47}, t) \quad | \quad -(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19}, t) \quad | \quad -(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27}, t) \quad | \quad -(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31}, t) \quad | \quad -(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35}, t) \end{array} \right] \\ -(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G, t) \quad | \quad -(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39}, t) \quad | \quad -(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{13} \\ \\ & \frac{dT_{45}}{dt} \\ &= (b_{45})^{(9)}T_{44} - \left[ \begin{array}{l} (b'_{45})^{(9)} \left[ \begin{array}{l} -(b''_{45})^{(9)}(G_{47}, t) \quad | \quad -(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19}, t) \quad | \quad -(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27}, t) \quad | \quad -(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31}, t) \quad | \quad -(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35}, t) \end{array} \right] \\ -(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G, t) \quad | \quad -(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39}, t) \quad | \quad -(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{14} \\ \\ & \frac{dT_{46}}{dt} \\ &= (b_{46})^{(9)}T_{45} - \left[ \begin{array}{l} (b'_{46})^{(9)} \left[ \begin{array}{l} -(b''_{46})^{(9)}(G_{47}, t) \quad | \quad -(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19}, t) \quad | \quad -(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23}, t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27}, t) \quad | \quad -(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31}, t) \quad | \quad -(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35}, t) \end{array} \right] \\ -(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G, t) \quad | \quad -(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39}, t) \quad | \quad -(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43}, t) \end{array} \right] T_{15} \end{aligned}$$

Where  $-(b''_{44})^{(9)}(G_{47}, t)$ ,  $-(b''_{45})^{(9)}(G_{47}, t)$ ,  $-(b''_{46})^{(9)}(G_{47}, t)$  are first detrition coefficients for category 1, 2 and 3  
 $-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19}, t)$ ,  $-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19}, t)$  are second detrition coefficients for category 1, 2 and 3  
 $-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23}, t)$ ,  $-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23}, t)$  are third detrition coefficients for category 1, 2 and 3  
 $-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27}, t)$ ,  $-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27}, t)$  are fourth detrition coefficients for category 1, 2 and 3  
 $-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31}, t)$ ,  $-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31}, t)$  are fifth detrition coefficients for category 1, 2 and 3  
 $-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35}, t)$ ,  $-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35}, t)$  are sixth detrition coefficients for category 1, 2 and 3  
 $-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G, t)$ ,  $-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G, t)$  are seventh detrition coefficients for category 1, 2 and 3  
 $-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39}, t)$ ,  $-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39}, t)$  are eighth detrition coefficients for category 1, 2 and 3  
 $-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43}, t)$ ,  $-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43}, t)$  are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, i, j = 13,14,15$$

The functions  $(a''_i)^{(1)}, (b''_i)^{(1)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(1)}, (r_i)^{(1)}$ :

$$(a_i'')^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(1)}(T_{14}, t) = (p_i)^{(1)} \tag{98}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(1)}(G, t) = (r_i)^{(1)}$$

**Definition of**  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$  :

Where  $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$  are positive constants and  $i = 13, 14, 15$

They satisfy Lipschitz condition:

$$|(a_i'')^{(1)}(T'_{14}, t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T'_{14} - T_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} |G - G'| e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(1)}(T'_{14}, t)$  and  $(a_i'')^{(1)}(T_{14}, t)$ .  $(T'_{14}, t)$  and  $(T_{14}, t)$  are points belonging to the interval  $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$ . It is to be noted that  $(a_i'')^{(1)}(T_{14}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{13})^{(1)} = 1$  then the function  $(a_i'')^{(1)}(T_{14}, t)$ , the first augmentation coefficient attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$  :

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$ , are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

**Definition of**  $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$  :

There exists two constants  $(\hat{P}_{13})^{(1)}$  and  $(\hat{Q}_{13})^{(1)}$  which together With  $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$  and  $(\hat{B}_{13})^{(1)}$  and the constants  $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13, 14, 15$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, i, j = 16, 17, 18$$

The functions  $(a_i'')^{(2)}, (b_i'')^{(2)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(2)}, (r_i)^{(2)}$ :

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \tag{102}$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \tag{103}$$

$$\lim_{T_2 \rightarrow \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \tag{104}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(2)}(G_{19}, t) = (r_i)^{(2)} \tag{105}$$

**Definition of**  $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$  : 106

Where  $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$  are positive constants and  $i = 16, 17, 18$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17}' - T_{17}| e^{-(\hat{M}_{16})^{(2)}t} \tag{107}$$

$$|(b_i'')^{(2)}(G_{19}', t) - (b_i'')^{(2)}(G_{19}, t)| < (\hat{k}_{16})^{(2)} |(G_{19}') - (G_{19})| e^{-(\hat{M}_{16})^{(2)}t} \tag{108}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(2)}(T_{17}', t)$  and  $(a_i'')^{(2)}(T_{17}, t)$ .  $(T_{17}', t)$  and  $(T_{17}, t)$  are points belonging to the interval  $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$ . It is to be noted that  $(a_i'')^{(2)}(T_{17}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{16})^{(2)} = 1$  then the function  $(a_i'')^{(2)}(T_{17}, t)$ , the first augmentation coefficient attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$  :

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$ , are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}} , \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

**Definition of**  $(\hat{P}_{16})^{(2)}, (\hat{Q}_{16})^{(2)}$  :

There exists two constants  $(\hat{P}_{16})^{(2)}$  and  $(\hat{Q}_{16})^{(2)}$  which together with  $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$  and  $(\hat{B}_{16})^{(2)}$  and the constants  $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16, 17, 18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \tag{110}$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \tag{111}$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, i, j = 20, 21, 22 \quad 112$$

The functions  $(a_i'')^{(3)}, (b_i'')^{(3)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(3)}, (r_i)^{(3)}$ :

$$\begin{aligned} (a_i'')^{(3)}(T_{21}, t) &\leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)} \\ (b_i'')^{(3)}(G_{23}, t) &\leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)} \\ \lim_{T_2 \rightarrow \infty} (a_i'')^{(3)}(T_{21}, t) &= (p_i)^{(3)} \\ \lim_{G \rightarrow \infty} (b_i'')^{(3)}(G_{23}, t) &= (r_i)^{(3)} \end{aligned} \quad 113$$

**Definition of**  $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$ :

Where  $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$  are positive constants and  $i = 20, 21, 22$

They satisfy Lipschitz condition:

$$\begin{aligned} |(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| &\leq (\hat{k}_{20})^{(3)} |T_{21} - T_{21}'| e^{-(\hat{M}_{20})^{(3)}t} \\ |(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| &< (\hat{k}_{20})^{(3)} |G_{23} - G_{23}'| e^{-(\hat{M}_{20})^{(3)}t} \end{aligned} \quad 114$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(3)}(T_{21}', t)$  and  $(a_i'')^{(3)}(T_{21}, t)$ .  $(T_{21}', t)$  and  $(T_{21}, t)$  are points belonging to the interval  $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$ . It is to be noted that  $(a_i'')^{(3)}(T_{21}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{20})^{(3)} = 1$  then the function  $(a_i'')^{(3)}(T_{21}, t)$ , the first augmentation coefficient attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$ :

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$ , are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1 \quad 115$$

There exists two constants  $(\hat{P}_{20})^{(3)}$  and  $(\hat{Q}_{20})^{(3)}$  which together with  $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$  and  $(\hat{B}_{20})^{(3)}$  and the constants  $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$ , satisfy the inequalities

$$\begin{aligned} \frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] &< 1 \\ \frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] &< 1 \end{aligned} \quad 116$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, i, j = 24, 25, 26 \quad 117$$

The functions  $(a_i'')^{(4)}, (b_i'')^{(4)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(4)}, (r_i)^{(4)}$ :

$$\begin{aligned} (a_i'')^{(4)}(T_{25}, t) &\leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)} \\ (b_i'')^{(4)}((G_{27}), t) &\leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)} \\ \lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) &= (p_i)^{(4)} \\ \lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) &= (r_i)^{(4)} \end{aligned} \quad 118$$

**Definition of**  $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$ :

Where  $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$  are positive constants and  $i = 24, 25, 26$

They satisfy Lipschitz condition: 119

$$\begin{aligned} |(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| &\leq (\hat{k}_{24})^{(4)} |T_{25}' - T_{25}| e^{-(\hat{M}_{24})^{(4)}t} \\ |(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| &< (\hat{k}_{24})^{(4)} |(G_{27})' - (G_{27})| e^{-(\hat{M}_{24})^{(4)}t} \end{aligned}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a_i'')^{(4)}(T_{25}', t)$  and  $(a_i'')^{(4)}(T_{25}, t)$ .  $(T_{25}', t)$  and  $(T_{25}, t)$  are points belonging to the interval  $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$ . It is to be noted that  $(a_i'')^{(4)}(T_{25}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{24})^{(4)} = 1$  then the function  $(a_i'')^{(4)}(T_{25}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$ : 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$ , are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

**Definition of**  $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$ : 121

There exists two constants  $(\hat{P}_{24})^{(4)}$  and  $(\hat{Q}_{24})^{(4)}$  which together with  $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$  and  $(\hat{B}_{24})^{(4)}$  and the constants  $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [ (b_i)^{(4)} + (b'_i)^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)} ] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a'_i)^{(5)}, (a''_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (b''_i)^{(5)} > 0, i, j = 28,29,30 \tag{122}$$

The functions  $(a''_i)^{(5)}, (b''_i)^{(5)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(5)}, (r_i)^{(5)}$ :

$$(a''_i)^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b''_i)^{(5)}(G_{31}, t) \leq (r_i)^{(5)} \leq (b'_i)^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(5)}(T_{29}, t) = (p_i)^{(5)} \tag{123}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

**Definition of**  $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$ :

Where  $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$  are positive constants and  $i = 28,29,30$

They satisfy Lipschitz condition:

124

$$|(a''_i)^{(5)}(T'_{29}, t) - (a''_i)^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29} - T'_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b''_i)^{(5)}(G'_{31}, t) - (b''_i)^{(5)}(G_{31}, t)| < (\hat{k}_{28})^{(5)} |(G_{31}) - (G'_{31})| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(5)}(T'_{29}, t)$  and  $(a''_i)^{(5)}(T_{29}, t)$ .  $(T'_{29}, t)$  and  $(T_{29}, t)$  are points belonging to the interval  $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$ . It is to be noted that  $(a''_i)^{(5)}(T_{29}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{28})^{(5)} = 1$  then the function  $(a''_i)^{(5)}(T_{29}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$ :

125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$ , are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

**Definition of**  $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$ :

126

There exists two constants  $(\hat{P}_{28})^{(5)}$  and  $(\hat{Q}_{28})^{(5)}$  which together with  $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$  and  $(\hat{B}_{28})^{(5)}$  and the constants  $(a_i)^{(5)}, (a'_i)^{(5)}, (b_i)^{(5)}, (b'_i)^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a'_i)^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b'_i)^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a'_i)^{(6)}, (a''_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (b''_i)^{(6)} > 0, i, j = 32, 33, 34 \tag{127}$$

The functions  $(a''_i)^{(6)}, (b''_i)^{(6)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(6)}, (r_i)^{(6)}$ :

$$(a''_i)^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b''_i)^{(6)}(G_{35}, t) \leq (r_i)^{(6)} \leq (b'_i)^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(6)}(T_{33}, t) = (p_i)^{(6)} \tag{128}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(6)}(G_{35}, t) = (r_i)^{(6)}$$

**Definition of**  $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$ :

Where  $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$  are positive constants and  $i = 32, 33, 34$

They satisfy Lipschitz condition:

$$|(a''_i)^{(6)}(T'_{33}, t) - (a''_i)^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T'_{33}| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b''_i)^{(6)}(G'_{35}, t) - (b''_i)^{(6)}(G_{35}, t)| < (\hat{k}_{32})^{(6)} |(G_{35}) - (G'_{35})| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(6)}(T'_{33}, t)$  and  $(a''_i)^{(6)}(T_{33}, t)$ .  $(T'_{33}, t)$  and  $(T_{33}, t)$  are points belonging to the interval  $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$ . It is to be noted that  $(a''_i)^{(6)}(T_{33}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{32})^{(6)} = 1$  then the function  $(a''_i)^{(6)}(T_{33}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$ : 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$ , are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

**Definition of**  $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$ : 130

There exists two constants  $(\hat{P}_{32})^{(6)}$  and  $(\hat{Q}_{32})^{(6)}$  which together with  $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$  and  $(\hat{B}_{32})^{(6)}$  and the

constants  $(a_i)^{(6)}, (a'_i)^{(6)}, (b_i)^{(6)}, (b'_i)^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34$ , satisfy the inequalities

$$\frac{1}{(\widehat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a'_i)^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\widehat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b'_i)^{(6)} + (\widehat{B}_{32})^{(6)} + (\widehat{Q}_{32})^{(6)} (\widehat{k}_{32})^{(6)}] < 1$$

Where we suppose

(G)  $(a_i)^{(7)}, (a'_i)^{(7)}, (a''_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (b''_i)^{(7)} > 0, i, j = 36,37,38$  131

(H) The functions  $(a''_i)^{(7)}, (b''_i)^{(7)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(7)}, (r_i)^{(7)}$ :

$$(a''_i)^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\widehat{A}_{36})^{(7)}$$

$$(b''_i)^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b'_i)^{(7)} \leq (\widehat{B}_{36})^{(7)}$$

132

(I)  $\lim_{T_2 \rightarrow \infty} (a''_i)^{(7)}(T_{37}, t) = (p_i)^{(7)}$   
 (J)

$$\lim_{G \rightarrow \infty} (b''_i)^{(7)}(G_{39}, t) = (r_i)^{(7)}$$

**Definition of**  $(\widehat{A}_{36})^{(7)}, (\widehat{B}_{36})^{(7)}$  :

Where  $(\widehat{A}_{36})^{(7)}, (\widehat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}$  are positive constants and  $i = 36,37,38$

They satisfy Lipschitz condition:

133

$$|(a''_i)^{(7)}(T'_{37}, t) - (a''_i)^{(7)}(T_{37}, t)| \leq (\widehat{k}_{36})^{(7)} |T'_{37} - T_{37}| e^{-(\widehat{M}_{36})^{(7)}t}$$

$$|(b''_i)^{(7)}(G'_{39}, t) - (b''_i)^{(7)}(G_{39}, t)| < (\widehat{k}_{36})^{(7)} |(G_{39}) - (G'_{39})| e^{-(\widehat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(7)}(T'_{37}, t)$  and  $(a''_i)^{(7)}(T_{37}, t)$ .  $(T'_{37}, t)$  and  $(T_{37}, t)$  are points belonging to the interval  $[(\widehat{k}_{36})^{(7)}, (\widehat{M}_{36})^{(7)}]$ . It is to be noted that  $(a''_i)^{(7)}(T_{37}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\widehat{M}_{36})^{(7)} = 1$  then the function  $(a''_i)^{(7)}(T_{37}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\widehat{M}_{36})^{(7)}, (\widehat{k}_{36})^{(7)}$  :

134

(K)  $(\widehat{M}_{36})^{(7)}, (\widehat{k}_{36})^{(7)}$ , are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

**Definition of**  $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$  : 135

(L) There exists two constants  $(\hat{P}_{36})^{(7)}$  and  $(\hat{Q}_{36})^{(7)}$  which together with  $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)}$  and  $(\hat{B}_{36})^{(7)}$  and the constants  $(a_i)^{(7)}, (a'_i)^{(7)}, (b_i)^{(7)}, (b'_i)^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$ , satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(a_i)^{(7)} + (a'_i)^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}} [(b_i)^{(7)} + (b'_i)^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a'_i)^{(8)}, (a''_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (b''_i)^{(8)} > 0, i, j = 40,41,42 \quad 136$$

The functions  $(a''_i)^{(8)}, (b''_i)^{(8)}$  are positive continuous increasing and bounded

**Definition of**  $(p_i)^{(8)}, (r_i)^{(8)}$ : 137

$$(a''_i)^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \quad 138$$

$$(b''_i)^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b'_i)^{(8)} \leq (\hat{B}_{40})^{(8)} \quad 139$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(8)}(T_{41}, t) = (p_i)^{(8)} \quad 140$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(8)}((G_{43}), t) = (r_i)^{(8)} \quad 141$$

**Definition of**  $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$  :

Where  $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}$  are positive constants and  $i = 40,41,42$

They satisfy Lipschitz condition:

$$|(a''_i)^{(8)}(T'_{41}, t) - (a''_i)^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T'_{41} - T_{41}| e^{-(\hat{M}_{40})^{(8)}t} \quad 142$$

$$|(b''_i)^{(8)}((G'_{43}), t) - (b''_i)^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} |(G'_{43}) - (G_{43})| e^{-(\hat{M}_{40})^{(8)}t} \quad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(8)}(T'_{41}, t)$  and  $(a''_i)^{(8)}(T_{41}, t)$ .  $(T'_{41}, t)$  and  $(T_{41}, t)$  are points belonging to the interval  $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$ . It is to be noted that  $(a''_i)^{(8)}(T_{41}, t)$  is uniformly continuous. In the eventuality of the fact, that if  $(\hat{M}_{40})^{(8)} = 1$  then the function  $(a''_i)^{(8)}(T_{41}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\widehat{M}_{40})^{(8)}, (\widehat{k}_{40})^{(8)}$  :

$(\widehat{M}_{40})^{(8)}, (\widehat{k}_{40})^{(8)}$ , are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} , \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1 \tag{144}$$

**Definition of**  $(\widehat{P}_{40})^{(8)}, (\widehat{Q}_{40})^{(8)}$  :

There exists two constants  $(\widehat{P}_{40})^{(8)}$  and  $(\widehat{Q}_{40})^{(8)}$  which together with  $(\widehat{M}_{40})^{(8)}, (\widehat{k}_{40})^{(8)}, (\widehat{A}_{40})^{(8)}$   $(\widehat{B}_{40})^{(8)}$  and the constants  $(a_i)^{(8)}, (a'_i)^{(8)}, (b_i)^{(8)}, (b'_i)^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40, 41, 42$ , Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}} [ (a_i)^{(8)} + (a'_i)^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)} (\widehat{k}_{40})^{(8)} ] < 1 \tag{145}$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}} [ (b_i)^{(8)} + (b'_i)^{(8)} + (\widehat{B}_{40})^{(8)} + (\widehat{Q}_{40})^{(8)} (\widehat{k}_{40})^{(8)} ] < 1 \tag{146}$$

Where we suppose

$$(a_i)^{(9)}, (a'_i)^{(9)}, (a''_i)^{(9)}, (b_i)^{(9)}, (b'_i)^{(9)}, (b''_i)^{(9)} > 0, i, j = 44, 45, 46 \tag{146}$$

A

The functions  $(a''_i)^{(9)}, (b''_i)^{(9)}$  are positive continuous increasing and bounded.

**Definition of**  $(p_i)^{(9)}, (r_i)^{(9)}$ :

$$(a''_i)^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\widehat{A}_{44})^{(9)}$$

$$(b''_i)^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b'_i)^{(9)} \leq (\widehat{B}_{44})^{(9)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

**Definition of**  $(\widehat{A}_{44})^{(9)}, (\widehat{B}_{44})^{(9)}$  :

Where  $(\widehat{A}_{44})^{(9)}, (\widehat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}$  are positive constants and  $i = 44, 45, 46$

They satisfy Lipschitz condition:

$$|(a''_i)^{(9)}(T'_{45}, t) - (a''_i)^{(9)}(T_{45}, t)| \leq (\widehat{k}_{44})^{(9)} |T'_{45} - T_{45}| e^{-(\widehat{M}_{44})^{(9)}t}$$

$$|(b''_i)^{(9)}((G'_{47}), t) - (b''_i)^{(9)}((G_{47}), t)| < (\widehat{k}_{44})^{(9)} |(G'_{47}) - (G_{47})| e^{-(\widehat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions  $(a''_i)^{(9)}(T'_{45}, t)$  and  $(a''_i)^{(9)}(T_{45}, t)$ .  $(T'_{45}, t)$  and  $(T_{45}, t)$  are points belonging to the interval  $[(\widehat{k}_{44})^{(9)}, (\widehat{M}_{44})^{(9)}]$ . It is to be noted that  $(a''_i)^{(9)}(T_{45}, t)$  is uniformly continuous. In the eventuality of

the fact, that if  $(\widehat{M}_{44})^{(9)} = 1$  then the function  $(a_i'')^{(9)}(T_{45}, t)$ , the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

**Definition of**  $(\widehat{M}_{44})^{(9)}, (\widehat{k}_{44})^{(9)}$  :

$(\widehat{M}_{44})^{(9)}, (\widehat{k}_{44})^{(9)}$ , are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} , \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$$

**Definition of**  $(\widehat{P}_{44})^{(9)}, (\widehat{Q}_{44})^{(9)}$  :

There exists two constants  $(\widehat{P}_{44})^{(9)}$  and  $(\widehat{Q}_{44})^{(9)}$  which together with  $(\widehat{M}_{44})^{(9)}, (\widehat{k}_{44})^{(9)}, (\widehat{A}_{44})^{(9)}$  and  $(\widehat{B}_{44})^{(9)}$  and the constants  $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44, 45, 46$ , satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}} [(a_i)^{(9)} + (a_i')^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}} [(b_i)^{(9)} + (b_i')^{(9)} + (\widehat{B}_{44})^{(9)} + (\widehat{Q}_{44})^{(9)}(\widehat{k}_{44})^{(9)}] < 1$$

**Theorem 1:** if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} , \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} , \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 2 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

**Definition of**  $G_i(0), T_i(0)$

$$G_i(t) \leq (\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} , G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} , T_i(0) = T_i^0 > 0$$

**Theorem 3 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \leq (\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} , G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} , T_i(0) = T_i^0 > 0$$

**Theorem 4 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} , \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 5 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 6 :** if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 7:** if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 8:** if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

A

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Theorem 9:** if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

**Definition of**  $G_i(0), T_i(0)$  :

$$G_i(t) \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

**Proof:** Consider operator  $\mathcal{A}^{(1)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy 154

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)}, \quad 155$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[ (a_{13})^{(1)} G_{14}(s_{(13)}) - \left( (a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[ (a_{14})^{(1)} G_{13}(s_{(13)}) - \left( (a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[ (a_{15})^{(1)} G_{14}(s_{(13)}) - \left( (a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[ (b_{13})^{(1)} T_{14}(s_{(13)}) - \left( (b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[ (b_{14})^{(1)} T_{13}(s_{(13)}) - \left( (b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[ (b_{15})^{(1)} T_{14}(s_{(13)}) - \left( (b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where  $s_{(13)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:** 159

Consider operator  $\mathcal{A}^{(2)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[ (a_{16})^{(2)} G_{17}(s_{(16)}) - \left( (a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[ (a_{17})^{(2)} G_{16}(s_{(16)}) - \left( (a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[ (a_{18})^{(2)} G_{17}(s_{(16)}) - \left( (a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[ (b_{16})^{(2)} T_{17}(s_{(16)}) - \left( (b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[ (b_{17})^{(2)} T_{16}(s_{(16)}) - \left( (b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[ (b_{18})^{(2)} T_{17}(s_{(16)}) - \left( (b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where  $s_{(16)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(3)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$\begin{aligned} G_i(0) &= G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)}, \\ 0 &\leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t} \\ 0 &\leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t} \end{aligned}$$

By

161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[ (a_{20})^{(3)} G_{21}(s_{(20)}) - \left( (a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[ (a_{21})^{(3)} G_{20}(s_{(20)}) - \left( (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[ (a_{22})^{(3)} G_{21}(s_{(20)}) - \left( (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[ (b_{20})^{(3)} T_{21}(s_{(20)}) - \left( (b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[ (b_{21})^{(3)} T_{20}(s_{(20)}) - \left( (b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[ (b_{22})^{(3)} T_{21}(s_{(20)}) - \left( (b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where  $s_{(20)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:** Consider operator  $\mathcal{A}^{(4)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$$

By

162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[ (a_{24})^{(4)} G_{25}(s_{(24)}) - \left( (a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[ (a_{25})^{(4)} G_{24}(s_{(24)}) - \left( (a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[ (a_{26})^{(4)} G_{25}(s_{(24)}) - \left( (a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[ (b_{24})^{(4)} T_{25}(s_{(24)}) - \left( (b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[ (b_{25})^{(4)} T_{24}(s_{(24)}) - \left( (b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[ (b_{26})^{(4)} T_{25}(s_{(24)}) - \left( (b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where  $s_{(24)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:** Consider operator  $\mathcal{A}^{(5)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}$$

By

163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[ (a_{28})^{(5)} G_{29}(s_{(28)}) - \left( (a'_{28})^{(5)} + a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[ (a_{29})^{(5)} G_{28}(s_{(28)}) - \left( (a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[ (a_{30})^{(5)} G_{29}(s_{(28)}) - \left( (a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[ (b_{28})^{(5)} T_{29}(s_{(28)}) - \left( (b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[ (b_{29})^{(5)} T_{28}(s_{(28)}) - \left( (b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[ (b_{30})^{(5)} T_{29}(s_{(28)}) - \left( (b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where  $s_{(28)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(6)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(M_{32})^{(6)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(M_{32})^{(6)}t}$$

By

164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[ (a_{32})^{(6)} G_{33}(s_{(32)}) - \left( (a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[ (a_{33})^{(6)} G_{32}(s_{(32)}) - \left( (a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[ (a_{34})^{(6)} G_{33}(s_{(32)}) - \left( (a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[ (b_{32})^{(6)} T_{33}(s_{(32)}) - \left( (b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[ (b_{33})^{(6)} T_{32}(s_{(32)}) - \left( (b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[ (b_{34})^{(6)} T_{33}(s_{(32)}) - \left( (b'_{34})^{(6)} - (b''_{34})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)}$$

Where  $s_{(32)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(7)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{36})^{(7)}, T_i^0 \leq (\hat{Q}_{36})^{(7)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{36})^{(7)} e^{(M_{36})^{(7)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)}t}$$

By

165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[ (a_{36})^{(7)} G_{37}(s_{(36)}) - \left( (a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[ (a_{37})^{(7)} G_{36}(s_{(36)}) - \left( (a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[ (a_{38})^{(7)} G_{37}(s_{(36)}) - \left( (a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)}) \right) G_{38}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[ (b_{36})^{(7)} T_{37}(s_{(36)}) - \left( (b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{36}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[ (b_{37})^{(7)} T_{36}(s_{(36)}) - \left( (b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{37}(s_{(36)}) \right] ds_{(36)}$$

$$\bar{T}_{38}(t) = T_{38}^0 + \int_0^t \left[ (b_{38})^{(7)} T_{37}(s_{(36)}) - \left( (b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)}) \right) T_{38}(s_{(36)}) \right] ds_{(36)}$$

Where  $s_{(36)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(8)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{40})^{(8)}, T_i^0 \leq (\hat{Q}_{40})^{(8)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}t}$$

By

166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[ (a_{40})^{(8)} G_{41}(s_{(40)}) - \left( (a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[ (a_{41})^{(8)} G_{40}(s_{(40)}) - \left( (a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[ (a_{42})^{(8)} G_{41}(s_{(40)}) - \left( (a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)}) \right) G_{42}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[ (b_{40})^{(8)} T_{41}(s_{(40)}) - \left( (b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)}) \right) T_{40}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[ (b_{41})^{(8)} T_{40}(s_{(40)}) - \left( (b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)}) \right) T_{41}(s_{(40)}) \right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[ (b_{42})^{(8)} T_{41}(s_{(40)}) - \left( (b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)}) \right) T_{42}(s_{(40)}) \right] ds_{(40)}$$

Where  $s_{(40)}$  is the integrand that is integrated over an interval  $(0, t)$

**Proof:**

Consider operator  $\mathcal{A}^{(9)}$  defined on the space of sextuples of continuous functions  $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$  which satisfy

166  
A

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{44})^{(9)}, T_i^0 \leq (\hat{Q}_{44})^{(9)},$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[ (a_{44})^{(9)} G_{45}(s_{(44)}) - \left( (a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)}) \right) G_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[ (a_{45})^{(9)} G_{44}(s_{(44)}) - \left( (a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)}) \right) G_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[ (a_{46})^{(9)} G_{45}(s_{(44)}) - \left( (a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)}) \right) G_{46}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[ (b_{44})^{(9)} T_{45}(s_{(44)}) - \left( (b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)}) \right) T_{44}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[ (b_{45})^{(9)} T_{44}(s_{(44)}) - \left( (b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)}) \right) T_{45}(s_{(44)}) \right] ds_{(44)}$$

$$\bar{T}_{46}(t) = T_{46}^0 + \int_0^t \left[ (b_{46})^{(9)} T_{45}(s_{(44)}) - \left( (b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)}) \right) T_{46}(s_{(44)}) \right] ds_{(44)}$$

Where  $s_{(44)}$  is the integrand that is integrated over an interval  $(0, t)$

The operator  $\mathcal{A}^{(1)}$  maps the space of functions satisfying Equations into itself. Indeed it is obvious that

167

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[ (a_{13})^{(1)} \left( G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}s_{(13)}} \right) \right] ds_{(13)} =$$

$$\left( 1 + (a_{13})^{(1)}t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left( e^{(\hat{M}_{13})^{(1)}t} - 1 \right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0)e^{-(M_{13})^{(1)}t} \leq \frac{(a_{13})^{(1)}}{(M_{13})^{(1)}} \left[ ((\hat{P}_{13})^{(1)} + G_{14}^0)e^{-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}} + (\hat{P}_{13})^{(1)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 1

Analogous inequalities hold also for  $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator  $\mathcal{A}^{(2)}$  maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[ (a_{16})^{(2)} \left( G_{17}^0 + (\hat{P}_{16})^{(2)} e^{(M_{16})^{(2)}s(16)} \right) \right] ds_{(16)} \tag{169}$$

$$= (1 + (a_{16})^{(2)}t)G_{17}^0 + \frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(M_{16})^{(2)}} \left( e^{(M_{16})^{(2)}t} - 1 \right)$$

From which it follows that 170

$$(G_{16}(t) - G_{16}^0)e^{-(M_{16})^{(2)}t} \leq \frac{(a_{16})^{(2)}}{(M_{16})^{(2)}} \left[ ((\hat{P}_{16})^{(2)} + G_{17}^0)e^{-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for  $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator  $\mathcal{A}^{(3)}$  maps the space of functions satisfying Equations into itself. Indeed it is obvious that 171

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[ (a_{20})^{(3)} \left( G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(M_{20})^{(3)}s(20)} \right) \right] ds_{(20)} =$$

$$(1 + (a_{20})^{(3)}t)G_{21}^0 + \frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(M_{20})^{(3)}} \left( e^{(M_{20})^{(3)}t} - 1 \right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^0)e^{-(M_{20})^{(3)}t} \leq \frac{(a_{20})^{(3)}}{(M_{20})^{(3)}} \left[ ((\hat{P}_{20})^{(3)} + G_{21}^0)e^{-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for  $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator  $\mathcal{A}^{(4)}$  maps the space of functions satisfying into itself. Indeed it is obvious that 173

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[ (a_{24})^{(4)} \left( G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(M_{24})^{(4)}s(24)} \right) \right] ds_{(24)} =$$

$$(1 + (a_{24})^{(4)}t)G_{25}^0 + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(M_{24})^{(4)}} \left( e^{(M_{24})^{(4)}t} - 1 \right)$$

From which it follows that 174

$$(G_{24}(t) - G_{24}^0)e^{-(M_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\widehat{M}_{24})^{(4)}} \left[ ((\widehat{P}_{24})^{(4)} + G_{25}^0)e^{\left(-\frac{(\widehat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\widehat{P}_{24})^{(4)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 4

The operator  $\mathcal{A}^{(5)}$  maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[ (a_{28})^{(5)} \left( G_{29}^0 + (\widehat{P}_{28})^{(5)} e^{(M_{28})^{(5)}s_{(28)}} \right) \right] ds_{(28)} =$$

$$(1 + (a_{28})^{(5)}t)G_{29}^0 + \frac{(a_{28})^{(5)}(\widehat{P}_{28})^{(5)}}{(\widehat{M}_{28})^{(5)}} \left( e^{(M_{28})^{(5)}t} - 1 \right)$$

From which it follows that

175

$$(G_{28}(t) - G_{28}^0)e^{-(M_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\widehat{M}_{28})^{(5)}} \left[ ((\widehat{P}_{28})^{(5)} + G_{29}^0)e^{\left(-\frac{(\widehat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\widehat{P}_{28})^{(5)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 5

The operator  $\mathcal{A}^{(6)}$  maps the space of functions satisfying Equations into itself. Indeed it is obvious that

176

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[ (a_{32})^{(6)} \left( G_{33}^0 + (\widehat{P}_{32})^{(6)} e^{(M_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$(1 + (a_{32})^{(6)}t)G_{33}^0 + \frac{(a_{32})^{(6)}(\widehat{P}_{32})^{(6)}}{(\widehat{M}_{32})^{(6)}} \left( e^{(M_{32})^{(6)}t} - 1 \right)$$

From which it follows that

177

$$(G_{32}(t) - G_{32}^0)e^{-(M_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\widehat{M}_{32})^{(6)}} \left[ ((\widehat{P}_{32})^{(6)} + G_{33}^0)e^{\left(-\frac{(\widehat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\widehat{P}_{32})^{(6)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 6

Analogous inequalities hold also for  $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

(b) The operator  $\mathcal{A}^{(7)}$  maps the space of functions satisfying Equations into itself. Indeed it is obvious that

178

$$G_{36}(t) \leq G_{36}^0 + \int_0^t \left[ (a_{36})^{(7)} \left( G_{37}^0 + (\widehat{P}_{36})^{(7)} e^{(M_{36})^{(7)}s_{(36)}} \right) \right] ds_{(36)} =$$

$$(1 + (a_{36})^{(7)}t)G_{37}^0 + \frac{(a_{36})^{(7)}(\widehat{P}_{36})^{(7)}}{(\widehat{M}_{36})^{(7)}} \left( e^{(M_{36})^{(7)}t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0)e^{-(\hat{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[ ((\hat{P}_{36})^{(7)} + G_{37}^0)e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 7

The operator  $\mathcal{A}^{(8)}$  maps the space of functions satisfying Equations into itself. Indeed it is obvious that

$$G_{40}(t) \leq G_{40}^0 + \int_0^t \left[ (a_{40})^{(8)} \left( G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)}s_{(40)}} \right) \right] ds_{(40)} = \tag{180}$$

$$\left( 1 + (a_{40})^{(8)}t \right) G_{41}^0 + \frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left( e^{(\hat{M}_{40})^{(8)}t} - 1 \right)$$

From which it follows that

$$(G_{40}(t) - G_{40}^0)e^{-(\hat{M}_{40})^{(8)}t} \leq \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[ ((\hat{P}_{40})^{(8)} + G_{41}^0)e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 8

Analogous inequalities hold also for  $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator  $\mathcal{A}^{(9)}$  maps the space of functions satisfying 34,35,36 into itself. Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[ (a_{44})^{(9)} \left( G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)}s_{(44)}} \right) \right] ds_{(44)} =$$

$$\left( 1 + (a_{44})^{(9)}t \right) G_{45}^0 + \frac{(a_{44})^{(9)}(\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left( e^{(\hat{M}_{44})^{(9)}t} - 1 \right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0)e^{-(\hat{M}_{44})^{(9)}t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[ ((\hat{P}_{44})^{(9)} + G_{45}^0)e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)} \right]$$

$(G_i^0)$  is as defined in the statement of theorem 9

Analogous inequalities hold also for  $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take  $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$  and to choose  $\tag{182}$

$(\hat{P}_{13})^{(1)}$  and  $(\hat{Q}_{13})^{(1)}$  large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[ (\hat{P}_{13})^{(1)} + ((\hat{P}_{13})^{(1)} + G_j^0)e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{13})^{(1)} \tag{183}$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}} \left[ ((\widehat{Q}_{13})^{(1)} + T_j^0) e^{-\left(\frac{(\widehat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\widehat{Q}_{13})^{(1)} \right] \leq (\widehat{Q}_{13})^{(1)} \tag{184}$$

In order that the operator  $\mathcal{A}^{(1)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(1)}$  is a contraction with respect to the metric 185

$$d\left((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)})\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{13})^{(1)}t} \right\}$$

Indeed if we denote

**Definition of  $\tilde{G}, \tilde{T}$  :**  $(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$

It results

$$\begin{aligned} |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\ &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} + \\ &(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} + \\ &G_{13}^{(2)} |(a''_{13})^{(1)} (T_{14}^{(1)}, s_{(13)}) - (a''_{13})^{(1)} (T_{14}^{(2)}, s_{(13)})| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)} \end{aligned}$$

Where  $s_{(13)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)}t} &\leq \frac{1}{(\widehat{M}_{13})^{(1)}} ((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} \\ &+ (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)}) d\left((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)})\right) \end{aligned} \tag{186}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 1:** The fact that we supposed  $(a''_{13})^{(1)}$  and  $(b''_{13})^{(1)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$  and  $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(1)}$  and  $(b''_i)^{(1)}$ ,  $i = 13, 14, 15$  depend only on  $T_{14}$  and respectively on

$G$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 2:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(1)} - (a_i'')^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(1)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{13})^{(1)})_1, ((\widehat{M}_{13})^{(1)})_2$  and  $((\widehat{M}_{13})^{(1)})_3$  : 187

**Remark 3:** if  $G_{13}$  is bounded, the same property holds also for  $G_{14}$  and  $G_{15}$ . indeed if

$G_{13} < ((\widehat{M}_{13})^{(1)})_1$  it follows  $\frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a_{14}')^{(1)}G_{14}$  and by integrating

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14}')^{(1)}((\widehat{M}_{13})^{(1)})_1 / (a_{14}')^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15}')^{(1)}((\widehat{M}_{13})^{(1)})_2 / (a_{15}')^{(1)}$$

If  $G_{14}$  or  $G_{15}$  is bounded, the same property follows for  $G_{13}$ ,  $G_{15}$  and  $G_{13}$ ,  $G_{14}$  respectively.

**Remark 4:** If  $G_{13}$  is bounded, from below, the same property holds for  $G_{14}$  and  $G_{15}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{14}$  is bounded from below. 188

**Remark 5:** If  $T_{13}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(1)}(G(t), t)) = (b_{14}')^{(1)}$  then  $T_{14} \rightarrow \infty$ . 189

**Definition of**  $(m)^{(1)}$  and  $\varepsilon_1$  :

Indeed let  $t_1$  be so that for  $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then  $\frac{dT_{14}}{dt} \geq (a_{14}')^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$  which leads to

$$T_{14} \geq \left( \frac{(a_{14}')^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_1 t} = \frac{1}{2} \text{ it results}$$

$$T_{14} \geq \left( \frac{(a_{14}')^{(1)}(m)^{(1)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_1} \text{ By taking now } \varepsilon_1 \text{ sufficiently small one sees that } T_{14} \text{ is unbounded.}$$

The same property holds for  $T_{15}$  if  $\lim_{t \rightarrow \infty} ((b_{15}'')^{(1)}(G(t), t)) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take  $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$  and to choose 190

$(\widehat{P}_{16})^{(2)}$  and  $(\widehat{Q}_{16})^{(2)}$  large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[ (\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{16})^{(2)} \tag{191}$$

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[ ((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)} \tag{192}$$

In order that the operator  $\mathcal{A}^{(2)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself 193

The operator  $\mathcal{A}^{(2)}$  is a contraction with respect to the metric 194

$$d \left( ((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote 195

**Definition of  $\tilde{G}_{19}, \tilde{T}_{19}$  :**  $(\tilde{G}_{19}, \tilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\begin{aligned} |\tilde{G}_{16}^{(1)} - \tilde{G}_{16}^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} | (a''_{16})^{(2)} (T_{17}^{(1)}, s_{(16)}) - (a''_{16})^{(2)} (T_{17}^{(2)}, s_{(16)}) | e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} \} ds_{(16)} \end{aligned}$$

Where  $s_{(16)}$  represents integrand that is integrated over the interval  $[0, t]$  197

From the hypotheses it follows

$$\begin{aligned} &|(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\bar{M}_{16})^{(2)}t} \\ &\leq \frac{1}{(\bar{M}_{16})^{(2)}} \left( (a_{16})^{(2)} + (a'_{16})^{(2)} + (\hat{A}_{16})^{(2)} \right) \\ &+ (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)} d \left( ((G_{19})^{(1)}, (T_{19})^{(1)}); (G_{19})^{(2)}, (T_{19})^{(2)} \right) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows 198

**Remark 6:** The fact that we supposed  $(a''_{16})^{(2)}$  and  $(b''_{16})^{(2)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 199

$(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$  and  $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a_i'')^{(2)}$  and  $(b_i'')^{(2)}$ ,  $i = 16, 17, 18$  depend only on  $T_{17}$  and respectively on  $(G_{19})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 7:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(2)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{16})^{(2)})_1$ ,  $((\widehat{M}_{16})^{(2)})_2$  and  $((\widehat{M}_{16})^{(2)})_3$  : 201

**Remark 8:** if  $G_{16}$  is bounded, the same property have also  $G_{17}$  and  $G_{18}$ . indeed if

$G_{16} < ((\widehat{M}_{16})^{(2)})_1$  it follows  $\frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a_{17}')^{(2)}G_{17}$  and by integrating

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)}((\widehat{M}_{16})^{(2)})_1 / (a_{17}')^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)}((\widehat{M}_{16})^{(2)})_2 / (a_{18}')^{(2)}$$

If  $G_{17}$  or  $G_{18}$  is bounded, the same property follows for  $G_{16}$ ,  $G_{18}$  and  $G_{16}$ ,  $G_{17}$  respectively.

**Remark 9:** If  $G_{16}$  is bounded, from below, the same property holds for  $G_{17}$  and  $G_{18}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{17}$  is bounded from below. 202

**Remark 10:** If  $T_{16}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$  then  $T_{17} \rightarrow \infty$ . 203

**Definition of**  $(m)^{(2)}$  and  $\varepsilon_2$  :

Indeed let  $t_2$  be so that for  $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then  $\frac{dT_{17}}{dt} \geq (a_{17}')^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$  which leads to 204

$$T_{17} \geq \left(\frac{(a_{17}')^{(2)}(m)^{(2)}}{\varepsilon_2}\right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$T_{17} \geq \left(\frac{(a_{17}')^{(2)}(m)^{(2)}}{2}\right)$ ,  $t = \log \frac{2}{\varepsilon_2}$  By taking now  $\varepsilon_2$  sufficiently small one sees that  $T_{17}$  is unbounded. 205

The same property holds for  $T_{18}$  if  $\lim_{t \rightarrow \infty} (b_{18}'')^{(2)}((G_{19})(t), t) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take  $\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1$  and to choose 207

$(\hat{P}_{20})^{(3)}$  and  $(\hat{Q}_{20})^{(3)}$  large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[ (\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[ ((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator  $\mathcal{A}^{(3)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself 210

The operator  $\mathcal{A}^{(3)}$  is a contraction with respect to the metric 211

$$d \left( ((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \}$$

Indeed if we denote 212

**Definition of  $\widetilde{G}_{23}, \widetilde{T}_{23}$  :**  $((\widetilde{G}_{23}), (\widetilde{T}_{23})) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$\begin{aligned} |\widetilde{G}_{20}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} | (a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)}) | e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned}$$

Where  $s_{(20)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G_{23}^{(1)} - G_{23}^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \frac{1}{(\bar{M}_{20})^{(3)}} \left( (a_{20})^{(3)} + (a'_{20})^{(3)} + (\hat{A}_{20})^{(3)} \right) \\ &+ (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)} d \left( ((G_{23})^{(1)}, (T_{23})^{(1)}); (G_{23})^{(2)}, (T_{23})^{(2)} \right) \end{aligned} \quad 214$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 11:** The fact that we supposed  $(a''_{20})^{(3)}$  and  $(b''_{20})^{(3)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate a condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$  and  $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$  respectively of  $\mathbb{R}_+$ . 215

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(3)}$  and  $(b''_i)^{(3)}$ ,  $i = 20, 21, 22$  depend only on  $T_{21}$  and respectively on  $(G_{23})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 12:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(3)} t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{20})^{(3)})_1$ ,  $((\widehat{M}_{20})^{(3)})_2$  and  $((\widehat{M}_{20})^{(3)})_3$  : 217

**Remark 13:** if  $G_{20}$  is bounded, the same property have also  $G_{21}$  and  $G_{22}$ . indeed if

$G_{20} < ((\widehat{M}_{20})^{(3)})_1$  it follows  $\frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a'_{21})^{(3)} G_{21}$  and by integrating

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21})^{(3)} ((\widehat{M}_{20})^{(3)})_1 / (a'_{21})^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22})^{(3)} ((\widehat{M}_{20})^{(3)})_2 / (a'_{22})^{(3)}$$

If  $G_{21}$  or  $G_{22}$  is bounded, the same property follows for  $G_{20}$ ,  $G_{22}$  and  $G_{20}$ ,  $G_{21}$  respectively.

**Remark 14:** If  $G_{20}$  is bounded, from below, the same property holds for  $G_{21}$  and  $G_{22}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{21}$  is bounded from below. 218

**Remark 15:** If  $T_{20}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b'_i)^{(3)} ((G_{23})(t), t)) = (b'_{21})^{(3)}$  then  $T_{21} \rightarrow \infty$ . 219

**Definition of**  $(m)^{(3)}$  and  $\varepsilon_3$  :

Indeed let  $t_3$  be so that for  $t > t_3$

$$(b_{21})^{(3)} - (b''_i)^{(3)} ((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then  $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)} (m)^{(3)} - \varepsilon_3 T_{21}$  which leads to 220

$$T_{21} \geq \left( \frac{(a_{21})^{(3)} (m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$$T_{21} \geq \left( \frac{(a_{21})^{(3)} (m)^{(3)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_3} \text{ By taking now } \varepsilon_3 \text{ sufficiently small one sees that } T_{21} \text{ is unbounded.}$$

The same property holds for  $T_{22}$  if  $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)} ((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take  $\frac{(a_i)^{(4)}}{(\overline{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\overline{M}_{24})^{(4)}} < 1$  and to choose 221

$(\widehat{P}_{24})^{(4)}$  and  $(\widehat{Q}_{24})^{(4)}$  large to have

$$\frac{(a_i)^{(4)}}{(\overline{M}_{24})^{(4)}} \left[ (\widehat{P}_{24})^{(4)} + ((\widehat{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{((\widehat{P}_{24})^{(4)} + G_j^0)}{G_j^0}\right)} \right] \leq (\widehat{P}_{24})^{(4)} \tag{222}$$

$$\frac{(b_i)^{(4)}}{(\overline{M}_{24})^{(4)}} \left[ ((\widehat{Q}_{24})^{(4)} + T_j^0) e^{-\left(\frac{((\widehat{Q}_{24})^{(4)} + T_j^0)}{T_j^0}\right)} + (\widehat{Q}_{24})^{(4)} \right] \leq (\widehat{Q}_{24})^{(4)} \tag{223}$$

In order that the operator  $\mathcal{A}^{(4)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself 224

The operator  $\mathcal{A}^{(4)}$  is a contraction with respect to the metric 225

$$d \left( ((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\overline{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\overline{M}_{24})^{(4)}t} \}$$

Indeed if we denote

**Definition of  $(\widetilde{G}_{27}), (\widetilde{T}_{27})$ :**  $((\widetilde{G}_{27}), (\widetilde{T}_{27})) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\widetilde{G}_{24}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\overline{M}_{24})^{(4)}s_{(24)}} e^{(\overline{M}_{24})^{(4)}s_{(24)}} ds_{(24)} + \\ &\int_0^t \{ (a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\overline{M}_{24})^{(4)}s_{(24)}} e^{-(\overline{M}_{24})^{(4)}s_{(24)}} + \\ & (a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\overline{M}_{24})^{(4)}s_{(24)}} e^{(\overline{M}_{24})^{(4)}s_{(24)}} + \\ & G_{24}^{(2)} | (a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)}) | e^{-(\overline{M}_{24})^{(4)}s_{(24)}} e^{(\overline{M}_{24})^{(4)}s_{(24)}} \} ds_{(24)} \end{aligned}$$

Where  $s_{(24)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses on Equations it follows

$$\begin{aligned}
 & |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\widehat{M}_{24})^{(4)}t} & 226 \\
 & \leq \frac{1}{(\widehat{M}_{24})^{(4)}} \left( (a_{24})^{(4)} + (a'_{24})^{(4)} + (\widehat{A}_{24})^{(4)} \right. \\
 & \quad \left. + (\widehat{P}_{24})^{(4)} (\widehat{k}_{24})^{(4)} \right) d \left( ((G_{27})^{(1)}, (T_{27})^{(1)}); (G_{27})^{(2)}, (T_{27})^{(2)} \right)
 \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 16:** The fact that we supposed  $(a''_{24})^{(4)}$  and  $(b''_{24})^{(4)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$  and  $(\widehat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t}$  respectively of  $\mathbb{R}_+$ . 227

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(4)}$  and  $(b''_i)^{(4)}$ ,  $i = 24, 25, 26$  depend only on  $T_{25}$  and respectively on  $(G_{27})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 17:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  228

it results

$$G_i(t) \geq G_i^0 e^{\left[ - \int_0^t \{ (a'_i)^{(4)} - (a''_i)^{(4)}(T_{25}(s_{(24)}), s_{(24)}) \} ds_{(24)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(4)}t} > 0 \text{ for } t > 0$$

**Definition of**  $(\widehat{M}_{24})^{(4)}_1, (\widehat{M}_{24})^{(4)}_2$  and  $(\widehat{M}_{24})^{(4)}_3$  : 229

**Remark 18:** if  $G_{24}$  is bounded, the same property have also  $G_{25}$  and  $G_{26}$ . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$  it follows  $\frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25}$  and by integrating

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way, one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If  $G_{25}$  or  $G_{26}$  is bounded, the same property follows for  $G_{24}$ ,  $G_{26}$  and  $G_{24}$ ,  $G_{25}$  respectively.

**Remark 19:** If  $G_{24}$  is bounded, from below, the same property holds for  $G_{25}$  and  $G_{26}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{25}$  is bounded from below. 230

**Remark 20:** If  $T_{24}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$  then  $T_{25} \rightarrow \infty$ . 231

**Definition of**  $(m)^{(4)}$  and  $\varepsilon_4$  :

Indeed let  $t_4$  be so that for  $t > t_4$

$$(b_{25})^{(4)} - (b''_i)^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then  $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$  which leads to 232

$$T_{25} \geq \left( \frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$$

If we take  $t$  such that  $e^{-\varepsilon_4 t} = \frac{1}{2}$  it results

$$T_{25} \geq \left( \frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_4}$$

By taking now  $\varepsilon_4$  sufficiently small one sees that  $T_{25}$  is unbounded.

The same property holds for  $T_{26}$  if  $\lim_{t \rightarrow \infty} (G_{27})''(t, t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for  $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take  $\frac{(a_i)^{(5)}}{(\overline{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\overline{M}_{28})^{(5)}} < 1$  and to choose 233

$(\widehat{P}_{28})^{(5)}$  and  $(\widehat{Q}_{28})^{(5)}$  large to have

$$\frac{(a_i)^{(5)}}{(\overline{M}_{28})^{(5)}} \left[ (\widehat{P}_{28})^{(5)} + ((\widehat{P}_{28})^{(5)} + G_j^0) e^{-\left( \frac{(\widehat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\overline{M}_{28})^{(5)}} \left[ ((\widehat{Q}_{28})^{(5)} + T_j^0) e^{-\left( \frac{(\widehat{Q}_{28})^{(5)} + T_j^0}{T_j^0} \right)} + (\widehat{Q}_{28})^{(5)} \right] \leq (\widehat{Q}_{28})^{(5)} \quad 235$$

In order that the operator  $\mathcal{A}^{(5)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(5)}$  is a contraction with respect to the metric 236

$$d \left( ((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\overline{M}_{28})^{(5)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\overline{M}_{28})^{(5)}t} \right\}$$

Indeed if we denote

**Definition of**  $(\widetilde{G}_{31}), (\widetilde{T}_{31})$ :  $((\widetilde{G}_{31}), (\widetilde{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$|\widetilde{G}_{28}^{(1)} - \widetilde{G}_{28}^{(2)}| \leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\overline{M}_{28})^{(5)}s(28)} e^{(\overline{M}_{28})^{(5)}s(28)} ds(28) +$$

$$\int_0^t \{ (a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\overline{M}_{28})^{(5)}s(28)} e^{-(\overline{M}_{28})^{(5)}s(28)} +$$

$$(a''_{28})^{(5)}(T_{29}^{(1)}, s(28)) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\overline{M}_{28})^{(5)}s(28)} e^{(\overline{M}_{28})^{(5)}s(28)} +$$

$$G_{28}^{(2)} |(a''_{28})^{(5)}(T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)}(T_{29}^{(2)}, s_{(28)})| e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} e^{(\widehat{M}_{28})^{(5)}s_{(28)}} ds_{(28)}$$

Where  $s_{(28)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses on it follows

$$\begin{aligned} & |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\widehat{M}_{28})^{(5)}t} & 237 \\ & \leq \frac{1}{(\widehat{M}_{28})^{(5)}} ((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} \\ & + (\widehat{P}_{28})^{(5)} (\widehat{k}_{28})^{(5)}) d((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)}) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 21:** The fact that we supposed  $(a''_{28})^{(5)}$  and  $(b''_{28})^{(5)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$  and  $(\widehat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t}$  respectively of  $\mathbb{R}_+$ . 238

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(5)}$  and  $(b''_i)^{(5)}$ ,  $i = 28, 29, 30$  depend only on  $T_{29}$  and respectively on  $(G_{31})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 22:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  239

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(5)} - (a''_i)^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(5)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$  and  $((\widehat{M}_{28})^{(5)})_3$  : 240

**Remark 23:** if  $G_{28}$  is bounded, the same property have also  $G_{29}$  and  $G_{30}$ . indeed if

$G_{28} < ((\widehat{M}_{28})^{(5)})$  it follows  $\frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)}G_{29}$  and by integrating

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way, one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If  $G_{29}$  or  $G_{30}$  is bounded, the same property follows for  $G_{28}$ ,  $G_{30}$  and  $G_{28}$ ,  $G_{29}$  respectively.

**Remark 24:** If  $G_{28}$  is bounded, from below, the same property holds for  $G_{29}$  and  $G_{30}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{29}$  is bounded from below. 241

**Remark 25:** If  $T_{28}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$  then  $T_{29} \rightarrow \infty$ . 242

**Definition of**  $(m)^{(5)}$  and  $\varepsilon_5$  :

Indeed let  $t_5$  be so that for  $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$$

Then  $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$  which leads to 243

$$T_{29} \geq \left( \frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5} \right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$$T_{29} \geq \left( \frac{(a_{29})^{(5)}(m)^{(5)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_5} \text{ By taking now } \varepsilon_5 \text{ sufficiently small one sees that } T_{29} \text{ is unbounded.}$$

The same property holds for  $T_{30}$  if  $\lim_{t \rightarrow \infty} (b_{30}'')^{(5)}((G_{31})(t), t) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for  $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take  $\frac{(a_i)^{(6)}}{(\bar{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\bar{M}_{32})^{(6)}} < 1$  and to choose 244

$(\hat{P}_{32})^{(6)}$  and  $(\hat{Q}_{32})^{(6)}$  large to have

$$\frac{(a_i)^{(6)}}{(\bar{M}_{32})^{(6)}} \left[ (\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left( \frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\bar{M}_{32})^{(6)}} \left[ ((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left( \frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator  $\mathcal{A}^{(6)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(6)}$  is a contraction with respect to the metric 247

$$d \left( ((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{32})^{(6)} t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{32})^{(6)} t} \right\}$$

Indeed if we denote

**Definition of**  $(\widetilde{G}_{35}), (\widetilde{T}_{35})$  :  $((\widetilde{G}_{35}), (\widetilde{T}_{35})) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$

It results

$$|\widetilde{G}_{32}^{(1)} - \widetilde{G}_i^{(2)}| \leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\bar{M}_{32})^{(6)} s_{(32)}} e^{(\bar{M}_{32})^{(6)} s_{(32)}} ds_{(32)} +$$

$$\int_0^t \{(a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} + (a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}} + G_{32}^{(2)} |(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\widehat{M}_{32})^{(6)} s_{(32)}} e^{(\widehat{M}_{32})^{(6)} s_{(32)}}\} ds_{(32)}$$

Where  $s_{(32)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses it follows

$$\begin{aligned} & |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\widehat{M}_{32})^{(6)} t} \\ & \leq \frac{1}{(\widehat{M}_{32})^{(6)}} ((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)}) \\ & + (\widehat{P}_{32})^{(6)} (\widehat{k}_{32})^{(6)} d \left( ((G_{35})^{(1)}, (T_{35})^{(1)}); (G_{35})^{(2)}, (T_{35})^{(2)}) \right) \end{aligned} \tag{248}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 26:** The fact that we supposed  $(a''_{32})^{(6)}$  and  $(b''_{32})^{(6)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$  and  $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)} t}$  respectively of  $\mathbb{R}_+$ . 249

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(6)}$  and  $(b''_i)^{(6)}$ ,  $i = 32, 33, 34$  depend only on  $T_{33}$  and respectively on  $(G_{35})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 27:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  250

it results

$$G_i(t) \geq G_i^0 e^{[-\int_0^t \{(a'_i)^{(6)} - (a''_i)^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(6)} t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$  and  $((\widehat{M}_{32})^{(6)})_3$  : 251

**Remark 28:** if  $G_{32}$  is bounded, the same property have also  $G_{33}$  and  $G_{34}$ . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$  it follows  $\frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)} G_{33}$  and by integrating

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)} ((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way, one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)} ((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If  $G_{33}$  or  $G_{34}$  is bounded, the same property follows for  $G_{32}$ ,  $G_{34}$  and  $G_{32}$ ,  $G_{33}$  respectively.

**Remark 29:** If  $G_{32}$  is bounded, from below, the same property holds for  $G_{33}$  and  $G_{34}$ . The proof is analogous with the preceding one. An analogous property is true if  $G_{33}$  is bounded from below. 252

**Remark 30:** If  $T_{32}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t)) = (b_{33}')^{(6)}$  then  $T_{33} \rightarrow \infty$ . 253

**Definition of**  $(m)^{(6)}$  and  $\varepsilon_6$  :

Indeed let  $t_6$  be so that for  $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then  $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$  which leads to 254

$$T_{33} \geq \left( \frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$$

If we take  $t$  such that  $e^{-\varepsilon_6 t} = \frac{1}{2}$  it results

$$T_{33} \geq \left( \frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_6}$$

By taking now  $\varepsilon_6$  sufficiently small one sees that  $T_{33}$  is unbounded.

The same property holds for  $T_{34}$  if  $\lim_{t \rightarrow \infty} (b_{34}'')^{(6)}((G_{35})(t), t(t), t) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for  $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$  255

It is now sufficient to take  $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$  and to choose  $(\widehat{P}_{36})^{(7)}$  and  $(\widehat{Q}_{36})^{(7)}$  large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[ (\widehat{P}_{36})^{(7)} + ((\widehat{P}_{36})^{(7)} + G_j^0) e^{-\left( \frac{((\widehat{P}_{36})^{(7)} + G_j^0)}{G_j^0} \right)} \right] \leq (\widehat{P}_{36})^{(7)}$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} \left[ ((\widehat{Q}_{36})^{(7)} + T_j^0) e^{-\left( \frac{((\widehat{Q}_{36})^{(7)} + T_j^0)}{T_j^0} \right)} + (\widehat{Q}_{36})^{(7)} \right] \leq (\widehat{Q}_{36})^{(7)}$$

In order that the operator  $\mathcal{A}^{(7)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(7)}$  is a contraction with respect to the metric 258

$$d \left( ((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)}) \right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\widehat{M}_{36})^{(7)}t} \right\}$$

Indeed if we denote

**Definition of**  $(\widehat{G}_{39}), (\widehat{T}_{39}) : ((\widehat{G}_{39}), (\widehat{T}_{39})) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$

It results

$$\begin{aligned}
 |\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| \leq & \int_0^t (a_{36})^{(7)} |G_{37}^{(1)} - G_{37}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} ds_{(36)} + \\
 & \int_0^t \{ (a'_{36})^{(7)} |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{-(\bar{M}_{36})^{(7)}s_{(36)}} + \\
 & (a''_{36})^{(7)}(T_{37}^{(1)}, s_{(36)}) |G_{36}^{(1)} - G_{36}^{(2)}| e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} + \\
 & G_{36}^{(2)} | (a''_{36})^{(7)}(T_{37}^{(1)}, s_{(36)}) - (a''_{36})^{(7)}(T_{37}^{(2)}, s_{(36)}) | e^{-(\bar{M}_{36})^{(7)}s_{(36)}} e^{(\bar{M}_{36})^{(7)}s_{(36)}} \} ds_{(36)}
 \end{aligned}$$

Where  $s_{(36)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses on it follows

$$\begin{aligned}
 |(G_{39})^{(1)} - (G_{39})^{(2)}| e^{-(\bar{M}_{36})^{(7)}t} & \tag{259} \\
 \leq \frac{1}{(\bar{M}_{36})^{(7)}} & ((a_{36})^{(7)} + (a'_{36})^{(7)} + (\bar{A}_{36})^{(7)} \\
 + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}) & d((G_{39})^{(1)}, (T_{39})^{(1)}; (G_{39})^{(2)}, (T_{39})^{(2)})
 \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 31:** The fact that we supposed  $(a''_{36})^{(7)}$  and  $(b''_{36})^{(7)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$  and  $(\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)}t}$  respectively of  $\mathbb{R}_+$ . 260

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(7)}$  and  $(b''_i)^{(7)}$ ,  $i = 36, 37, 38$  depend only on  $T_{37}$  and respectively on  $(G_{39})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 32:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  261

it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(7)} - (a''_i)^{(7)}(T_{37}(s_{(36)}), s_{(36)}) \} ds_{(36)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(7)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\bar{M}_{36})^{(7)})_1, ((\bar{M}_{36})^{(7)})_2$  and  $((\bar{M}_{36})^{(7)})_3$  : 262

**Remark 33:** if  $G_{36}$  is bounded, the same property have also  $G_{37}$  and  $G_{38}$ . indeed if

$G_{36} < (\bar{M}_{36})^{(7)}$  it follows  $\frac{dG_{37}}{dt} \leq ((\bar{M}_{36})^{(7)})_1 - (a'_{37})^{(7)}G_{37}$  and by integrating

$$G_{37} \leq ((\bar{M}_{36})^{(7)})_2 = G_{37}^0 + 2(a_{37})^{(7)}((\bar{M}_{36})^{(7)})_1 / (a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq ((\widehat{M}_{36})^{(7)})_3 = G_{38}^0 + 2(a_{38})^{(7)}((\widehat{M}_{36})^{(7)})_2 / (a'_{38})^{(7)}$$

If  $G_{37}$  or  $G_{38}$  is bounded, the same property follows for  $G_{36}$  ,  $G_{38}$  and  $G_{36}$  ,  $G_{37}$  respectively.

**Remark 34:** If  $G_{36}$  is bounded, from below, the same property holds for  $G_{37}$  and  $G_{38}$  . The proof is analogous with the preceding one. An analogous property is true if  $G_{37}$  is bounded from below. 263

**Remark 35:** If  $T_{36}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b_i'')^{(7)} ((G_{39})(t), t)) = (b'_{37})^{(7)}$  then  $T_{37} \rightarrow \infty$ . 264

**Definition of**  $(m)^{(7)}$  and  $\varepsilon_7$  :

Indeed let  $t_7$  be so that for  $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)} ((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then  $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$  which leads to 265

$$T_{37} \geq \left( \frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7} \right) (1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_7 t} = \frac{1}{2} \text{ it results}$$

$$T_{37} \geq \left( \frac{(a_{37})^{(7)}(m)^{(7)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_7} \text{ By taking now } \varepsilon_7 \text{ sufficiently small one sees that } T_{37} \text{ is unbounded.}$$

The same property holds for  $T_{38}$  if  $\lim_{t \rightarrow \infty} ((b_{38}'')^{(7)} ((G_{39})(t), t)) = (b'_{38})^{(7)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take  $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}, \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$  and to choose 266  
 $(\widehat{P}_{40})^{(8)}$  and  $(\widehat{Q}_{40})^{(8)}$  large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[ (\widehat{P}_{40})^{(8)} + ((\widehat{P}_{40})^{(8)} + G_j^0) e^{-\left( \frac{((\widehat{P}_{40})^{(8)} + G_j^0)}{G_j^0} \right)} \right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} \left[ ((\widehat{Q}_{40})^{(8)} + T_j^0) e^{-\left( \frac{((\widehat{Q}_{40})^{(8)} + T_j^0)}{T_j^0} \right)} + (\widehat{Q}_{40})^{(8)} \right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator  $\mathcal{A}^{(8)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying Equations into itself

The operator  $\mathcal{A}^{(8)}$  is a contraction with respect to the metric

$$d \left( ((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)}) \right) = \quad 269$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{40})^{(8)}t} \}$$

Indeed if we denote 270

**Definition of**  $(\widetilde{G}_{43}), (\widetilde{T}_{43})$  :  $((\widetilde{G}_{43}), (\widetilde{T}_{43})) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$\begin{aligned} |\tilde{G}_{40}^{(1)} - \tilde{G}_{40}^{(2)}| &\leq \int_0^t (a_{40})^{(8)} |G_{41}^{(1)} - G_{41}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} ds_{(40)} + \\ &\int_0^t \{ (a'_{40})^{(8)} |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{-(\bar{M}_{40})^{(8)}s_{(40)}} + \\ (a''_{40})^{(8)}(T_{41}^{(1)}, s_{(40)}) |G_{40}^{(1)} - G_{40}^{(2)}| e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} + \\ G_{40}^{(2)} | (a''_{40})^{(8)}(T_{41}^{(1)}, s_{(40)}) - (a''_{40})^{(8)}(T_{41}^{(2)}, s_{(40)}) | e^{-(\bar{M}_{40})^{(8)}s_{(40)}} e^{(\bar{M}_{40})^{(8)}s_{(40)}} \} ds_{(40)} \end{aligned}$$

Where  $s_{(40)}$  represents integrand that is integrated over the interval  $[0, t]$  272

From the hypotheses it follows

$$\begin{aligned} |(G_{43})^{(1)} - (G_{43})^{(2)}| e^{-(\bar{M}_{40})^{(8)}t} & \leq \frac{1}{(\bar{M}_{40})^{(8)}} ((a_{40})^{(8)} + (a'_{40})^{(8)} + (\tilde{A}_{40})^{(8)}) \\ & + (\tilde{P}_{40})^{(8)} (\tilde{k}_{40})^{(8)} d \left( ((G_{43})^{(1)}, (T_{43})^{(1)}); (G_{43})^{(2)}, (T_{43})^{(2)} \right) \end{aligned}$$
273

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis the result follows

**Remark 36:** The fact that we supposed  $(a''_{40})^{(8)}$  and  $(b''_{40})^{(8)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$  and  $(\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)}t}$  respectively of  $\mathbb{R}_+$ . 274

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a''_i)^{(8)}$  and  $(b''_i)^{(8)}$ ,  $i = 40, 41, 42$  depend only on  $T_{41}$  and respectively on  $(G_{43})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 37** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$  275

it results

$$G_i(t) \geq G_i^0 e^{\left[ - \int_0^t \{ (a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)}) \} ds_{(40)} \right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(8)}t} > 0 \text{ for } t > 0$$

**Definition of**  $(\widehat{M}_{40})^{(8)}_1, (\widehat{M}_{40})^{(8)}_2$  and  $(\widehat{M}_{40})^{(8)}_3$  : 276

**Remark 38:** if  $G_{40}$  is bounded, the same property have also  $G_{41}$  and  $G_{42}$  . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$  it follows  $\frac{dG_{41}}{dt} \leq ((\widehat{M}_{40})^{(8)})_1 - (a'_{41})^{(8)}G_{41}$  and by integrating

$$G_{41} \leq ((\widehat{M}_{40})^{(8)})_2 = G_{41}^0 + 2(a_{41})^{(8)}((\widehat{M}_{40})^{(8)})_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq ((\widehat{M}_{40})^{(8)})_3 = G_{42}^0 + 2(a_{42})^{(8)}((\widehat{M}_{40})^{(8)})_2 / (a'_{42})^{(8)}$$

If  $G_{41}$  or  $G_{42}$  is bounded, the same property follows for  $G_{40}$  ,  $G_{42}$  and  $G_{40}$  ,  $G_{41}$  respectively.

**Remark 39:** If  $G_{40}$  is bounded, from below, the same property holds for  $G_{41}$  and  $G_{42}$  . The proof is 277  
 analogous with the preceding one. An analogous property is true if  $G_{41}$  is bounded from below.

**Remark 40:** If  $T_{40}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$  then  $T_{41} \rightarrow \infty$ . 278

**Definition of**  $(m)^{(8)}$  and  $\varepsilon_8$  :

Indeed let  $t_8$  be so that for  $t > t_8$

$$(b_{41})^{(8)} - (b''_i)^{(8)}((G_{43})(t), t) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then  $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$  which leads to 279

$$T_{41} \geq \left( \frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8} \right) (1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$$

If we take  $t$  such that  $e^{-\varepsilon_8 t} = \frac{1}{2}$  it results

$$T_{41} \geq \left( \frac{(a_{41})^{(8)}(m)^{(8)}}{2} \right), \quad t = \log \frac{2}{\varepsilon_8}$$

By taking now  $\varepsilon_8$  sufficiently small one sees that  $T_{41}$  is unbounded.

The same property holds for  $T_{42}$  if  $\lim_{t \rightarrow \infty} ((b''_{42})^{(8)}((G_{43})(t), t(t), t)) = (b'_{42})^{(8)}$

It is now sufficient to take  $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$  and to choose  $(\widehat{P}_{44})^{(9)}$  and  $(\widehat{Q}_{44})^{(9)}$  large to have 279  
 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[ (\widehat{P}_{44})^{(9)} + ((\widehat{P}_{44})^{(9)} + G_j^0) e^{-\left( \frac{((\widehat{P}_{44})^{(9)} + G_j^0)}{G_j^0} \right)} \right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \left[ ((\widehat{Q}_{44})^{(9)} + T_j^0) e^{-\left( \frac{((\widehat{Q}_{44})^{(9)} + T_j^0)}{T_j^0} \right)} + (\widehat{Q}_{44})^{(9)} \right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator  $\mathcal{A}^{(9)}$  transforms the space of sextuples of functions  $G_i, T_i$  satisfying 39,35,36

into itself

The operator  $\mathcal{A}^{(9)}$  is a contraction with respect to the metric

$$d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right), \left((G_{47})^{(2)}, (T_{47})^{(2)}\right)\right) = \sup_i \left\{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{44})^{(9)}t} \right\}$$

Indeed if we denote

**Definition of**  $(\widetilde{G_{47}}, \widetilde{T_{47}}) : ((\widetilde{G_{47}}, \widetilde{T_{47}})) = \mathcal{A}^{(9)}((G_{47}), (T_{47}))$

It results

$$\begin{aligned} |\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{44})^{(9)} |G_{45}^{(1)} - G_{45}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} ds_{(44)} + \\ &\int_0^t \{ (a'_{44})^{(9)} |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{-(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &(a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) |G_{44}^{(1)} - G_{44}^{(2)}| e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} + \\ &G_{44}^{(2)} | (a''_{44})^{(9)} (T_{45}^{(1)}, s_{(44)}) - (a''_{44})^{(9)} (T_{45}^{(2)}, s_{(44)}) | e^{-(\bar{M}_{44})^{(9)}s_{(44)}} e^{(\bar{M}_{44})^{(9)}s_{(44)}} \} ds_{(44)} \end{aligned}$$

Where  $s_{(44)}$  represents integrand that is integrated over the interval  $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\begin{aligned} |(G_{47})^{(1)} - G^{(2)}| e^{-(\bar{M}_{44})^{(9)}t} &\leq \frac{1}{(\bar{M}_{44})^{(9)}} \left( (a_{44})^{(9)} + (a'_{44})^{(9)} + (\hat{A}_{44})^{(9)} \right) \\ &+ (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)} d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}\right); (G_{47})^{(2)}, (T_{47})^{(2)}\right) \end{aligned}$$

And analogous inequalities for  $G_i$  and  $T_i$ . Taking into account the hypothesis (39,35,36) the result follows

**Remark 41:** The fact that we supposed  $(a''_{44})^{(9)}$  and  $(b''_{44})^{(9)}$  depending also on  $t$  can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by  $(\hat{P}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$  and  $(\hat{Q}_{44})^{(9)} e^{(\bar{M}_{44})^{(9)}t}$  respectively of  $\mathbb{R}_+$ .

If instead of proving the existence of the solution on  $\mathbb{R}_+$ , we have to prove it only on a compact then it suffices to consider that  $(a'_i)^{(9)}$  and  $(b'_i)^{(9)}$ ,  $i = 44, 45, 46$  depend only on  $T_{45}$  and respectively on  $(G_{47})$  (and not on  $t$ ) and hypothesis can be replaced by a usual Lipschitz condition.

**Remark 42:** There does not exist any  $t$  where  $G_i(t) = 0$  and  $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{ (a'_i)^{(9)} - (a''_i)^{(9)} (T_{45}(s_{(44)}), s_{(44)}) \} ds_{(44)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(9)}t} > 0 \text{ for } t > 0$$

**Definition of**  $((\widehat{M}_{44})^{(9)})_1, ((\widehat{M}_{44})^{(9)})_2$  and  $((\widehat{M}_{44})^{(9)})_3$  :

**Remark 43:**if  $G_{44}$  is bounded, the same property have also  $G_{45}$  and  $G_{46}$  . indeed if

$G_{44} < ((\widehat{M}_{44})^{(9)})$  it follows  $\frac{dG_{45}}{dt} \leq ((\widehat{M}_{44})^{(9)})_1 - (a'_{45})^{(9)}G_{45}$  and by integrating

$$G_{45} \leq ((\widehat{M}_{44})^{(9)})_2 = G_{45}^0 + 2(a_{45})^{(9)}((\widehat{M}_{44})^{(9)})_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq ((\widehat{M}_{44})^{(9)})_3 = G_{46}^0 + 2(a_{46})^{(9)}((\widehat{M}_{44})^{(9)})_2 / (a'_{46})^{(9)}$$

If  $G_{45}$  or  $G_{46}$  is bounded, the same property follows for  $G_{44}$  ,  $G_{46}$  and  $G_{44}$  ,  $G_{45}$  respectively.

**Remark 44:** If  $G_{44}$  is bounded, from below, the same property holds for  $G_{45}$  and  $G_{46}$  . The proof is analogous with the preceding one. An analogous property is true if  $G_{45}$  is bounded from below.

**Remark 45:** If  $T_{44}$  is bounded from below and  $\lim_{t \rightarrow \infty} ((b''_i)^{(9)}((G_{47})(t), t)) = (b'_{45})^{(9)}$  then  $T_{45} \rightarrow \infty$ .

**Definition of**  $(m)^{(9)}$  and  $\varepsilon_9$  :

Indeed let  $t_9$  be so that for  $t > t_9$

$$(b_{45})^{(9)} - (b''_i)^{(9)}((G_{47})(t), t) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then  $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$  which leads to

$$T_{45} \geq \left( \frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9} \right) (1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$$

If we take  $t$  such that  $e^{-\varepsilon_9 t} = \frac{1}{2}$  it results

$$T_{45} \geq \left( \frac{(a_{45})^{(9)}(m)^{(9)}}{2} \right), t = \log \frac{2}{\varepsilon_9}$$

By taking now  $\varepsilon_9$  sufficiently small one sees that  $T_{45}$  is unbounded.

The same property holds for  $T_{46}$  if  $\lim_{t \rightarrow \infty} (b''_{46})^{(9)}((G_{47})(t), t) = (b'_{46})^{(9)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

**Behavior of the solutions of equation**

280

**Theorem** If we denote and define

**Definition of**  $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$  :

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$  four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

**Definition of**  $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$  :

281

By  $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$  and respectively  $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$  the roots of the equations

$$(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0 \text{ and } (b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$$

**Definition of**  $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$  : 282

By  $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$  and respectively  $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$  the roots of the equations  $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$  and  $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

**Definition of**  $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$  :- 283

If we define  $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$  by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \quad \text{if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \quad \text{if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

and 
$$(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \quad \text{if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously 284

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \quad \text{if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \quad \text{if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

and 
$$(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}$ , if  $(\bar{u}_1)^{(1)} < (u_0)^{(1)}$  where  $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where  $(p_i)^{(1)}$  is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left( \frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \right) [e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t}] + G_{15}^0 e^{-(S_2)^{(1)}t} \leq G_{15}(t) \quad 286$$

$$\leq \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} [e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t}] + G_{15}^0 e^{-(a'_{15})^{(1)}t}$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)}T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})} \left[ e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \tag{289}$$

$$\frac{(a_{15})^{(1)}T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[ e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

**Definition of**  $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$ :- 290

Where  $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

**Behavior of the solutions of equation** 291

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$  : 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$  four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)} \tag{293}$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \tag{294}$$

**Definition of**  $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$  : 295

By  $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$  and respectively  $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$  the roots 296

of the equations  $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$  297

and  $(b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$  and 298

**Definition of**  $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$  : 299

By  $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$  and respectively  $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$  the 300

roots of the equations  $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$  301

and  $(b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$  302

**Definition of**  $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$  :- 303

If we define  $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$  by 304

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \quad \text{if } (v_0)^{(2)} < (v_1)^{(2)} \tag{305}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \quad \text{if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \tag{306}$$

and 
$$\boxed{(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \quad \mathbf{if}(\bar{v}_1)^{(2)} < (v_0)^{(2)} \tag{307}$$

and analogously 308

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \quad \mathbf{if}(u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \mathbf{if}(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

and 
$$\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \quad \mathbf{if}(\bar{u}_1)^{(2)} < (u_0)^{(2)} \tag{309}$$

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{(S_1)^{(2)} - (p_{16})^{(2)} t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$  is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} - (p_{16})^{(2)} t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \tag{311}$$

$$\left( \frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})} \right) \left[ e^{((S_1)^{(2)} - (p_{16})^{(2)}) t} - e^{-(S_2)^{(2)} t} \right] + G_{18}^0 e^{-(S_2)^{(2)} t} \leq G_{18}(t) \tag{312}$$

$$\leq \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[ e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)}) t}} \tag{313}$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)}) t} \tag{314}$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[ e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \leq T_{18}(t) \leq \tag{315}$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[ e^{((R_1)^{(2)} + (r_{16})^{(2)}) t} - e^{-(R_2)^{(2)} t} \right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

**Definition of**  $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$ :- 316

Where  $(S_1)^{(2)} = (a_{16})^{(2)} (m_2)^{(2)} - (a'_{16})^{(2)}$  317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

**Behavior of the solutions**

319

**Theorem 3:** If we denote and define

**Definition of**  $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$  :

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$  four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}(G_{23}, t) \leq -(\tau_1)^{(3)}$$

**Definition of**  $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$  :

320

By  $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$  and respectively  $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$  the roots of the equations

$$(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0 \text{ and}$$

By  $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$  and respectively  $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$  the

$$\text{roots of the equations } (a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$$

$$\text{and } (b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$$

**Definition of**  $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$  :-

321

If we define  $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$  by

$$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \quad \text{if } (v_0)^{(3)} < (v_1)^{(3)}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \quad \text{if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$$

$$\text{and } (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$$

$$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \quad \text{if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$$

and analogously

322

$$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \quad \text{if } (u_0)^{(3)} < (u_1)^{(3)}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \quad \text{if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \quad \text{and } (u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}$$

$$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \quad \text{if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$  is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t} \quad 323$$

$$\begin{aligned} & \left( \frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[ e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \quad 324 \\ & \leq \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[ e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t} \end{aligned}$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)} ((R_1)^{(3)} - (b'_{22})^{(3)})} \left[ e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)} ((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[ e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$$

**Definition of**  $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$ :- 328

$$\text{Where } (S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

**Behavior of the solutions of equation**

**Theorem:** If we denote and define

**Definition of**  $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$  :

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$  four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

**Definition of**  $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$  : 329

By  $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$  and respectively  $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$  the roots of the equations

$$(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$  : 330

By  $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$  and respectively  $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$  the roots of the equations  $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$  and  $(b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

**Definition of**  $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$  :-

If we define  $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$  by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \quad \text{if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \quad \text{if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

and  $(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \quad \text{if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \quad \text{if } (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \quad \text{if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

and  $(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \quad \text{if } (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where  $(p_i)^{(4)}$  is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left( \frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[ e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \right) \leq G_{26}(t) \leq \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[ e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[ e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)}T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[ e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

**Definition of**  $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$ :- 337

$$\begin{aligned} \text{Where } (S_1)^{(4)} &= (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)} \\ (S_2)^{(4)} &= (a_{26})^{(4)} - (p_{26})^{(4)} \\ (R_1)^{(4)} &= (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)} \\ (R_2)^{(4)} &= (b'_{26})^{(4)} - (r_{26})^{(4)} \end{aligned}$$

**Behavior of the solutions of equation** 338

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$  :

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$  four constants satisfying

$$\begin{aligned} -(\sigma_2)^{(5)} &\leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)} \\ -(\tau_2)^{(5)} &\leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)} \end{aligned}$$

**Definition of**  $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$  : 339

By  $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$  and respectively  $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$  the roots of the equations

$$\begin{aligned} (a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} &= 0 \\ \text{and } (b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} &= 0 \text{ and} \end{aligned}$$

**Definition of**  $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$  : 340

By  $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$  and respectively  $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$  the roots of the equations  $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$  and  $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

**Definition of**  $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$  :-

If we define  $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$  by

$$\begin{aligned} (m_2)^{(5)} &= (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \quad \text{if } (v_0)^{(5)} < (v_1)^{(5)} \\ (m_2)^{(5)} &= (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \quad \text{if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)}, \\ \text{and } (v_0)^{(5)} &= \frac{G_{28}^0}{G_{29}^0} \end{aligned}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \quad \text{if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously 341

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \quad \text{if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

and  $(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)} \text{ where } (u_1)^{(5)}, (\bar{u}_1)^{(5)}$$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where  $(p_i)^{(5)}$  is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left( \frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \left[ e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \leq G_{30}(t) \right. \\ \left. \leq \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[ e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t} \right) \quad 344$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[ e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[ e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

**Definition of**  $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$ :- 348

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)} (m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)} (\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

**Behavior of the solutions of equation** 349

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$  :

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$  four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

**Definition of**  $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$  : 350

By  $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$  and respectively  $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$  the roots of the equations  
 $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$   
 and  $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$  and

**Definition of**  $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$  : 351

By  $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$  and respectively  $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$  the roots of the equations  $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$   
 and  $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

**Definition of**  $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$  :-

If we define  $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$  by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \quad \text{if } (v_0)^{(6)} < (v_1)^{(6)}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \quad \text{if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

and  $(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \quad \text{if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously 352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \quad \text{if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \quad \text{if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

and  $(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \quad \text{if } (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(S_1)^{(6)}t}$$

where  $(p_i)^{(6)}$  is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left( \frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} ((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})} \left[ e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t} \right] + G_{34}^0 e^{-(S_2)^{(6)}t} \leq G_{34}(t) \right. \\ \left. \leq \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} ((S_1)^{(6)} - (a'_{34})^{(6)})} \left[ e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)}t} \right) \quad 355$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} ((R_1)^{(6)} - (b'_{34})^{(6)})} \left[ e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \leq T_{34}(t) \leq \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} ((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})} \left[ e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t} \right] + T_{34}^0 e^{-(R_2)^{(6)}t}$$

**Definition of**  $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$ :- 359

Where  $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

**Behavior of the solutions of equation**

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$  :

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$  four constants satisfying

$$-(\sigma_2)^{(7)} \leq -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \leq -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \leq -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}((G_{39}), t) - (b''_{37})^{(7)}((G_{39}), t) \leq -(\tau_1)^{(7)}$$

**Definition of**  $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$  : 361

By  $(v_1)^{(7)} > 0, (v_2)^{(7)} < 0$  and respectively  $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$  the roots of the equations

$$(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$$

and  $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$  and

**Definition of**  $(\bar{v}_1)^{(7)}, (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$  : 362

By  $(\bar{v}_1)^{(7)} > 0, (\bar{v}_2)^{(7)} < 0$  and respectively  $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$  the

roots of the equations  $(a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$

and  $(b_{37})^{(7)}(u^{(7)})^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

**Definition of**  $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$  :-

If we define  $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$  by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \quad \text{if } (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \quad \text{if } (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

and  $(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \quad \text{if } (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \quad \text{if } (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \quad \text{if } (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and  $(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \quad \text{if } (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)}t}$$

where  $(p_i)^{(7)}$  is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t}$$

$$\left( \frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)} ((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})} \right) [e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t}] + G_{38}^0 e^{-(S_2)^{(7)}t} \leq G_{38}(t)$$

$$\leq \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)} ((S_1)^{(7)} - (a'_{38})^{(7)})} [e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t}] + G_{38}^0 e^{-(a'_{38})^{(7)}t}$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}}$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}$$

$$\frac{(b_{38})^{(7)}T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})} \left[ e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t} \right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \leq T_{38}(t) \leq \tag{369}$$

$$\frac{(a_{38})^{(7)}T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})} \left[ e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t} \right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

**Definition of**  $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$ :- 370

Where  $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$$

$$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

**Behavior of the solutions of equation** 371

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$  :

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$  four constants satisfying

$$-(\sigma_2)^{(8)} \leq -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \leq -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \leq -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}((G_{43}), t) - (b''_{41})^{(8)}((G_{43}), t) \leq -(\tau_1)^{(8)}$$

**Definition of**  $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$  : 372

By  $(v_1)^{(8)} > 0, (v_2)^{(8)} < 0$  and respectively  $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$  the roots of the equations

$$(a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$$

and  $(b_{41})^{(8)}(u^{(8)})^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$  and

**Definition of**  $(\bar{v}_1)^{(8)}, (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$  :

By  $(\bar{v}_1)^{(8)} > 0, (\bar{v}_2)^{(8)} < 0$  and respectively  $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$  the

$$\text{roots of the equations } (a_{41})^{(8)}(\bar{v}^{(8)})^2 + (\sigma_2)^{(8)}\bar{v}^{(8)} - (a_{40})^{(8)} = 0$$

$$\text{and } (b_{41})^{(8)}(\bar{u}^{(8)})^2 + (\tau_2)^{(8)}\bar{u}^{(8)} - (b_{40})^{(8)} = 0$$

**Definition of**  $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$  :-

If we define  $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$  by

$$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}, \quad \mathbf{if} (v_0)^{(8)} < (v_1)^{(8)}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}, \quad \mathbf{if} (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)},$$

$$\text{and } \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$$

$$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}, \quad \mathbf{if} (\bar{v}_1)^{(8)} < (v_0)^{(8)}$$

and analogously

374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \quad \mathbf{if} (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \quad \mathbf{if} (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

$$\text{and } \boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \quad \mathbf{if} (\bar{u}_1)^{(8)} < (u_0)^{(8)} \text{ where } (u_1)^{(8)}, (\bar{u}_1)^{(8)}$$

Then the solution of global equations satisfies the inequalities

375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{40}(t) \leq G_{40}^0 e^{(S_1)^{(8)}t}$$

where  $(p_i)^{(8)}$  is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \leq G_{41}(t) \leq \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\begin{aligned} & \left( \frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)} ((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})} \left[ e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t} \right] + G_{42}^0 e^{-(S_2)^{(8)}t} \right) \leq G_{42}(t) \quad 377 \\ & \leq \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)} ((S_1)^{(8)} - (a'_{42})^{(8)})} \left[ e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t} \right] + G_{42}^0 e^{-(a'_{42})^{(8)}t} \end{aligned}$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \leq T_{40}(t) \leq \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)} ((R_1)^{(8)} - (b'_{42})^{(8)})} \left[ e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t} \right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \leq T_{42}(t) \leq \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)} ((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})} \left[ e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t} \right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

**Definition of**  $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$ :-

381

Where  $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

**Behavior of the solutions of equation 37 to 92**

382

**Theorem 2:** If we denote and define

**Definition of**  $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$  :

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$  four constants satisfying

$$-(\sigma_2)^{(9)} \leq -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \leq -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \leq -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}((G_{47}), t) - (b''_{45})^{(9)}((G_{47}), t) \leq -(\tau_1)^{(9)}$$

**Definition of**  $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$  :

By  $(v_1)^{(9)} > 0, (v_2)^{(9)} < 0$  and respectively  $(u_1)^{(9)} > 0, (u_2)^{(9)} < 0$  the roots of the equations

$$(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$$

$$\text{and } (b_{45})^{(9)}(u^{(9)})^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0 \text{ and}$$

**Definition of**  $(\bar{v}_1)^{(9)}, (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$  :

By  $(\bar{v}_1)^{(9)} > 0, (\bar{v}_2)^{(9)} < 0$  and respectively  $(\bar{u}_1)^{(9)} > 0, (\bar{u}_2)^{(9)} < 0$  the

roots of the equations  $(a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and  $(b_{45})^{(9)}(u^{(9)})^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

**Definition of**  $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$  :-

If we define  $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$  by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \quad \text{if } (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \quad \text{if } (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

$$\text{and } (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \quad \text{if } (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(m_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \quad \text{if } (u_0)^{(9)} < (u_1)^{(9)}$$

$$(m_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \quad \text{if } (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

and  $(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$ , if  $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$  where  $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$  are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)}t}$$

where  $(p_i)^{(9)}$  is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)}t}$$

(

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})} \left[ e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)}t} \right] + G_{46}^0 e^{-(S_2)^{(9)}t} \leq G_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a_{46})^{(9)})} \left[ e^{(S_1)^{(9)}t} - e^{-(a'_{46})^{(9)}t} \right] + G_{46}^0 e^{-(a'_{46})^{(9)}t}$$

$$T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)}t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})} \left[ e^{(R_1)^{(9)}t} - e^{-(b'_{46})^{(9)}t} \right] + T_{46}^0 e^{-(b'_{46})^{(9)}t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})} \left[ e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)}t} \right] + T_{46}^0 e^{-(R_2)^{(9)}t}$$

**Definition of**  $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$ :-

Where  $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

**Proof:** From global equations we obtain

383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left( (a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t) \right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

**Definition of**  $v^{(1)}$  :-

$$v^{(1)} = \frac{G_{13}}{G_{14}}$$

It follows

$$-\left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \leq \frac{dv^{(1)}}{dt} \leq -\left((a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$  :-

For  $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}}{1 + (C)^{(1)}e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_0)^{(1)})t]}} , \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

it follows  $(v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$

In the same manner , we get

384

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)}e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}$$

From which we deduce  $(v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$

If  $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$  we find like in the previous case,

385

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)}(v_2)^{(1)}e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}}{1 + (C)^{(1)}e^{[-(a_{14})^{(1)}((v_1)^{(1)} - (v_2)^{(1)})t]}} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)}e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (\bar{v}_1)^{(1)}$$

If  $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$  , we obtain

386

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)}(\bar{v}_2)^{(1)}e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}}{1 + (\bar{C})^{(1)}e^{[-(a_{14})^{(1)}((\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)})t]}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(1)}(t)$  :-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)} , \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(1)}(t)$  :-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If  $(a''_{13})^{(1)} = (a''_{14})^{(1)}$ , then  $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$  and in this case  $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$  if in addition  $(v_0)^{(1)} = (v_1)^{(1)}$  then  $v^{(1)}(t) = (v_0)^{(1)}$  and as a consequence  $G_{13}(t) = (v_0)^{(1)}G_{14}(t)$  this also defines  $(v_0)^{(1)}$  for the special case

Analogously if  $(b''_{13})^{(1)} = (b''_{14})^{(1)}$ , then  $(\tau_1)^{(1)} = (\tau_2)^{(1)}$  and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$  if in addition  $(u_0)^{(1)} = (u_1)^{(1)}$  then  $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$  This is an important consequence of the relation between  $(v_1)^{(1)}$  and  $(\bar{v}_1)^{(1)}$ , and definition of  $(u_0)^{(1)}$ .

**Proof :** From global equations we obtain 387

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left( (a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t) \right) - (a''_{17})^{(2)}(T_{17}, t)v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

**Definition of**  $v^{(2)}$  :- 388

$$\boxed{v^{(2)} = \frac{G_{16}}{G_{17}}}$$

It follows 389

$$- \left( (a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} \right) \leq \frac{dv^{(2)}}{dt} \leq - \left( (a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} \right)$$

From which one obtains 390

**Definition of**  $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$  :-

$$\text{For } 0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_0)^{(2)})t]}} , \quad \boxed{(C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

$$\text{it follows } (v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$$

In the same manner , we get 391

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce  $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$  392

$$\text{If } 0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)} \text{ we find like in the previous case,} \quad \text{393}$$

$$\begin{aligned}
 (v_1)^{(2)} &\leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}}{1 + (C)^{(2)} e^{[-(a_{17})^{(2)}((v_1)^{(2)} - (v_2)^{(2)})t]}} \leq v^{(2)}(t) \leq \\
 &\frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (\bar{v}_1)^{(2)}
 \end{aligned}$$

If  $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$ , we obtain 394

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}}{1 + (\bar{C})^{(2)} e^{[-(a_{17})^{(2)}((\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)})t]}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c), we have

**Definition of  $v^{(2)}(t)$  :-** 395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}$$

In a completely analogous way, we obtain 396

**Definition of  $u^{(2)}(t)$  :-**

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :** 397

If  $(a_{16}'' )^{(2)} = (a_{17}'' )^{(2)}$ , then  $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$  and in this case  $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$  if in addition  $(v_0)^{(2)} = (v_1)^{(2)}$  then  $v^{(2)}(t) = (v_0)^{(2)}$  and as a consequence  $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if  $(b_{16}'' )^{(2)} = (b_{17}'' )^{(2)}$ , then  $(\tau_1)^{(2)} = (\tau_2)^{(2)}$  and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$  if in addition  $(u_0)^{(2)} = (u_1)^{(2)}$  then  $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$  This is an important consequence of the relation between  $(v_1)^{(2)}$  and  $(\bar{v}_1)^{(2)}$

**Proof :** From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left( (a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

**Definition of**  $v^{(3)}$  :- 
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$
 399

It follows

$$-\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For  $0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_0)^{(3)})t]}}$$

$$(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

it follows  $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}$$

$$(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

**Definition of**  $(\bar{v}_1)^{(3)}$  :-

From which we deduce  $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

If  $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$  we find like in the previous case, 402

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}}{1 + (C)^{(3)}e^{[-(a_{21})^{(3)}((v_1)^{(3)} - (v_2)^{(3)})t]}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (\bar{v}_1)^{(3)}$$

If  $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$  , we obtain 403

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (\bar{C})^{(3)}e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(3)}(t)$  :-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(3)}(t)$  :-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}$ , then  $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$  and in this case  $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$  if in addition  $(v_0)^{(3)} = (v_1)^{(3)}$  then  $v^{(3)}(t) = (v_0)^{(3)}$  and as a consequence  $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if  $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}$ , then  $(\tau_1)^{(3)} = (\tau_2)^{(3)}$  and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$  if in addition  $(u_0)^{(3)} = (u_1)^{(3)}$  then  $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$  This is an important consequence of the relation between  $(v_1)^{(3)}$  and  $(\bar{v}_1)^{(3)}$

**Proof:** From global equations we obtain

404

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left( (a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

**Definition of**  $v^{(4)}$  :-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

It follows

$$- \left( (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left( (a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$  :-

$$\text{For } 0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}$$

$$\text{it follows } (v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$$

In the same manner , we get

405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}}{4 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)})t]}} , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}$$

From which we deduce  $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If  $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$  we find like in the previous case,

406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)}-(v_2)^{(4)})t]}}{1 + (C)^{(4)}e^{[-(a_{25})^{(4)}((v_1)^{(4)}-(v_2)^{(4)})t]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)})t]}} \leq (\bar{v}_1)^{(4)}$$

If  $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$ , we obtain

407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)})t]}}{1 + (\bar{C})^{(4)}e^{[-(a_{25})^{(4)}((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)})t]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(4)}(t)$  :-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(4)}(t)$  :-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a_{24})''^{(4)} = (a_{25})''^{(4)}$ , then  $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$  and in this case  $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$  if in addition  $(v_0)^{(4)} = (v_1)^{(4)}$  then  $v^{(4)}(t) = (v_0)^{(4)}$  and as a consequence  $G_{24}(t) = (v_0)^{(4)}G_{25}(t)$  **this also defines  $(v_0)^{(4)}$  for the special case.**

Analogously if  $(b_{24})''^{(4)} = (b_{25})''^{(4)}$ , then  $(\tau_1)^{(4)} = (\tau_2)^{(4)}$  and then  $(u_1)^{(4)} = (\bar{u}_1)^{(4)}$  if in addition  $(u_0)^{(4)} = (u_1)^{(4)}$  then  $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$  This is an important consequence of the relation between  $(v_1)^{(4)}$  and  $(\bar{v}_1)^{(4)}$ , **and definition of  $(u_0)^{(4)}$ .**

408

**Proof :** From global equations we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left( (a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t) \right) - (a''_{29})^{(5)}(T_{29}, t)v^{(5)} - (a_{29})^{(5)}v^{(5)}$$

**Definition of**  $v^{(5)}$  :-  $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left( (a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left( (a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$  :-

$$\text{For } 0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}$$

it follows  $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

409

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}$$

From which we deduce  $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

If  $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$  we find like in the previous case,

410

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

If  $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$  , we obtain

411

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(5)}(t)$  :-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)} , \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(5)}(t)$  :-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)} , \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a_{28})^{(5)} = (a_{29})^{(5)}$ , then  $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$  and in this case  $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$  if in addition  $(v_0)^{(5)} = (v_5)^{(5)}$  then  $v^{(5)}(t) = (v_0)^{(5)}$  and as a consequence  $G_{28}(t) = (v_0)^{(5)}G_{29}(t)$  **this also defines  $(v_0)^{(5)}$  for the special case .**

Analogously if  $(b''_{28})^{(5)} = (b''_{29})^{(5)}$ , then  $(\tau_1)^{(5)} = (\tau_2)^{(5)}$  and then  $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$  if in addition  $(u_0)^{(5)} = (u_1)^{(5)}$  then  $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$  This is an important consequence of the relation between  $(v_1)^{(5)}$  and  $(\bar{v}_1)^{(5)}$ , and definition of  $(u_0)^{(5)}$ .

**Proof:** From global equations we obtain

412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left( (a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

**Definition of**  $v^{(6)}$  :- 
$$v^{(6)} = \frac{G_{32}}{G_{33}}$$

It follows

$$- \left( (a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left( (a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$  :-

For  $0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}(v_1)^{(6)} - (v_0)^{(6)}]t}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}(v_1)^{(6)} - (v_0)^{(6)}]t}}, \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

$$\text{it follows } (v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$$

In the same manner, we get

413

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}}, \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

From which we deduce  $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

If  $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$  we find like in the previous case,

414

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}(v_1)^{(6)} - (v_2)^{(6)}]t}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}(v_1)^{(6)} - (v_2)^{(6)}]t}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}} \leq (\bar{v}_1)^{(6)}$$

If  $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$ , we obtain

415

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}(\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)}]t}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c), we have

**Definition of**  $v^{(6)}(t)$  :-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(6)}(t)$  :-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{32})^{(6)} = (a''_{33})^{(6)}$ , then  $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$  and in this case  $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$  if in addition  $(v_0)^{(6)} = (v_1)^{(6)}$  then  $v^{(6)}(t) = (v_0)^{(6)}$  and as a consequence  $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$  **this also defines  $(v_0)^{(6)}$  for the special case.**

Analogously if  $(b''_{32})^{(6)} = (b''_{33})^{(6)}$ , then  $(\tau_1)^{(6)} = (\tau_2)^{(6)}$  and then  $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$  if in addition  $(u_0)^{(6)} = (u_1)^{(6)}$  then  $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$  This is an important consequence of the relation between  $(v_1)^{(6)}$  and  $(\bar{v}_1)^{(6)}$ , **and definition of  $(u_0)^{(6)}$ .**

416

**Proof :** From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left( (a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t) \right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

**Definition of**  $v^{(7)}$  :-  $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$- \left( (a_{37})^{(7)}(v^{(7)})^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} \right) \leq \frac{dv^{(7)}}{dt} \leq - \left( (a_{37})^{(7)}(v^{(7)})^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$  :-

$$\text{For } 0 < \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_0)^{(7)}]t}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_0)^{(7)}]t}}, \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}$$

$$\text{it follows } (v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$$

In the same manner, we get

417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}$$

From which we deduce  $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If  $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{v}_1)^{(7)}$  we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_2)^{(7)}]t}}{1 + (C)^{(7)} e^{[-(a_{37})^{(7)}(v_1)^{(7)} - (v_2)^{(7)}]t}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}} \leq (\bar{v}_1)^{(7)}$$

If  $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$ , we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}}{1 + (\bar{C})^{(7)} e^{[-(a_{37})^{(7)}(\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}]t}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c), we have

**Definition of  $v^{(7)}(t)$  :-**

$$(m_2)^{(7)} \leq v^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

**Definition of  $u^{(7)}(t)$  :-** 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a_{36}''^{(7)}) = (a_{37}''^{(7)})$ , then  $(\sigma_1)^{(7)} = (\sigma_2)^{(7)}$  and in this case  $(v_1)^{(7)} = (\bar{v}_1)^{(7)}$  if in addition  $(v_0)^{(7)} = (v_1)^{(7)}$  then  $v^{(7)}(t) = (v_0)^{(7)}$  and as a consequence  $G_{36}(t) = (v_0)^{(7)}G_{37}(t)$  **this also defines  $(v_0)^{(7)}$  for the special case .**

Analogously if  $(b_{36}''^{(7)}) = (b_{37}''^{(7)})$ , then  $(\tau_1)^{(7)} = (\tau_2)^{(7)}$  and then  $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$  if in addition  $(u_0)^{(7)} = (u_1)^{(7)}$  then  $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$  This is an important consequence of the relation between  $(v_1)^{(7)}$  and  $(\bar{v}_1)^{(7)}$ , **and definition of  $(u_0)^{(7)}$ .**

**Proof:** From global equations we obtain

421

$$\frac{dv^{(8)}}{dt} = (a_{40})^{(8)} - \left( (a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t) \right) - (a''_{41})^{(8)}(T_{41}, t)v^{(8)} - (a_{41})^{(8)}v^{(8)}$$

**Definition of**  $v^{(8)}$  :- 
$$v^{(8)} = \frac{G_{40}}{G_{41}}$$

It follows

$$-\left( (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} \right) \leq \frac{dv^{(8)}}{dt} \leq -\left( (a_{41})^{(8)}(v^{(8)})^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$  :-

For  $0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}(v_1)^{(8)} - (v_0)^{(8)}]t}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}(v_1)^{(8)} - (v_0)^{(8)}]t}}, \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}$$

it follows  $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get

422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}}, \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}$$

From which we deduce  $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If  $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$  we find like in the previous case,

423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)} e^{[-(a_{41})^{(8)}(v_1)^{(8)} - (v_2)^{(8)}]t}}{1 + (C)^{(8)} e^{[-(a_{41})^{(8)}(v_1)^{(8)} - (v_2)^{(8)}]t}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}} \leq (\bar{v}_1)^{(8)}$$

If  $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$ , we obtain

424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}}{1 + (\bar{C})^{(8)} e^{[-(a_{41})^{(8)}(\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}]t}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

**Definition of**  $v^{(8)}(t)$  :-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

**Definition of**  $u^{(8)}(t)$  :-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{40})^{(8)} = (a''_{41})^{(8)}$ , then  $(\sigma_1)^{(8)} = (\sigma_2)^{(8)}$  and in this case  $(v_1)^{(8)} = (\bar{v}_1)^{(8)}$  if in addition  $(v_0)^{(8)} = (v_1)^{(8)}$  then  $v^{(8)}(t) = (v_0)^{(8)}$  and as a consequence  $G_{40}(t) = (v_0)^{(8)}G_{41}(t)$  **this also defines  $(v_0)^{(8)}$  for the special case.**

Analogously if  $(b''_{40})^{(8)} = (b''_{41})^{(8)}$ , then  $(\tau_1)^{(8)} = (\tau_2)^{(8)}$  and then  $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$  if in addition  $(u_0)^{(8)} = (u_1)^{(8)}$  then  $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$  This is an important consequence of the relation between  $(v_1)^{(8)}$  and  $(\bar{v}_1)^{(8)}$ , **and definition of  $(u_0)^{(8)}$ .**

**Proof :** From 99,20,44,22,23,44 we obtain

424  
A

$$\frac{dv^{(9)}}{dt} = (a_{44})^{(9)} - \left( (a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t) \right) - (a''_{45})^{(9)}(T_{45}, t)v^{(9)} - (a_{45})^{(9)}v^{(9)}$$

**Definition of**  $v^{(9)}$  :-  $\boxed{v^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$- \left( (a_{45})^{(9)}(v^{(9)})^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} \right) \leq \frac{dv^{(9)}}{dt} \leq - \left( (a_{45})^{(9)}(v^{(9)})^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} \right)$$

From which one obtains

**Definition of**  $(\bar{v}_1)^{(9)}, (v_0)^{(9)}$  :-

For  $0 < \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (v_1)^{(9)} < (\bar{v}_1)^{(9)}$

$$v^{(9)}(t) \geq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}}{1 + (C)^{(9)} e^{[-(a_{45})^{(9)}((v_1)^{(9)} - (v_0)^{(9)})t]}} , \quad \boxed{(C)^{(9)} = \frac{(v_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (v_2)^{(9)}}$$

it follows  $(v_0)^{(9)} \leq v^{(9)}(t) \leq (v_9)^{(9)}$

In the same manner , we get

$$v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)} e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)} - (\bar{v}_2)^{(9)})t]}} , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{v}_1)^{(9)} - (v_0)^{(9)}}{(v_0)^{(9)} - (\bar{v}_2)^{(9)}}$$

From which we deduce  $(v_0)^{(9)} \leq v^{(9)}(t) \leq (\bar{v}_1)^{(9)}$

If  $0 < (v_1)^{(9)} < (v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{v}_1)^{(9)}$  we find like in the previous case,

$$(v_1)^{(9)} \leq \frac{(v_1)^{(9)} + (C)^{(9)}(v_2)^{(9)}e^{[-(a_{45})^{(9)}((v_1)^{(9)}-(v_2)^{(9)})t]}}{1 + (C)^{(9)}e^{[-(a_{45})^{(9)}((v_1)^{(9)}-(v_2)^{(9)})t]}} \leq v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)}e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)}-(\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)}e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)}-(\bar{v}_2)^{(9)})t]}} \leq (\bar{v}_1)^{(9)}$$

If  $0 < (v_1)^{(9)} \leq (\bar{v}_1)^{(9)} \leq \boxed{(v_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$ , we obtain

$$(v_1)^{(9)} \leq v^{(9)}(t) \leq \frac{(\bar{v}_1)^{(9)} + (\bar{C})^{(9)}(\bar{v}_2)^{(9)}e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)}-(\bar{v}_2)^{(9)})t]}}{1 + (\bar{C})^{(9)}e^{[-(a_{45})^{(9)}((\bar{v}_1)^{(9)}-(\bar{v}_2)^{(9)})t]}} \leq (v_0)^{(9)}$$

And so with the notation of the first part of condition (c), we have

**Definition of  $v^{(9)}(t)$  :-**

$$(m_2)^{(9)} \leq v^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{v^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

**Definition of  $u^{(9)}(t)$  :-**

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

**Particular case :**

If  $(a''_{44})^{(9)} = (a''_{45})^{(9)}$ , then  $(\sigma_1)^{(9)} = (\sigma_2)^{(9)}$  and in this case  $(v_1)^{(9)} = (\bar{v}_1)^{(9)}$  if in addition  $(v_0)^{(9)} = (v_1)^{(9)}$  then  $v^{(9)}(t) = (v_0)^{(9)}$  and as a consequence  $G_{44}(t) = (v_0)^{(9)}G_{45}(t)$  **this also defines  $(v_0)^{(9)}$  for the special case .**

Analogously if  $(b''_{44})^{(9)} = (b''_{45})^{(9)}$ , then  $(\tau_1)^{(9)} = (\tau_2)^{(9)}$  and then  $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$  if in addition  $(u_0)^{(9)} = (u_1)^{(9)}$  then  $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$  This is an important consequence of the relation between  $(v_1)^{(9)}$  and  $(\bar{v}_1)^{(9)}$ , **and definition of  $(u_0)^{(9)}$ .**

We can prove the following

425

**Theorem :** If  $(a_i'')^{(1)}$  and  $(b_i'')^{(1)}$  are independent on  $t$ , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with  $(p_{13})^{(1)}, (r_{14})^{(1)}$  as defined by equation are satisfied , then the system

**Theorem :** If  $(a_i'')^{(2)}$  and  $(b_i'')^{(2)}$  are independent on  $t$  , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \quad 427$$

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0 , \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

with  $(p_{16})^{(2)}, (r_{17})^{(2)}$  as defined by equation are satisfied , then the system

**Theorem :** If  $(a_i'')^{(3)}$  and  $(b_i'')^{(3)}$  are independent on  $t$  , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0 ,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with  $(p_{20})^{(3)}, (r_{21})^{(3)}$  as defined by equation are satisfied , then the system

We can prove the following 432

**Theorem :** If  $(a_i'')^{(4)}$  and  $(b_i'')^{(4)}$  are independent on  $t$  , and the conditions with the notations

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0 ,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with  $(p_{24})^{(4)}, (r_{25})^{(4)}$  as defined by equation are satisfied , then the system

**Theorem :** If  $(a_i'')^{(5)}$  and  $(b_i'')^{(5)}$  are independent on  $t$  , and the conditions with the notations 433

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0 ,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with  $(p_{28})^{(5)}, (r_{29})^{(5)}$  as defined by equation are satisfied , then the system

**Theorem** If  $(a_i'')^{(6)}$  and  $(b_i'')^{(6)}$  are independent on  $t$ , and the conditions with the notations 434

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$

with  $(p_{32})^{(6)}, (r_{33})^{(6)}$  as defined by equation are satisfied, then the system

**Theorem :** If  $(a_i'')^{(7)}$  and  $(b_i'')^{(7)}$  are independent on  $t$ , and the conditions with the notations 435

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$$

$$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0,$$

$$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$$

with  $(p_{36})^{(7)}, (r_{37})^{(7)}$  as defined by equation are satisfied, then the system

**Theorem :** If  $(a_i'')^{(8)}$  and  $(b_i'')^{(8)}$  are independent on  $t$ , and the conditions with the notations 436

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$$

$$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0,$$

$$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$$

with  $(p_{40})^{(8)}, (r_{41})^{(8)}$  as defined by equation are satisfied, then the system

**Theorem :** If  $(a_i'')^{(9)}$  and  $(b_i'')^{(9)}$  are independent on  $t$ , and the conditions (with the notations 436  
 45,46,27,28) A

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$$

$$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0,$$

$$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$$

with  $(p_{44})^{(9)}, (\tau_{45})^{(9)}$  as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27})]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27})]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27})]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - [(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - [(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - [(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - [(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - [(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - [(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - [(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})]G_{44} = 0 \quad 484$$

A

$$(a_{45})^{(9)}G_{44} - [(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - [(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})]T_{46} = 0$$

**Proof:** 485

(a) Indeed the first two equations have a nontrivial solution  $G_{13}, G_{14}$  if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) \\ + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

**Proof:** 486

(b) Indeed the first two equations have a nontrivial solution  $G_{16}, G_{17}$  if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) \\ + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

**Proof:** 487

(a) Indeed the first two equations have a nontrivial solution  $G_{20}, G_{21}$  if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

**Proof:** 488

(a) Indeed the first two equations have a nontrivial solution  $G_{24}, G_{25}$  if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

**Proof:** 489

(a) Indeed the first two equations have a nontrivial solution  $G_{28}, G_{29}$  if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

**Proof:** 490

(a) Indeed the first two equations have a nontrivial solution  $G_{32}, G_{33}$  if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

**Proof:** 491

(a) Indeed the first two equations have a nontrivial solution  $G_{36}, G_{37}$  if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

**Proof:** 492

(a) Indeed the first two equations have a nontrivial solution  $G_{40}, G_{41}$  if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

**Proof:** 492

(a) Indeed the first two equations have a nontrivial solution  $G_{44}, G_{45}$  if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

A

**Definition and uniqueness of  $\Gamma_{14}^*$  :-** 493

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(1)}(T_{14})$  being increasing, it follows that there exists a unique  $T_{14}^*$  for which  $f(T_{14}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^*)]}$$

**Definition and uniqueness of  $\Gamma_{17}^*$  :-** 494

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(2)}(T_{17})$  being increasing, it follows that there exists a unique  $T_{17}^*$  for which  $f(T_{17}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^*)]}$$
495

**Definition and uniqueness of  $\Gamma_{21}^*$  :-** 496

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(1)}(T_{21})$  being increasing, it follows that there exists a unique  $T_{21}^*$  for which  $f(T_{21}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^*)]}$$

**Definition and uniqueness of  $\Gamma_{25}^*$  :-** 497

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(4)}(T_{25})$  being increasing, it follows that there exists a unique  $T_{25}^*$  for which  $f(T_{25}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^*)]}$$

**Definition and uniqueness of  $\Gamma_{29}^*$  :-** 498

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(5)}(T_{29})$  being increasing, it follows that there exists a unique  $T_{29}^*$  for which  $f(T_{29}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

**Definition and uniqueness of  $\Gamma_{33}^*$  :-** 499

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(6)}(T_{33})$  being increasing, it follows that there exists a unique  $T_{33}^*$  for which  $f(T_{33}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

**Definition and uniqueness of  $\Gamma_{37}^*$  :-** 500

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a''_i)^{(7)}(T_{37})$  being increasing, it follows that there exists a unique  $T_{37}^*$  for which  $f(T_{37}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

**Definition and uniqueness of  $\Gamma_{41}^*$  :-**

501

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(8)}(T_{41})$  being increasing, it follows that there exists a unique  $T_{41}^*$  for which  $f(T_{41}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)}G_{41}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)}G_{41}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^*)]}$$

**Definition and uniqueness of  $\Gamma_{45}^*$  :-**

501

A

After hypothesis  $f(0) < 0, f(\infty) > 0$  and the functions  $(a_i'')^{(9)}(T_{45})$  being increasing, it follows that there exists a unique  $T_{45}^*$  for which  $f(T_{45}^*) = 0$ . With this value, we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)}G_{45}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)}G_{45}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions  $G_{13}, G_{14}$  if

502

$$\begin{aligned} \varphi(G) &= (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - \\ &[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0 \end{aligned}$$

Where in  $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{14}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{14}^*$  such that  $\varphi(G^*) = 0$

By the same argument, the equations admit solutions  $G_{16}, G_{17}$  if

503

$$\begin{aligned} \varphi(G_{19}) &= (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - \\ &[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0 \end{aligned}$$

Where in  $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{17}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{17}^*$  such that  $\varphi((G_{19})^*) = 0$

504

By the same argument, the equations admit solutions  $G_{20}, G_{21}$  if

505

$$\begin{aligned} \varphi(G_{23}) &= (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - \\ &[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0 \end{aligned}$$

Where in  $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is

a decreasing function in  $G_{21}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{21}^*$  such that  $\varphi((G_{21})^*) = 0$

By the same argument, the equations admit solutions  $G_{24}, G_{25}$  if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - [(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in  $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{25}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{25}^*$  such that  $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions  $G_{28}, G_{29}$  if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - [(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in  $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{29}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{29}^*$  such that  $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions  $G_{32}, G_{33}$  if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - [(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in  $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{33}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{33}^*$  such that  $\varphi(G_{35}^*) = 0$

By the same argument, the equations admit solutions  $G_{36}, G_{37}$  if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - [(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in  $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{37}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{37}^*$  such that  $\varphi(G_{39}^*) = 0$

By the same argument, the equations admit solutions  $G_{40}, G_{41}$  if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - [(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in  $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{41}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{41}^*$  such that  $\varphi(G_{43}^*) = 0$

By the same argument, the equations 92,93 admit solutions  $G_{44}, G_{45}$  if

$$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$$

$$[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in  $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$  must be replaced by their values from 96. It is easy to see that  $\varphi$  is a decreasing function in  $G_{45}$  taking into account the hypothesis  $\varphi(0) > 0, \varphi(\infty) < 0$  it follows that there exists a unique  $G_{45}^*$  such that  $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution

511

$G_{14}^*$  given by  $\varphi(G^*) = 0, T_{14}^*$  given by  $f(T_{14}^*) = 0$  and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

$G_{17}^*$  given by  $\varphi((G_{19})^*) = 0, T_{17}^*$  given by  $f(T_{17}^*) = 0$  and

512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} , G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]}$$

513

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} , T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]}$$

514

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

515

$G_{21}^*$  given by  $\varphi((G_{23})^*) = 0, T_{21}^*$  given by  $f(T_{21}^*) = 0$  and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} , G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution

516

$G_{25}^*$  given by  $\varphi(G_{27}) = 0, T_{25}^*$  given by  $f(T_{25}^*) = 0$  and

$$G_{24}^* = \frac{(a_{24})^{(4)}G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} , G_{26}^* = \frac{(a_{26})^{(4)}G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)}T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} , T_{26}^* = \frac{(b_{26})^{(4)}T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]}$$

517

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

$G_{29}^*$  given by  $\varphi((G_{31})^*) = 0$ ,  $T_{29}^*$  given by  $f(T_{29}^*) = 0$  and

$$G_{28}^* = \frac{(a_{28})^{(5)}G_{29}^*}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} , G_{30}^* = \frac{(a_{30})^{(5)}G_{29}^*}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)}T_{29}^*}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^*)]} , T_{30}^* = \frac{(b_{30})^{(5)}T_{29}^*}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^*)]}$$
519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

$G_{33}^*$  given by  $\varphi((G_{35})^*) = 0$ ,  $T_{33}^*$  given by  $f(T_{33}^*) = 0$  and

$$G_{32}^* = \frac{(a_{32})^{(6)}G_{33}^*}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} , G_{34}^* = \frac{(a_{34})^{(6)}G_{33}^*}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)}T_{33}^*}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^*)]} , T_{34}^* = \frac{(b_{34})^{(6)}T_{33}^*}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^*)]}$$
521

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

$G_{37}^*$  given by  $\varphi((G_{39})^*) = 0$ ,  $T_{37}^*$  given by  $f(T_{37}^*) = 0$  and

$$G_{36}^* = \frac{(a_{36})^{(7)}G_{37}^*}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^*)]} , G_{38}^* = \frac{(a_{38})^{(7)}G_{37}^*}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)}T_{37}^*}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^*)]} , T_{38}^* = \frac{(b_{38})^{(7)}T_{37}^*}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

$G_{41}^*$  given by  $\varphi((G_{43})^*) = 0$ ,  $T_{41}^*$  given by  $f(T_{41}^*) = 0$  and

$$G_{40}^* = \frac{(a_{40})^{(8)}G_{41}^*}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^*)]} , G_{42}^* = \frac{(a_{42})^{(8)}G_{41}^*}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)}T_{41}^*}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^*)]} , T_{42}^* = \frac{(b_{42})^{(8)}T_{41}^*}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99 523

$G_{45}^*$  given by  $\varphi((G_{47})^*) = 0$ ,  $T_{45}^*$  given by  $f(T_{45}^*) = 0$  and A

$$G_{44}^* = \frac{(a_{44})^{(9)}G_{45}^*}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^*)]} , G_{46}^* = \frac{(a_{46})^{(9)}G_{45}^*}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)}T_{45}^*}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^*)]} , T_{46}^* = \frac{(b_{46})^{(9)}T_{45}^*}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^*)]}$$

**ASYMPTOTIC STABILITY ANALYSIS**

524

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(1)}$  and  $(b_i'')^{(1)}$  Belong to  $C^{(1)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

**Proof:**Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + G_i , T_i = T_i^* + T_i$$

$$\frac{\partial (a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})G_{13} + (a_{13})^{(1)}G_{14} - (q_{13})^{(1)}G_{13}^*T_{14} \tag{525}$$

$$\frac{dG_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})G_{14} + (a_{14})^{(1)}G_{13} - (q_{14})^{(1)}G_{14}^*T_{14} \tag{526}$$

$$\frac{dG_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})G_{15} + (a_{15})^{(1)}G_{14} - (q_{15})^{(1)}G_{15}^*T_{14} \tag{527}$$

$$\frac{dT_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})T_{13} + (b_{13})^{(1)}T_{14} + \sum_{j=13}^{15} (s_{(13)(j)})T_{13}^*G_j \tag{528}$$

$$\frac{dT_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})T_{14} + (b_{14})^{(1)}T_{13} + \sum_{j=13}^{15} (s_{(14)(j)})T_{14}^*G_j \tag{529}$$

$$\frac{dT_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})T_{15} + (b_{15})^{(1)}T_{14} + \sum_{j=13}^{15} (s_{(15)(j)})T_{15}^*G_j \tag{530}$$

**ASYMPTOTIC STABILITY ANALYSIS**

531

**Theorem 4:**If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(2)}$  and  $(b_i'')^{(2)}$  Belong to  $C^{(2)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable

**Proof:** Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \tag{532}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{533}$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{534}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{535}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{536}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(16)(j)})\mathbb{T}_{16}^*\mathbb{G}_j \tag{537}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18} (s_{(17)(j)})\mathbb{T}_{17}^*\mathbb{G}_j \tag{538}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18} (s_{(18)(j)})\mathbb{T}_{18}^*\mathbb{G}_j \tag{539}$$

**ASYMPTOTIC STABILITY ANALYSIS** 540

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(3)}$  and  $(b_i'')^{(3)}$  belong to  $C^{(3)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $\mathbb{G}_i, \mathbb{T}_i$  :-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \quad , \quad \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{541}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{542}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{543}$$

$$\frac{dT_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})T_{20} + (b_{20})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(20)(j)}T_{20}^*G_j) \quad 544$$

$$\frac{dT_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})T_{21} + (b_{21})^{(3)}T_{20} + \sum_{j=20}^{22} (s_{(21)(j)}T_{21}^*G_j) \quad 545$$

$$\frac{dT_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})T_{22} + (b_{22})^{(3)}T_{21} + \sum_{j=20}^{22} (s_{(22)(j)}T_{22}^*G_j) \quad 546$$

**ASYMPTOTIC STABILITY ANALYSIS** 547

**Theorem 4:** If the conditions of the previous theorem are satisfied and if the functions  $(a''_i)^{(4)}$  and  $(b''_i)^{(4)}$  belong to  $C^{(4)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :- 548

$$G_i = G_i^* + G_i, \quad T_i = T_i^* + T_i$$

$$\frac{\partial (a''_{25})^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)}, \quad \frac{\partial (b''_i)^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{dG_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})G_{24} + (a_{24})^{(4)}G_{25} - (q_{24})^{(4)}G_{24}^*T_{25} \quad 549$$

$$\frac{dG_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})G_{25} + (a_{25})^{(4)}G_{24} - (q_{25})^{(4)}G_{25}^*T_{25} \quad 550$$

$$\frac{dG_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})G_{26} + (a_{26})^{(4)}G_{25} - (q_{26})^{(4)}G_{26}^*T_{25} \quad 551$$

$$\frac{dT_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})T_{24} + (b_{24})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(24)(j)}T_{24}^*G_j) \quad 552$$

$$\frac{dT_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})T_{25} + (b_{25})^{(4)}T_{24} + \sum_{j=24}^{26} (s_{(25)(j)}T_{25}^*G_j) \quad 553$$

$$\frac{dT_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})T_{26} + (b_{26})^{(4)}T_{25} + \sum_{j=24}^{26} (s_{(26)(j)}T_{26}^*G_j) \quad 554$$

**ASYMPTOTIC STABILITY ANALYSIS** 555

**Theorem 5:** If the conditions of the previous theorem are satisfied and if the functions  $(a''_i)^{(5)}$  and  $(b''_i)^{(5)}$  belong to  $C^{(5)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $\mathbb{G}_i, \mathbb{T}_i$  :- 556

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}''^{(5)})}{\partial T_{29}} (T_{29}^*) = (q_{29})^{(5)} \quad , \quad \frac{\partial (b_i''^{(5)})}{\partial G_j} ((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)})T_{28}^*\mathbb{G}_j \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)})T_{29}^*\mathbb{G}_j \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)})T_{30}^*\mathbb{G}_j \quad 562$$

**ASYMPTOTIC STABILITY ANALYSIS** 563

**Theorem 6:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i''^{(6)})$  and  $(b_i''^{(6)})$  belong to  $C^{(6)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $\mathbb{G}_i, \mathbb{T}_i$  :- 564

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}''^{(6)})}{\partial T_{33}} (T_{33}^*) = (q_{33})^{(6)} \quad , \quad \frac{\partial (b_i''^{(6)})}{\partial G_j} ((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^* \mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34} (s_{(32)(j)})T_{32}^* \mathbb{G}_j \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34} (s_{(33)(j)})T_{33}^* \mathbb{G}_j \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34} (s_{(34)(j)})T_{34}^* \mathbb{G}_j \quad 570$$

**ASYMPTOTIC STABILITY ANALYSIS** 571

**Theorem 7:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(7)}$  and  $(b_i'')^{(7)}$  belong to  $C^{(7)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $\mathbb{G}_i, \mathbb{T}_i$  :- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \quad , \quad \frac{\partial (b_i'')^{(7)}}{\partial G_j}((G_{39})^{**}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -((a'_{36})^{(7)} + (p_{36})^{(7)})\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^* \mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -((a'_{37})^{(7)} + (p_{37})^{(7)})\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^* \mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -((a'_{38})^{(7)} + (p_{38})^{(7)})\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^* \mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -((b'_{36})^{(7)} - (r_{36})^{(7)})\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38} (s_{(36)(j)})T_{36}^* \mathbb{G}_j \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -((b'_{37})^{(7)} - (r_{37})^{(7)})\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38} (s_{(37)(j)})T_{37}^* \mathbb{G}_j \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -((b'_{38})^{(7)} - (r_{38})^{(7)})\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38} (s_{(38)(j)})T_{38}^* \mathbb{G}_j \quad 579$$

Obviously, these values represent an equilibrium solution

**ASYMPTOTIC STABILITY ANALYSIS**

**Theorem 8:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(8)}$  and  $(b_i'')^{(8)}$  belong to  $C^{(8)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :- 580

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)}, \quad \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -((a'_{40})^{(8)} + (p_{40})^{(8)})\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \tag{581}$$

$$\frac{d\mathbb{G}_{41}}{dt} = -((a'_{41})^{(8)} + (p_{41})^{(8)})\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \tag{582}$$

$$\frac{d\mathbb{G}_{42}}{dt} = -((a'_{42})^{(8)} + (p_{42})^{(8)})\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \tag{583}$$

$$\frac{d\mathbb{T}_{40}}{dt} = -((b'_{40})^{(8)} - (r_{40})^{(8)})\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42} (s_{(40)(j)})T_{40}^*\mathbb{G}_j \tag{584}$$

$$\frac{d\mathbb{T}_{41}}{dt} = -((b'_{41})^{(8)} - (r_{41})^{(8)})\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42} (s_{(41)(j)})T_{41}^*\mathbb{G}_j \tag{585}$$

$$\frac{d\mathbb{T}_{42}}{dt} = -((b'_{42})^{(8)} - (r_{42})^{(8)})\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42} (s_{(42)(j)})T_{42}^*\mathbb{G}_j \tag{586}$$

**ASYMPTOTIC STABILITY ANALYSIS**

586  
A

**Theorem 9:** If the conditions of the previous theorem are satisfied and if the functions  $(a_i'')^{(9)}$  and  $(b_i'')^{(9)}$  belong to  $C^{(9)}(\mathbb{R}_+)$  then the above equilibrium point is asymptotically stable.

Proof: Denote

**Definition of**  $G_i, T_i$  :-

$$G_i = G_i^* + \mathbb{G}_i, \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)}, \quad \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{dG_{44}}{dt} = -((a'_{44})^{(9)} + (p_{44})^{(9)})G_{44} + (a_{44})^{(9)}G_{45} - (q_{44})^{(9)}G_{44}^*T_{45} \quad 586 \text{ B}$$

$$\frac{dG_{45}}{dt} = -((a'_{45})^{(9)} + (p_{45})^{(9)})G_{45} + (a_{45})^{(9)}G_{44} - (q_{45})^{(9)}G_{45}^*T_{45} \quad 586 \text{ C}$$

$$\frac{dG_{46}}{dt} = -((a'_{46})^{(9)} + (p_{46})^{(9)})G_{46} + (a_{46})^{(9)}G_{45} - (q_{46})^{(9)}G_{46}^*T_{45} \quad 586 \text{ D}$$

$$\frac{dT_{44}}{dt} = -((b'_{44})^{(9)} - (r_{44})^{(9)})T_{44} + (b_{44})^{(9)}T_{45} + \sum_{j=44}^{46} (s_{(44)(j)})T_{44}^*G_j \quad 586 \text{ E}$$

$$\frac{dT_{45}}{dt} = -((b'_{45})^{(9)} - (r_{45})^{(9)})T_{45} + (b_{45})^{(9)}T_{44} + \sum_{j=44}^{46} (s_{(45)(j)})T_{45}^*G_j \quad 586 \text{ F}$$

$$\frac{dT_{46}}{dt} = -((b'_{46})^{(9)} - (r_{46})^{(9)})T_{46} + (b_{46})^{(9)}T_{45} + \sum_{j=44}^{46} (s_{(46)(j)})T_{46}^*G_j \quad 586 \text{ G}$$

**The characteristic equation of this system is** 587

$$\begin{aligned} & ((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}) \\ & \left[ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right] \\ & \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \right) \\ & + \left( ((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \right) \\ & \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \right) \\ & \left( ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)})(\lambda)^{(1)} \right) \\ & \left( ((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)})(\lambda)^{(1)} \right) \\ & + \left( ((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)})(\lambda)^{(1)} \right)(q_{15})^{(1)}G_{15} \\ & + ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^* \\ & \left. \left( ((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \right) \right\} = 0 \end{aligned}$$

+

$$((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)})$$

$$\begin{aligned}
 & \left[ \left( (\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)} \right) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right] \\
 & \left( (\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)} \right) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \\
 & + \left( (\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)} \right) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \\
 & \left( (\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)} \right) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \\
 & \left( (\lambda)^{(2)} \right)^2 + \left( (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & \left( (\lambda)^{(2)} \right)^2 + \left( (b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left( (\lambda)^{(2)} \right)^2 + \left( (a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + \left( (\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)} \right) \left( (a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^* \right) \\
 & \left. \left( (\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)} \right) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & \left( (\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)} \right) \left\{ (\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)} \right\} \\
 & \left[ \left( (\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)} \right) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \\
 & \left( (\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \\
 & + \left( (\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)} \right) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(3)} G_{21}^* \\
 & \left( (\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \\
 & \left( (\lambda)^{(3)} \right)^2 + \left( (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left( (\lambda)^{(3)} \right)^2 + \left( (b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left( (\lambda)^{(3)} \right)^2 + \left( (a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + \left( (\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)} \right) \left( (a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^* \right) \\
 & \left. \left( (\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)})\{((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)}) \\
 & [(((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)})(q_{25})^{(4)}G_{25}^* + (a_{25})^{(4)}(q_{24})^{(4)}G_{24}^*)] \\
 & ((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)})s_{(25),(25)}T_{25}^* + (b_{25})^{(4)}s_{(24),(25)}T_{25}^* \\
 & + (((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)})(q_{24})^{(4)}G_{24}^* + (a_{24})^{(4)}(q_{25})^{(4)}G_{25}^*) \\
 & ((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)})s_{(25),(24)}T_{25}^* + (b_{25})^{(4)}s_{(24),(24)}T_{24}^* \\
 & (((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)})(\lambda)^{(4)}) \\
 & (((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)})(\lambda)^{(4)}) \\
 & + (((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)})(\lambda)^{(4)})(q_{26})^{(4)}G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)})(a_{26})^{(4)}(q_{25})^{(4)}G_{25}^* + (a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G_{24}^* \\
 & ((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)})s_{(25),(26)}T_{25}^* + (b_{25})^{(4)}s_{(24),(26)}T_{24}^* \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)})\{((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & [(((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)})(q_{29})^{(5)}G_{29}^* + (a_{29})^{(5)}(q_{28})^{(5)}G_{28}^*)] \\
 & ((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})s_{(29),(29)}T_{29}^* + (b_{29})^{(5)}s_{(28),(29)}T_{29}^* \\
 & + (((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)})(q_{28})^{(5)}G_{28}^* + (a_{28})^{(5)}(q_{29})^{(5)}G_{29}^*) \\
 & ((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})s_{(29),(28)}T_{29}^* + (b_{29})^{(5)}s_{(28),(28)}T_{28}^* \\
 & (((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)})(\lambda)^{(5)}) \\
 & (((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)})(\lambda)^{(5)}) \\
 & + (((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)})(\lambda)^{(5)})(q_{30})^{(5)}G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)})(a_{30})^{(5)}(q_{29})^{(5)}G_{29}^* + (a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G_{28}^* \\
 & ((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})s_{(29),(30)}T_{29}^* + (b_{29})^{(5)}s_{(28),(30)}T_{28}^* \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)})\{((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & [((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)})(q_{33})^{(6)}G_{33}^* + (a_{33})^{(6)}(q_{32})^{(6)}G_{32}^*] \\
 & ((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(33)}T_{33}^* + (b_{33})^{(6)}s_{(32),(33)}T_{33}^* \\
 & + ((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)})(q_{32})^{(6)}G_{32}^* + (a_{32})^{(6)}(q_{33})^{(6)}G_{33}^* \\
 & ((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(32)}T_{33}^* + (b_{33})^{(6)}s_{(32),(32)}T_{32}^* \\
 & (((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)})(\lambda)^{(6)}) \\
 & (((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)})(\lambda)^{(6)}) \\
 & + (((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)})(\lambda)^{(6)})(q_{34})^{(6)}G_{34} \\
 & + ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)})(a_{34})^{(6)}(q_{33})^{(6)}G_{33}^* + (a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G_{32}^* \\
 & ((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(34)}T_{33}^* + (b_{33})^{(6)}s_{(32),(34)}T_{32}^* \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)})\{((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)}) \\
 & [((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)})(q_{37})^{(7)}G_{37}^* + (a_{37})^{(7)}(q_{36})^{(7)}G_{36}^*] \\
 & ((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(37)}T_{37}^* + (b_{37})^{(7)}s_{(36),(37)}T_{37}^* \\
 & + ((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)})(q_{36})^{(7)}G_{36}^* + (a_{36})^{(7)}(q_{37})^{(7)}G_{37}^* \\
 & ((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(36)}T_{37}^* + (b_{37})^{(7)}s_{(36),(36)}T_{36}^* \\
 & (((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)})(\lambda)^{(7)}) \\
 & (((\lambda)^{(7)})^2 + ((b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)})(\lambda)^{(7)}) \\
 & + (((\lambda)^{(7)})^2 + ((a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)})(\lambda)^{(7)})(q_{38})^{(7)}G_{38} \\
 & + ((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)})(a_{38})^{(7)}(q_{37})^{(7)}G_{37}^* + (a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G_{36}^* \\
 & ((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(38)}T_{37}^* + (b_{37})^{(7)}s_{(36),(38)}T_{36}^* \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)})\{((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)}) \\
 & [((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)})(q_{41})^{(8)}G_{41}^* + (a_{41})^{(8)}(q_{40})^{(8)}G_{40}^*] \\
 & ((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)})s_{(41),(41)}T_{41}^* + (b_{41})^{(8)}s_{(40),(41)}T_{41}^* \\
 & + ((\lambda)^{(8)} + (a'_{41})^{(8)} + (p_{41})^{(8)})(q_{40})^{(8)}G_{40}^* + (a_{40})^{(8)}(q_{41})^{(8)}G_{41}^* \\
 & ((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)})s_{(41),(40)}T_{41}^* + (b_{41})^{(8)}s_{(40),(40)}T_{40}^* \\
 & (((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)})(\lambda)^{(8)}) \\
 & (((\lambda)^{(8)})^2 + ((b'_{40})^{(8)} + (b'_{41})^{(8)} - (r_{40})^{(8)} + (r_{41})^{(8)})(\lambda)^{(8)}) \\
 & + (((\lambda)^{(8)})^2 + ((a'_{40})^{(8)} + (a'_{41})^{(8)} + (p_{40})^{(8)} + (p_{41})^{(8)})(\lambda)^{(8)})(q_{42})^{(8)}G_{42} \\
 & + ((\lambda)^{(8)} + (a'_{40})^{(8)} + (p_{40})^{(8)})(a_{42})^{(8)}(q_{41})^{(8)}G_{41}^* + (a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G_{40}^* \\
 & ((\lambda)^{(8)} + (b'_{40})^{(8)} - (r_{40})^{(8)})s_{(41),(42)}T_{41}^* + (b_{41})^{(8)}s_{(40),(42)}T_{40}^* \} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(9)} + (b'_{46})^{(9)} - (r_{46})^{(9)})\{((\lambda)^{(9)} + (a'_{46})^{(9)} + (p_{46})^{(9)}) \\
 & [((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)})(q_{45})^{(9)}G_{45}^* + (a_{45})^{(9)}(q_{44})^{(9)}G_{44}^*] \\
 & ((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)})s_{(45),(45)}T_{45}^* + (b_{45})^{(9)}s_{(44),(45)}T_{45}^* \\
 & + ((\lambda)^{(9)} + (a'_{45})^{(9)} + (p_{45})^{(9)})(q_{44})^{(9)}G_{44}^* + (a_{44})^{(9)}(q_{45})^{(9)}G_{45}^* \\
 & ((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)})s_{(45),(44)}T_{45}^* + (b_{45})^{(9)}s_{(44),(44)}T_{44}^* \\
 & (((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)})(\lambda)^{(9)}) \\
 & (((\lambda)^{(9)})^2 + ((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)})(\lambda)^{(9)}) \\
 & + (((\lambda)^{(9)})^2 + ((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)})(\lambda)^{(9)})(q_{46})^{(9)}G_{46} \\
 & + ((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)})(a_{46})^{(9)}(q_{45})^{(9)}G_{45}^* + (a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G_{44}^* \\
 & ((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)})s_{(45),(46)}T_{45}^* + (b_{45})^{(9)}s_{(44),(46)}T_{44}^* \} = 0
 \end{aligned}$$

**And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.**

