

Nanoparticles as Alternative to Pesticides in Management Plant Diseases - A Review

AbdulHameed.M. Al-Samarrai

Department of Biological, Collage of Education, University Samarraa, Iraq

Abstract- For the past few decades, there has been a considerable in research interest in the area of natural product delivery using particulate for controlling plant pathology. The secondary metabolites in plants have been used in the formulation of nanoparticles through increase the effectiveness of therapeutic compounds used to reduce the spread of plant diseases, while minimizing side effects for being: rich source of bioactive chemicals, biodegradable in nature and non-polluting (eco-friendly). Particulate systems like Nanoparticles have been used a physical approach to alter and improve the effective to properties of some types of synthetic chemical pesticides or in the production of bio-pesticides directly. Here, we review various aspects of nanoparticles formulation, characterization, effect of their characteristics and their applications in management plant diseases.

Index Terms- nanoparticles, natural product, secondary metabolites, synthetic chemicals, management disease

I. INTRODUCTION

The need for biosynthesis of Nanoparticles rose as the physical and chemical processes were costly. So in the search of for cheaper pathways for nanoparticle synthesis, scientists used microorganisms and then plant extracts for synthesis. Nature has devised various processes for the synthesis of nano- and micro- length scaled inorganic materials which have contributed to the development of relatively new and largely unexplored area of research based on the biosynthesis of nonmaterial's (Mohanpuria et al., 2007). Biosynthesis of Nanoparticles is a kind of bottom up approach where the main reaction occurring is reduction/oxidation. The microbial enzymes or the plant phytochemicals with anti oxidant or reducing properties are usually responsible for reduction of metal compounds into their respective Nanoparticles.

The three main steps in the preparation of Nanoparticles that should be evaluated from a green chemistry perspective are the choice of the solvent medium used for the synthesis, the Biomimetic Synthesis of Nanoparticles: Science, Technology & Applicability 5 choice of an environmentally benign reducing agent and the choice of a non toxic material for the stabilization of the Nanoparticles. Most of the synthetic methods reported to date rely heavily on organic solvents. This is mainly due to the hydrophobicity of the capping agents used (Raveendran et al., 2002). Synthesis using bio-organisms is compatible with the green chemistry principles: the bio-organism is (i) eco-friendly as

are (ii) the reducing agent employed and (iii) the capping agent in the reaction (Li et al., 2007). These papers come to highlight on Bioactive Nanoparticles of plant which can be effective alternatives to from plant secondary metabolites as botanical biopesticides.

II. USE OF PLANTS TO SYNTHESIZE NANOPARTICALS

The advantage of using plants for the synthesis of nanoparticle is that they are easily available, safe to handle and possess a broad variability of metabolites that may aid in reduction. A number of plants are being currently investigated for their role in the of nanoparticle. Gold Nanoparticles with a size range of 2-20 nm have been synthesized using the live alfa alfa plants (Torresday et al., 2002). Nanoparticles of silver, nickel, cobalt, zinc and copper have also been synthesized inside the live plants of *Brassica juncea* (Indian mustard), *Medicago sativa* (Alfa) and *Heliantus annus* (Sunflower). Certain plants are known to accumulate higher concentrations of metals compared to others and such plants are termed as hyper accumulators. Of the plants investigated, *Brassica juncea* had better metal accumulating ability and later assimilating it as nanoparticle (Bali et al., 2006). Recently much work has been done with regard to plant assisted reduction of metal nanoparticle and the respective role of phytochemicals. The main phytochemicals responsible have been identified as terpenoids, flavones, ketones, aldehydes, amides and carboxylic acids in the light of IR spectroscopic studies. The main water soluble phytochemicals are flavones, organic acids and quinones which are responsible for immediate reduction.

The phytochemicals present in *Bryophyllum sp.* (Xerophytes), *Cyprus sp.* (Mesophytes) and *Hydrilla sp.* (Hydrophytes) were studied for their role in the synthesis of silver nanoparticle. The Xerophytes were found to contain emodin, an anthraquinone which could undergo redial tautomerization leading to the formation of silver Nanoparticles. The Mesophytes studied contained three types of benzoquinones, namely, cyperoquinone, dietchequinone and remirin. It was suggested that gentle warming followed by subsequent incubation resulted in the activation of quinones leading to particle size reduction. Catechol and protocatechaldehyde were reported in the hydrophyte studied along with other phytochemicals. It was reported that catechol under alkaline conditions gets transformed into protocatechaldehyde and finally into protocatecheuc acid. Both these processes liberated hydrogen and it was suggested that it played a role in the synthesis of thenanoparticles. The size of the Nanoparticles synthesized using xerophytes, mesophytes and hydrophytes were in the range of 2- 5nm (Jha et al., 2009).

Recently gold nanoparticles have been synthesized using the extracts of *Magnolia kobus* and *Diopyros kaki* leaf extracts. The effect of temperature on nanoparticle formation was investigated and it was reported that polydisperse particles with a size range of 5- 300nm was obtained at lower temperature while a higher temperature supported the formation of smaller and spherical particles (Song *et al.*, 2009). While fungi and bacteria require a comparatively longer incubation time for the reduction of metal ions, water soluble phytochemicals do it in a much lesser time.

Therefore compared to biomimetic synthesis of Nanoparticles: Science, Technology & Applicability 9 bacteria and fungi, plants are better candidates for the synthesis of Nanoparticles. Taking use of plant tissue culture techniques and downstream processing procedures, it is possible to synthesize metallic as well as oxide Nanoparticles on an industrial scale once issues like the metabolic status of the plant etc. are properly addressed.

III. SYNTHESIS OF NANOPARTICLES IN PLANT

Studies indicated that the reducing phytochemicals in the neem leaf consisted mainly of terpenoids. It was found that these reducing components also served as capping and stabilizing agents in addition to reduction as revealed from FT IR studies. The major advantage of using the neem leaves is that it is a commonly available medicinal plant and the antibacterial such best, bacteria and fungi in plant activity of the biosynthesized silver nanoparticle might have been enhanced as it was capped with the neem leaf extract. Biomimetic Synthesis of Nanoparticles. The major chemical constituents in the extract were identified as nimbin and quercetin this compound possible to be anti-pathological causes of plants. (Shankar *et al.*, 2004, Tripathy *et al.*, 2009). Study is because of the Increase in new resistant strains of insect, bacteria and fungi against most potent antibiotics. Has promoted research in the well known activity compounds, including Nanoparticles and was more pronounced against pest than organisms.

IV. APPLICATION OF NANOPARTICLES AGAINST ANTI-MICROBIOLOGY IN PLANTS

Silver Nanoparticles have shown promise against gram positive *S. aureus*. As Metal nanoparticle embedded paints have been synthesized using vegetable oils and have been found to have good ant microbiology activity (Kumar *et al.*, 2008).

The use of nano-sized silver particles as antimicrobial agents has become more common as technological advances make their production more economical. One of the potential applications in which silver can be utilized is in management of plant diseases. Since silver displays multiple modes of inhibitory action to microorganisms (Clement and Jarret 1994), it may be used for controlling various plant pathogens in a relatively safer way compared to synthetic fungicides (Kim *et al.*, 2006). Until now, limited research provided some evidence of the applicability of silver for controlling plant diseases (Kim *et al.*, 2006).





Fig. 1: Colony formation of *Bipolaris sorokiniana* on potato dextrose agar medium, 1 h post-treatment of spores with water, AgNO₃, AgCl, silver Nanoparticles [Ag(p)] and electrochemical silver [Ag(e)]. Images were captured after culture plates had been incubated for 6 days at 25°C.

(Kim *et al.*, 2009).

Various forms of silver ions and Nanoparticles were tested by Kim *et al.* (2009) in the current study to examine the antifungal activity on two plant-pathogenic fungi, *Bipolaris sorokiniana* and *Magnaporthe grisea*. In vitro petri dish assays indicated that silver ions and Nanoparticles had a significant effect on the colony formation of these two pathogens. Effective concentrations of the silver compounds inhibiting colony formation by 50% (EC₅₀) were higher for *B. sorokiniana* than for *M. grisea*. The inhibitory effect on colony formation significantly diminished after silver cations were neutralized with chloride ions. Growth chamber inoculation assays further confirmed that both ionic and nanoparticle silver significantly reduced these two fungal diseases on perennial ryegrass (*Lolium perenne*). Particularly, silver ions and Nanoparticles effectively reduced disease severity with an application at 3 h before spore inoculation, but their efficacy significantly diminished when applied at 24 h after inoculation. The in vitro and in planta evaluations of silver indicated that both silver ions and Nanoparticles influence colony formation of spores and disease progress of plant-pathogenic fungi. In planta efficacy of silver ions and Nanoparticles is much greater with preventative application, which may promote the direct contact of silver with spores and germ tubes, and inhibit their viability.



Marek *et al* (2010) studied a significant reduction in mycelial growth was observed for spores incubated with silver Nanoparticles. The sporulation test showed that, relative to control samples, the number of spores formed by mycelia increased in the culture after contact with silver Nanoparticles, especially on the nutrient-poor PDA medium. The 24 h incubation of FC spores with a 2.5 ppm solution of silver Nanoparticles greatly reduced the number of germinating fragments and sprout length relative to the control.

Silver nanoparticles (WA-CV-WA13B) at various concentrations were applied before and after disease outbreak in plants to determine antifungal activities. In the field tests, the application of 100 ppm silver nanoparticles showed the highest inhibition rate for both before and after the outbreak of disease on cucumbers and pumpkins. Also, the application of 100 ppm silver nanoparticles showed maximum inhibition for the growth of fungal hyphae and conidial germination in *in vivo* tests. Scanning electron microscope results indicated that the silver nanoparticles caused detrimental effects on both mycelial growth and conidial germination (Kabir *et al*, 2010).

The antibacterial potentiality of zinc oxide (ZnO) nanoparticles (NPs), compared with conventional ZnO powder, against nine bacterial strains, mostly foodborne including pathogens, was evaluated using qualitative and quantitative assays. ZnO NP was more efficient as antibacterial agent than powder. Gram-positive bacteria were generally more sensitive to ZnO than Gram negatives. The exposure of *Salmonella typhimurium* and *Staphylococcus aureus* to their relevant minimal inhibitory concentrations from ZnO NP reduced the cell number to zero within 8 and 4 h, respectively. Scanning electron micrographs of the treated bacteria with NPs exhibited that the disruptive effect of ZnO on *S. aureus* was vigorous as all treated cells were completely exploded or lysed after only 4h from exposure. Promising results of ZnO NP antibacterial activity suggest its usage in food systems as preservative agent after further required investigations and risk assessments (Ahamed *et al*;2011).

In study was planning to evaluate the potential of nano silver (NS) and nano dioxide titanium (TiO₂) to remove bacterial contaminants. Experiment involved NS and TiO₂ in MS medium. Tobacco explants were cultured on modified MS medium and evaluate after four weeks. This research shows that NS and TiO₂ had a good potential for removing of the

bacterial contaminants in tobacco plant tissue culture procedures (Kamran *et al*; 2011).

V. COLLOCATION

Use Nano particles for delivery of anti-microbiological or drug molecules will be at its helm in near future for therapy of all pathological sufferings of plants. There are myriad of Nanomaterials including polymeric nanoparticles, iron oxide nanoparticles, gold nanoparticles and silver ion which can be easily synthesized and exploited as pesticide. Antimicrobial activity of nano particles and after confirm the ability of nano to reduce the microorganism, we decide to using and adding NS to MS tissue culture media and the application of nanotechnology in the chemical and bio-based industries besides making a special reference on its usage in the nanoparticle formulations from plant secondary metabolites as botanical biopesticides .

REFERENCES

- [1] Ahmed A. T, Wael F. T, Shaaban. M, A, Mohammed F. S. (2001). Antibacterial action of zinc oxide nanoparticle agents' foodborne pathogens. *Journal of Food Safety*: 31, 2, pages 211-218, May 2011
- [2] Bali, R.; Razak, N.; Lumb, A. & Harris, A.T. (2006). The synthesis of metal nanoparticles inside live plants. *IEEE Xplore* DOI 10.1109/ICONN.2006.340592.
- [3] Clement, J. L., and Jarret, P. S. 1994. Antimicrobial silver. *Metal-Based Drugs* 1:467-482. Park, H.-J., Kim, S. H., Kim, H. J., and Choi, S.-H. 2006. A new composition of nanosized silica-silver for control of various plant diseases. *Plant Pathol. J.* 22:295-302.
- [4] 4-Jha, A.K.; Prasad, K.; Prasad, K. & Kulkarni, A.R. (2009). Plant system: Nature's nanofactory. *Colloids and Surfaces B: Biointerfaces* (inpress) doi: 10.1016/j.colsurfb.2009.05.018.
- [5] Li, S.; Shen, Y.; Xie, A.; Yu, X.; Qui, L.; Zhang, L. & Zhang, Q. (2007). Green synthesis of silver nanoparticles using *Capsicum annum* L. extract. *Green Chemistry*. 9. 852-858
- [6] Kabir Lamsall, Sang-Woo Kim¹, Jin Hee Jung¹, Yun Seok Kim¹, Kyoung Su Kim² and Youn Su Lee.(2010). Inhibition Effects of Silver Nanoparticles against Powdery Mildews on Cucumber and Pumpkin. *Microbiology* 39(1): 26-32 (2011).
- [7] Kamran. S., Forogh .M, Mahtab. E and Mohammad. (2011). *In Vitro* Antibacterial Activity of Nanomaterials for using in Tobacco Plants Tissue Culture. *World Academy of Science, Engineering and Technology* 79:372-373.
- [8] Marek Jan Kasproiewicz, Magdalena Koziół, and Anna Gorczyca The effect of silver nanoparticles on phytopathogenic spores of *Fusarium culmorum*. *Canadian Journal of Microbiology* Jan 2010, Volume 56, Number 3: 247-253.
- [9] Mohanpuria, P.; Rana, K.N. & Yadav, S.K (2008). Biosynthesis of nanoparticles: technological concepts and future applications. *Journal of Nanoparticle Research* 10.; 507- 517.
- [10] Raveendran, P.; Fu, J. & Wallen., S.L. (2003). Completely "Green" Synthesis and Stabilization of metal nanoparticles. *Journal of American Chemical Society*, 125(46); 13940-13941.
- [11] Shankar, S.S.; Rai, A.; Ahmad, A. & Sastry, M. (2004). Rapid synthesis of Au, Ag and bimetallic Au core- Ag shell nanoparticles using neem (*Azadirachta indica*) leaf broth. *Journal of colloid and interface science*. 275; 496-502.
- [12] Song, J.Y.; Jang, H.K. & Kim, B.S. (2009). Biological synthesis of gold nanoparticles using *Magnolia kobus* and *Diopyros kaki* leaf extracts. *Process Biochemistry* 44.; 1133- 1138.
- [13] Young-Ki Jo, Byung H. Kim, Geunhwa Jung. 2009. Antifungal Activity of Silver Ions and Nanoparticles on Phytopathogenic Fungi. *The American Phytopathological Society*.1037-1042. Vol. 93 No. 10.
- [14] Tripathy, A.; Raichur, A.M.; Chandrasekaran, N.; Prathna, T.C. & Mukherjee, A. (2009). Process variables in biomimetic synthesis of silver nanoparticles by aqueous extract of *Azadirachta indica* (Neem) leaves. *Journal of Nanoparticle Research*. Article in press.

AUTHORS

AbdulHameed.M. Alasmarrai, Department of Biological, Collage
of Education, University Samarraa.

Email id: thmoon_83@yahoo.com