

SOME CONTRIBUTIONS TO YANG MILLS THEORY FORTIFICATION –DISSIPATION MODELS

¹DR K N PRASANNA KUMAR, ²PROF B S KIRANAGI AND ³PROF C S BAGEWADI

ABSTRACT. We provide a series of Models for the problems that arise in Yang Mills Theory. No claim is made that the problem is solved. We do factorize the Yang Mills Theory and give a Model for the values of LHS and RHS of the yang Mills theory. We hope these forms the stepping stone for further factorizations and solutions to the subatomic denominations at Planck's scale. Work also throws light on some important factors like mass acquisition by symmetry breaking, relation between strong interaction and weak interaction, Lagrangian Invariance despite transformations, Gauge field, Noncommutative symmetry group of Gauge Theory and Yang Mills Theory itself.

We take in to consideration the following parameters, processes and concepts:

- (1) Acquisition of mass
- (2) Symmetry Breaking
- (3) Strong interaction
- (4) Unified Electroweak interaction
- (5) Continuous group of local transformations
- (6) Lagrangian Variance
- (7) Group generator in Gauge Theory
- (8) Vector field or Gauge field
- (9) Non commutative symmetry group in Gauge Theory
- (10) Yang Mills Theory (We repeat the same Bank's example. Individual debits and Credits are conservative so also the holistic one. Generalized theories are applied to various systems which are parameterized. And we live in 'measurement world'. Classification is done on the parameters of various systems to which the Theory is applied.).
- (11) First Term of the Lagrangian of the Yang Mills Theory(LHS)

$$\mathcal{L}_{gf} = -\frac{1}{4} \text{Tr}(F^2) = -\frac{1}{4} F^{\mu\nu\rho\sigma} F_{\mu\nu\rho\sigma}$$

- (12) RHS of the Yang Mills Theory

$$\mathcal{L}_{gf} = -\frac{1}{4} \text{Tr}(F^2) = -\frac{1}{4} F^{\mu\nu\rho\sigma} F_{\mu\nu\rho\sigma}$$

SYMMETRY BREAKING AND ACQUISITION OF MASS:

MODULE NUMBERED ONE

NOTATION :

G_{13} : CATEGORY ONE OF SYMMETRY BREAKING

G_{14} : CATEGORY TWO OF SYMMETRY BREAKING

G_{15} : CATEGORY THREE OF SYMMETRY BREAKING

T_{13} : CATEGORY ONE OF ACQUISITION OF MASS

T_{14} : CATEGORY TWO OF ACQUISITION OF MASS

T_{15} :CATEGORY THREE OF ACQUISITION OF MASS

UNIFIED ELECTROWEAK INTERACTION AND STRONG INTERACTION:

MODULE NUMBERED TWO:

G_{16} : CATEGORY ONE OF UNIFIED ELECTROWEAK INTERACTION

G_{17} : CATEGORY TWO OF UNIFIED ELECTROWEAK INTERACTION

G_{18} : CATEGORY THREE OF UNIFIED ELECTROWEAK INTERACTION

T_{16} :CATEGORY ONE OF STRONG INTERACTION

T_{17} : CATEGORY TWO OF STRONG INTERACTION

T_{18} : CATEGORY THREE OF STRONG INTERACTION

LAGRANGIAN INVARIANCE AND CONTINUOUS GROUP OF LOCAL TRANSFORMATIONS:

MODULE NUMBERED THREE:

G_{20} : CATEGORY ONE OF CONTINUOUS GROUP OF LOCAL TRANSFORMATIONS

G_{21} :CATEGORY TWO OF CONTINUOUS GROUP OF LOCAL TRANSFORMATIONS

G_{22} : CATEGORY THREE OF CONTINUOUS GROUP OF LOCAL TRANSFORMATION

T_{20} : CATEGORY ONE OF LAGRANGIAN INVARIANCE

T_{21} :CATEGORY TWO OF LAGRANGIAN INVARIANCE

T_{22} : CATEGORY THREE OF LAGRANGIAN INVARIANCE

GROUP GENERATOR OF GAUGE THEORY AND VECTOR FIELD(GAUGE FIELD):

: MODULE NUMBERED FOUR:

G_{24} : CATEGORY ONE OF GROUP GENERATOR OF GAUGE THEORY

G_{25} : CATEGORY TWO OF GROUP GENERATOR OF GAUGE THEORY

G_{26} : CATEGORY THREE OF GROUP GENERATOR OF GAUGE THEORY

T_{24} :CATEGORY ONE OF VECTOR FIELD NAMELY GAUGE FIELD

T_{25} :CATEGORY TWO OF GAUGE FIELD

T_{26} : CATEGORY THREE OF GAUGE FIELD

YANG MILLS THEORY AND NON COMMUTATIVE SYMMETRY GROUP IN GAUGE THEORY:

MODULE NUMBERED FIVE:

=====
=====
 G_{28} : CATEGORY ONE OF NON COMMUTATIVE SYMMETRY GROUP OF GAUGE THEORY

G_{29} : CATEGORY TWO OF NON COMMUTATIVE SYMMETRY GROUP OF GAUGE THEORY

G_{30} :CATEGORY THREE OF NON COMMUTATIVE SYMMETRY GROUP OF GAUGE THEORY

T_{28} : CATEGORY ONE OF YANG MILLS THEORY (Theory is applied to various subatomic particle systems and the classification is done based on the parametrization of these systems. There is not a single system known which is not characterized by some properties)

T_{29} :CATEGORY TWO OF YANG MILLS THEORY

T_{30} :CATEGORY THREE OF YANG MILLS THEORY

LHS OF THE YANG MILLS THEORY AND RHS OF THE YANG MILLS THEORY.TAKEN TO THE OTHER SIDE THE LHS WOULD DISSIPATE THE RHS WITH OR WITHOUT TIME LAG :

MODULE NUMBERED SIX:

$$\mathcal{L}_{gf} = -\frac{1}{4} \text{Tr}(F^2) = -\frac{1}{4} F^{\mu\nu a} F_{\mu\nu}^a$$

=====
=====
 G_{32} : CATEGORY ONE OF LHS OF YANG MILLS THEORY

G_{33} : CATEGORY TWO OF LHS OF YANG MILLS THEORY

G_{34} : CATEGORY THREE OF LHS OF YANG MILLS THEORY

T_{32} : CATEGORY ONE OF RHS OF YANG MILLS THEORY

T_{33} : CATEGORY TWO OF RHS OF YANG MILLS THEORY

T_{34} : CATEGORY THREE OF RHS OF YANG MILLS THEORY (Theory applied to various characterized systems and the systemic characterizations form the basis for the formulation of the classification).

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}, (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
 $(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}; (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}, (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
 $(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
 $(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$

are Accentuation coefficients

$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
 $(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}; (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
 $(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
 $(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}, (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$

are Dissipation coefficients

SYMMETRY BREAKING AND ACQUISITION OF MASS:

1

MODULE NUMBERED ONE

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)]G_{13} \quad 2$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)]G_{14} \quad 3$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)]G_{15} \quad 4$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)]T_{13} \quad 5$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)]T_{14} \quad 6$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)]T_{15} \quad 7$$

$$+(a''_{13})^{(1)}(T_{14}, t) = \text{First augmentation factor} \quad 8$$

$$-(b''_{13})^{(1)}(G, t) = \text{First detritions factor}$$

UNIFIED ELECTROWEAK INTERACTION AND STRONG INTERACTION:

9

MODULE NUMBERED TWO

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)]G_{16} \quad 10$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)]G_{17} \quad 11$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)]G_{18} \quad 12$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)]T_{16} \quad 13$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)]T_{17} \quad 14$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)]T_{18} \quad 15$$

$$+(a''_{16})^{(2)}(T_{17}, t) = \text{First augmentation factor} \quad 16$$

$$-(b''_{16})^{(2)}((G_{19}), t) = \text{First detritions factor} \quad 17$$

LAGRANGIAN INVARIANCE AND CONTINOUS GROUP OF LOCAL TRANSFORMATIONS: 18

MODULE NUMBERED THREE

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)]G_{20} \quad 19$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)]G_{21} \quad 20$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)]G_{22} \quad 21$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)]T_{20} \quad 22$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)]T_{21} \quad 23$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)]T_{22} \quad 24$$

$$+(a''_{20})^{(3)}(T_{21}, t) = \text{First augmentation factor}$$

$$-(b''_{20})^{(3)}(G_{23}, t) = \text{First detritions factor} \quad 25$$

26

GROUP GENERATOR OF GAUGE THEORY AND VECTOR FIELD(GAUGE FIELD):

: MODULE NUMBERED FOUR:

=====

==

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)]G_{24} \quad 27$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)]G_{25} \quad 28$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)]G_{26} \quad 29$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)]T_{24} \quad 30$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)]T_{25} \quad 31$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)]T_{26} \quad 32$$

$$+(a''_{24})^{(4)}(T_{25}, t) = \text{First augmentation factor} \quad 33$$

$$-(b''_{24})^{(4)}((G_{27}), t) = \text{First detritions factor} \quad 34$$

YANG MILLS THEORY AND NON COMMUTATIVE SYMMETRY GROUP IN GAUGE THEORY: 35

MODULE NUMBERED FIVE

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)]G_{28} \quad 36$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)]G_{29} \quad 37$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)]G_{30} \quad 38$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)]T_{28} \quad 39$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)]T_{29} \quad 40$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)]T_{30} \quad 41$$

$$+(a''_{28})^{(5)}(T_{29}, t) = \text{First augmentation factor} \quad 42$$

$$-(b''_{28})^{(5)}((G_{31}), t) = \text{First detritions factor} \quad 43$$

LHS OF THE YANG MILLS THEORY AND RHS OF THE YANG MILLS THEORY TAKEN TO THE OTHER SIDE THE LHS WOULD DISSIPATE THE RHS WITH OR WITHOUT TIME LAG: 44
45

MODULE NUMBERED SIX

:

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)} G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)] G_{32} \quad 46$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)} G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)] G_{33} \quad 47$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)} G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)] G_{34} \quad 48$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)} T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)] T_{32} \quad 49$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)} T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)] T_{33} \quad 50$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)} T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)] T_{34} \quad 51$$

$$+(a''_{32})^{(6)}(T_{33}, t) = \text{First augmentation factor} \quad 52$$

$$-(b''_{32})^{(6)}((G_{35}), t) = \text{First detritions factor} \quad 53$$

HOLISTIC CONCATENATE SYTEMAL EQUATIONS HENCEFORTH REFERRED TO AS "GLOBAL EQUATIONS" 54

We take in to consideration the following parameters, processes and concepts:

- (1) Acquisition of mass
- (2) Symmetry Breaking
- (3) Strong interaction
- (4) Unified Electroweak interaction
- (5) Continuous group of local transformations
- (6) Lagrangian Variance
- (7) Group generator in Gauge Theory
- (8) Vector field or Gauge field
- (9) Non commutative symmetry group in Gauge Theory
- (10) Yang Mills Theory (We repeat the same Bank's example. Individual debits and Credits are conservative so also the holistic one. Generalized theories are applied to various systems which are parameterized. And we live in 'measurement world'. Classification is done on the parameters of various systems to which the Theory is applied.)
- (11) First Term of the Lagrangian of the Yang Mills Theory(LHS)

$$\mathcal{L}_{gf} = -\frac{1}{4} \text{Tr}(F^2) = -\frac{1}{4} F^{\mu\nu\alpha} F_{\mu\nu}^{\alpha}$$

- (12) RHS of the Yang Mills Theory

$$\mathcal{L}_{gf} = -\frac{1}{4} \text{Tr}(F^2) = -\frac{1}{4} F^{\mu\nu\alpha} F_{\mu\nu}^{\alpha}$$

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)} G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)} & + (a''_{13})^{(1)}(T_{14}, t) & + (a''_{16})^{(2,2)}(T_{17}, t) & + (a''_{20})^{(3,3)}(T_{21}, t) \\ + (a''_{24})^{(4,4,4,4)}(T_{25}, t) & + (a''_{28})^{(5,5,5,5)}(T_{29}, t) & + (a''_{32})^{(6,6,6,6)}(T_{33}, t) & \end{array} \right] G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{|c|c|c|} \hline (a'_{14})^{(1)} & + (a''_{14})^{(1)}(T_{14}, t) & + (a''_{17})^{(2,2)}(T_{17}, t) & + (a''_{21})^{(3,3)}(T_{21}, t) \\ \hline \end{array} \right] G_{14} \tag{56}$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{|c|c|c|} \hline (a'_{15})^{(1)} & + (a''_{15})^{(1)}(T_{14}, t) & + (a''_{18})^{(2,2)}(T_{17}, t) & + (a''_{22})^{(3,3)}(T_{21}, t) \\ \hline \end{array} \right] G_{15} \tag{57}$$

Where $\boxed{(a'_{13})^{(1)}(T_{14}, t)}$, $\boxed{(a'_{14})^{(1)}(T_{14}, t)}$, $\boxed{(a'_{15})^{(1)}(T_{14}, t)}$ are first augmentation coefficients for category 1, 2 and 3 58

$\boxed{+(a''_{16})^{(2,2)}(T_{17}, t)}$, $\boxed{+(a''_{17})^{(2,2)}(T_{17}, t)}$, $\boxed{+(a''_{18})^{(2,2)}(T_{17}, t)}$ are second augmentation coefficient for category 1, 2 and 3 59

$\boxed{+(a''_{20})^{(3,3)}(T_{21}, t)}$, $\boxed{+(a''_{21})^{(3,3)}(T_{21}, t)}$, $\boxed{+(a''_{22})^{(3,3)}(T_{21}, t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{25})^{(4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a''_{26})^{(4,4,4,4)}(T_{25}, t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{29})^{(5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a''_{30})^{(5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{33})^{(6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a''_{34})^{(6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient for category 1, 2 and 3 60

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{array}{|c|c|c|} \hline (b'_{13})^{(1)} & - (b''_{13})^{(1)}(G, t) & - (b''_{16})^{(2,2)}(G_{19}, t) & - (b''_{20})^{(3,3)}(G_{23}, t) \\ \hline \end{array} \right] T_{13} \tag{61}$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{|c|c|c|} \hline (b'_{14})^{(1)} & - (b''_{14})^{(1)}(G, t) & - (b''_{17})^{(2,2)}(G_{19}, t) & - (b''_{21})^{(3,3)}(G_{23}, t) \\ \hline \end{array} \right] T_{14} \tag{62}$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{|c|c|c|} \hline (b'_{15})^{(1)} & - (b''_{15})^{(1)}(G, t) & - (b''_{18})^{(2,2)}(G_{19}, t) & - (b''_{22})^{(3,3)}(G_{23}, t) \\ \hline \end{array} \right] T_{15} \tag{63}$$

Where $\boxed{-(b'_{13})^{(1)}(G, t)}$, $\boxed{-(b'_{14})^{(1)}(G, t)}$, $\boxed{-(b'_{15})^{(1)}(G, t)}$ are first detrition coefficients for category 1, 2 and 3 64

$\boxed{-(b''_{16})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2,2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2,2)}(G_{19}, t)}$ are second detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3)}(G_{23}, t)}$ are third detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4)}(G_{27}, t)}$ are fourth detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5)}(G_{31}, t)}$ are fifth detritions coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6)}(G_{35}, t)}$ are sixth detritions coefficients for category 1, 2 and 3 65

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{|c|c|c|} \hline (a'_{16})^{(2)} & + (a''_{16})^{(2)}(T_{17}, t) & + (a''_{13})^{(1,1)}(T_{14}, t) & + (a''_{20})^{(3,3,3)}(T_{21}, t) \\ \hline \end{array} \right] G_{16} \tag{66}$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{|c|c|c|} \hline (a'_{17})^{(2)} & + (a''_{17})^{(2)}(T_{17}, t) & + (a''_{14})^{(1,1)}(T_{14}, t) & + (a''_{21})^{(3,3,3)}(T_{21}, t) \\ \hline \end{array} \right] G_{17} \tag{67}$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)} G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} \boxed{+(a''_{18})^{(2)}(T_{17}, t)} & \boxed{+(a'_{15})^{(1,1)}(T_{14}, t)} & \boxed{+(a'_{22})^{(3,3,3)}(T_{21}, t)} \\ \boxed{+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)} \end{array} \right] G_{18} \quad 68$$

Where $\boxed{+(a'_{16})^{(2)}(T_{17}, t)}$, $\boxed{+(a'_{17})^{(2)}(T_{17}, t)}$, $\boxed{+(a'_{18})^{(2)}(T_{17}, t)}$ are first augmentation coefficients for category 1, 2 and 3 69

$\boxed{+(a'_{13})^{(1,1)}(T_{14}, t)}$, $\boxed{+(a'_{14})^{(1,1)}(T_{14}, t)}$, $\boxed{+(a'_{15})^{(1,1)}(T_{14}, t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a'_{20})^{(3,3,3)}(T_{21}, t)}$, $\boxed{+(a'_{21})^{(3,3,3)}(T_{21}, t)}$, $\boxed{+(a'_{22})^{(3,3,3)}(T_{21}, t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a'_{24})^{(4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a'_{25})^{(4,4,4,4,4)}(T_{25}, t)}$, $\boxed{+(a'_{26})^{(4,4,4,4,4)}(T_{25}, t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a'_{28})^{(5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a'_{29})^{(5,5,5,5,5)}(T_{29}, t)}$, $\boxed{+(a'_{30})^{(5,5,5,5,5)}(T_{29}, t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a'_{32})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a'_{33})^{(6,6,6,6,6)}(T_{33}, t)}$, $\boxed{+(a'_{34})^{(6,6,6,6,6)}(T_{33}, t)}$ are sixth augmentation coefficient for category 1, 2 and 3 70

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} \boxed{-(b''_{16})^{(2)}(G_{19}, t)} & \boxed{-(b''_{13})^{(1,1)}(G, t)} & \boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)} \end{array} \right] T_{16} \quad 71$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)} T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} \boxed{-(b''_{17})^{(2)}(G_{19}, t)} & \boxed{-(b''_{14})^{(1,1)}(G, t)} & \boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)} \end{array} \right] T_{17} \quad 72$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)} T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} \boxed{-(b''_{18})^{(2)}(G_{19}, t)} & \boxed{-(b''_{15})^{(1,1)}(G, t)} & \boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)} \end{array} \right] T_{18} \quad 73$$

where $\boxed{-(b''_{16})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19}, t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19}, t)}$ are first detrition coefficients for category 1, 2 and 3 74

$\boxed{-(b''_{13})^{(1,1)}(G, t)}$, $\boxed{-(b''_{14})^{(1,1)}(G, t)}$, $\boxed{-(b''_{15})^{(1,1)}(G, t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{21})^{(3,3,3)}(G_{23}, t)}$, $\boxed{-(b''_{22})^{(3,3,3)}(G_{23}, t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27}, t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27}, t)}$ are fourth detritions coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31}, t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31}, t)}$ are fifth detritions coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35}, t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35}, t)}$ are sixth detritions coefficients for category 1,2 and 3 75

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)} G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} \boxed{+(a''_{20})^{(3)}(T_{21}, t)} & \boxed{+(a'_{16})^{(2,2,2)}(T_{17}, t)} & \boxed{+(a'_{13})^{(1,1,1)}(T_{14}, t)} \\ \boxed{+(a'_{24})^{(4,4,4,4,4,4)}(T_{25}, t)} & \boxed{+(a'_{28})^{(5,5,5,5,5,5)}(T_{29}, t)} & \boxed{+(a'_{32})^{(6,6,6,6,6,6)}(T_{33}, t)} \end{array} \right] G_{20} \quad 76$$

$$\frac{dG_{21}}{dt} = \quad 77$$

$$(a_{21})^{(3)}G_{20} - \left[\begin{array}{|c|} \hline (a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t) + (a'_{17})^{(2,2,2)}(T_{17}, t) + (a'_{14})^{(1,1,1)}(T_{14}, t) \\ \hline \end{array} \right] G_{21}$$

$$\frac{dG_{22}}{dt} = \left[\begin{array}{|c|} \hline (a'_{25})^{(4,4,4,4,4,4)}(T_{25}, t) + (a'_{29})^{(5,5,5,5,5,5)}(T_{29}, t) + (a'_{33})^{(6,6,6,6,6,6)}(T_{33}, t) \\ \hline \end{array} \right] G_{22}$$

$$(a_{22})^{(3)}G_{21} - \left[\begin{array}{|c|} \hline (a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t) + (a'_{18})^{(2,2,2)}(T_{17}, t) + (a'_{15})^{(1,1,1)}(T_{14}, t) \\ \hline \end{array} \right] G_{22}$$

$(a''_{20})^{(3)}(T_{21}, t), (a''_{21})^{(3)}(T_{21}, t), (a''_{22})^{(3)}(T_{21}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a''_{16})^{(2,2,2)}(T_{17}, t), (a''_{17})^{(2,2,2)}(T_{17}, t), (a''_{18})^{(2,2,2)}(T_{17}, t)$ are second augmentation coefficients for category 1, 2 and 3

$(a''_{13})^{(1,1,1)}(T_{14}, t), (a''_{14})^{(1,1,1)}(T_{14}, t), (a''_{15})^{(1,1,1)}(T_{14}, t)$ are third augmentation coefficients for category 1, 2 and 3

$(a''_{24})^{(4,4,4,4,4,4)}(T_{25}, t), (a''_{25})^{(4,4,4,4,4,4)}(T_{25}, t), (a''_{26})^{(4,4,4,4,4,4)}(T_{25}, t)$ are fourth augmentation coefficients for category 1, 2 and 3

$(a''_{28})^{(5,5,5,5,5,5)}(T_{29}, t), (a''_{29})^{(5,5,5,5,5,5)}(T_{29}, t), (a''_{30})^{(5,5,5,5,5,5)}(T_{29}, t)$ are fifth augmentation coefficients for category 1, 2 and 3

$(a''_{32})^{(6,6,6,6,6,6)}(T_{33}, t), (a''_{33})^{(6,6,6,6,6,6)}(T_{33}, t), (a''_{34})^{(6,6,6,6,6,6)}(T_{33}, t)$ are sixth augmentation coefficients for category 1, 2 and 3

$$(b_{20})^{(3)}T_{21} - \left[\begin{array}{|c|} \hline (b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b'_{16})^{(2,2,2)}(G_{19}, t) - (b'_{13})^{(1,1,1)}(G, t) \\ \hline \end{array} \right] T_{20}$$

$$(b_{21})^{(3)}T_{20} - \left[\begin{array}{|c|} \hline (b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t) - (b'_{17})^{(2,2,2)}(G_{19}, t) - (b'_{14})^{(1,1,1)}(G, t) \\ \hline \end{array} \right] T_{21}$$

$$(b_{22})^{(3)}T_{21} - \left[\begin{array}{|c|} \hline (b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t) - (b'_{18})^{(2,2,2)}(G_{19}, t) - (b'_{15})^{(1,1,1)}(G, t) \\ \hline \end{array} \right] T_{22}$$

$(b''_{20})^{(3)}(G_{23}, t), (b''_{21})^{(3)}(G_{23}, t), (b''_{22})^{(3)}(G_{23}, t)$ are first detritions coefficients for category 1, 2 and 3

$(b''_{16})^{(2,2,2)}(G_{19}, t), (b''_{17})^{(2,2,2)}(G_{19}, t), (b''_{18})^{(2,2,2)}(G_{19}, t)$ are second detritions coefficients for category 1, 2 and 3

$(b''_{13})^{(1,1,1)}(G, t), (b''_{14})^{(1,1,1)}(G, t), (b''_{15})^{(1,1,1)}(G, t)$ are third detrition coefficients for category 1, 2 and 3

$(b''_{24})^{(4,4,4,4,4,4)}(G_{27}, t), (b''_{25})^{(4,4,4,4,4,4)}(G_{27}, t), (b''_{26})^{(4,4,4,4,4,4)}(G_{27}, t)$ are fourth detritions coefficients for category 1, 2 and 3

$(b''_{28})^{(5,5,5,5,5,5)}(G_{31}, t), (b''_{29})^{(5,5,5,5,5,5)}(G_{31}, t), (b''_{30})^{(5,5,5,5,5,5)}(G_{31}, t)$ are fifth detritions coefficients for category 1, 2 and 3

$(b''_{32})^{(6,6,6,6,6,6)}(G_{35}, t), (b''_{33})^{(6,6,6,6,6,6)}(G_{35}, t), (b''_{34})^{(6,6,6,6,6,6)}(G_{35}, t)$ are sixth detritions coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{|c|c|c|c|} \hline (a'_{24})^{(4)} & + (a'_{24})^{(4)}(T_{25}, t) & + (a'_{28})^{(5,5)}(T_{29}, t) & + (a'_{32})^{(6,6)}(T_{33}, t) \\ \hline + (a'_{13})^{(1,1,1,1)}(T_{14}, t) & + (a'_{16})^{(2,2,2,2)}(T_{17}, t) & + (a'_{20})^{(3,3,3,3)}(T_{21}, t) & \\ \hline \end{array} \right] G_{24} \quad 87$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{|c|c|c|c|} \hline (a'_{25})^{(4)} & + (a'_{25})^{(4)}(T_{25}, t) & + (a'_{29})^{(5,5)}(T_{29}, t) & + (a'_{33})^{(6,6)}(T_{33}, t) \\ \hline + (a'_{14})^{(1,1,1,1)}(T_{14}, t) & + (a'_{17})^{(2,2,2,2)}(T_{17}, t) & + (a'_{21})^{(3,3,3,3)}(T_{21}, t) & \\ \hline \end{array} \right] G_{25} \quad 88$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{|c|c|c|c|} \hline (a'_{26})^{(4)} & + (a'_{26})^{(4)}(T_{25}, t) & + (a'_{30})^{(5,5)}(T_{29}, t) & + (a'_{34})^{(6,6)}(T_{33}, t) \\ \hline + (a'_{15})^{(1,1,1,1)}(T_{14}, t) & + (a'_{18})^{(2,2,2,2)}(T_{17}, t) & + (a'_{22})^{(3,3,3,3)}(T_{21}, t) & \\ \hline \end{array} \right] G_{26} \quad 89$$

Where $(a'_{24})^{(4)}(T_{25}, t)$, $(a'_{25})^{(4)}(T_{25}, t)$, $(a'_{26})^{(4)}(T_{25}, t)$ are first augmentation coefficients for category 1, 2 and 3

$(a'_{28})^{(5,5)}(T_{29}, t)$, $(a'_{29})^{(5,5)}(T_{29}, t)$, $(a'_{30})^{(5,5)}(T_{29}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a'_{32})^{(6,6)}(T_{33}, t)$, $(a'_{33})^{(6,6)}(T_{33}, t)$, $(a'_{34})^{(6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a'_{13})^{(1,1,1,1)}(T_{14}, t)$, $(a'_{14})^{(1,1,1,1)}(T_{14}, t)$, $(a'_{15})^{(1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$(a'_{16})^{(2,2,2,2)}(T_{17}, t)$, $(a'_{17})^{(2,2,2,2)}(T_{17}, t)$, $(a'_{18})^{(2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$(a'_{20})^{(3,3,3,3)}(T_{21}, t)$, $(a'_{21})^{(3,3,3,3)}(T_{21}, t)$, $(a'_{22})^{(3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, and 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{array}{|c|c|c|c|} \hline (b'_{24})^{(4)} & - (b'_{24})^{(4)}(G_{27}, t) & - (b'_{28})^{(5,5)}(G_{31}, t) & - (b'_{32})^{(6,6)}(G_{35}, t) \\ \hline - (b'_{13})^{(1,1,1,1)}(G, t) & - (b'_{16})^{(2,2,2,2)}(G_{19}, t) & - (b'_{20})^{(3,3,3,3)}(G_{23}, t) & \\ \hline \end{array} \right] T_{24} \quad 93$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{array}{|c|c|c|c|} \hline (b'_{25})^{(4)} & - (b'_{25})^{(4)}(G_{27}, t) & - (b'_{29})^{(5,5)}(G_{31}, t) & - (b'_{33})^{(6,6)}(G_{35}, t) \\ \hline - (b'_{14})^{(1,1,1,1)}(G, t) & - (b'_{17})^{(2,2,2,2)}(G_{19}, t) & - (b'_{21})^{(3,3,3,3)}(G_{23}, t) & \\ \hline \end{array} \right] T_{25} \quad 94$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{array}{|c|c|c|c|} \hline (b'_{26})^{(4)} & - (b'_{26})^{(4)}(G_{27}, t) & - (b'_{30})^{(5,5)}(G_{31}, t) & - (b'_{34})^{(6,6)}(G_{35}, t) \\ \hline - (b'_{15})^{(1,1,1,1)}(G, t) & - (b'_{18})^{(2,2,2,2)}(G_{19}, t) & - (b'_{22})^{(3,3,3,3)}(G_{23}, t) & \\ \hline \end{array} \right] T_{26} \quad 95$$

Where $-(b'_{24})^{(4)}(G_{27}, t)$, $-(b'_{25})^{(4)}(G_{27}, t)$, $-(b'_{26})^{(4)}(G_{27}, t)$ are first detrition coefficients for category 1, 2 and 3

$-(b'_{28})^{(5,5)}(G_{31}, t)$, $-(b'_{29})^{(5,5)}(G_{31}, t)$, $-(b'_{30})^{(5,5)}(G_{31}, t)$ are second detrition coefficients for category 1, 2 and 3

$-(b'_{32})^{(6,6)}(G_{35}, t)$, $-(b'_{33})^{(6,6)}(G_{35}, t)$, $-(b'_{34})^{(6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$-(b'_{13})^{(1,1,1,1)}(G, t)$, $-(b'_{14})^{(1,1,1,1)}(G, t)$, $-(b'_{15})^{(1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b'_{16})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{17})^{(2,2,2,2)}(G_{19}, t)$, $-(b'_{18})^{(2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b'_{20})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{21})^{(3,3,3,3)}(G_{23}, t)$, $-(b'_{22})^{(3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2 and 3

98

99

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\frac{(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)}{(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)} + \frac{(a''_{24})^{(4,4)}(T_{25}, t)}{(a'_{16})^{(2,2,2,2,2)}(T_{17}, t)} + \frac{(a''_{32})^{(6,6,6)}(T_{33}, t)}{(a'_{20})^{(3,3,3,3,3)}(T_{21}, t)} \right] G_{28}$$

100

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\frac{(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)}{(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)} + \frac{(a''_{25})^{(4,4)}(T_{25}, t)}{(a'_{17})^{(2,2,2,2,2)}(T_{17}, t)} + \frac{(a''_{33})^{(6,6,6)}(T_{33}, t)}{(a'_{21})^{(3,3,3,3,3)}(T_{21}, t)} \right] G_{29}$$

101

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\frac{(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)}{(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)} + \frac{(a''_{26})^{(4,4)}(T_{25}, t)}{(a'_{18})^{(2,2,2,2,2)}(T_{17}, t)} + \frac{(a''_{34})^{(6,6,6)}(T_{33}, t)}{(a'_{22})^{(3,3,3,3,3)}(T_{21}, t)} \right] G_{30}$$

Where $(a'_{28})^{(5)}(T_{29}, t)$, $(a'_{29})^{(5)}(T_{29}, t)$, $(a'_{30})^{(5)}(T_{29}, t)$ are first augmentation coefficients for category 1, 2 and 3

And $(a''_{24})^{(4,4)}(T_{25}, t)$, $(a''_{25})^{(4,4)}(T_{25}, t)$, $(a''_{26})^{(4,4)}(T_{25}, t)$ are second augmentation coefficient for category 1, 2 and 3

$(a''_{32})^{(6,6,6)}(T_{33}, t)$, $(a''_{33})^{(6,6,6)}(T_{33}, t)$, $(a''_{34})^{(6,6,6)}(T_{33}, t)$ are third augmentation coefficient for category 1, 2 and 3

$(a'_{13})^{(1,1,1,1,1)}(T_{14}, t)$, $(a'_{14})^{(1,1,1,1,1)}(T_{14}, t)$, $(a'_{15})^{(1,1,1,1,1)}(T_{14}, t)$ are fourth augmentation coefficients for category 1, 2, and 3

$(a'_{16})^{(2,2,2,2,2)}(T_{17}, t)$, $(a'_{17})^{(2,2,2,2,2)}(T_{17}, t)$, $(a'_{18})^{(2,2,2,2,2)}(T_{17}, t)$ are fifth augmentation coefficients for category 1, 2, and 3

$(a'_{20})^{(3,3,3,3,3)}(T_{21}, t)$, $(a'_{21})^{(3,3,3,3,3)}(T_{21}, t)$, $(a'_{22})^{(3,3,3,3,3)}(T_{21}, t)$ are sixth augmentation coefficients for category 1, 2, 3

103

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\frac{(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}, t)}{(b'_{13})^{(1,1,1,1,1)}(G, t)} + \frac{(b''_{24})^{(4,4)}(G_{27}, t)}{(b'_{16})^{(2,2,2,2,2)}(G_{19}, t)} + \frac{(b''_{32})^{(6,6,6)}(G_{35}, t)}{(b'_{20})^{(3,3,3,3,3)}(G_{23}, t)} \right] T_{28}$$

104

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\frac{(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}, t)}{(b'_{14})^{(1,1,1,1,1)}(G, t)} + \frac{(b''_{25})^{(4,4)}(G_{27}, t)}{(b'_{17})^{(2,2,2,2,2)}(G_{19}, t)} + \frac{(b''_{33})^{(6,6,6)}(G_{35}, t)}{(b'_{21})^{(3,3,3,3,3)}(G_{23}, t)} \right] T_{29}$$

105

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\frac{(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}, t)}{(b'_{15})^{(1,1,1,1,1)}(G, t)} + \frac{(b''_{26})^{(4,4)}(G_{27}, t)}{(b'_{18})^{(2,2,2,2,2)}(G_{19}, t)} + \frac{(b''_{34})^{(6,6,6)}(G_{35}, t)}{(b'_{22})^{(3,3,3,3,3)}(G_{23}, t)} \right] T_{30}$$

106

where $(b'_{28})^{(5)}(G_{31}, t)$, $(b'_{29})^{(5)}(G_{31}, t)$, $(b'_{30})^{(5)}(G_{31}, t)$ are first detrition coefficients for category 1, 2 and 3

$(b''_{24})^{(4,4)}(G_{27}, t)$, $(b''_{25})^{(4,4)}(G_{27}, t)$, $(b''_{26})^{(4,4)}(G_{27}, t)$ are second detrition coefficients for category 1, 2 and 3

$(b''_{32})^{(6,6,6)}(G_{35}, t)$, $(b''_{33})^{(6,6,6)}(G_{35}, t)$, $(b''_{34})^{(6,6,6)}(G_{35}, t)$ are third detrition coefficients for category 1, 2 and 3

$(b'_{13})^{(1,1,1,1,1)}(G, t)$, $(b'_{14})^{(1,1,1,1,1)}(G, t)$, $(b'_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2, and 3

$(b'_{16})^{(2,2,2,2,2)}(G_{19}, t)$, $(b'_{17})^{(2,2,2,2,2)}(G_{19}, t)$, $(b'_{18})^{(2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2, and 3

$(b'_{20})^{(3,3,3,3,3)}(G_{23}, t)$, $(b'_{21})^{(3,3,3,3,3)}(G_{23}, t)$, $(b'_{22})^{(3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2, and 3

108

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) & + (a''_{28})^{(5,5,5)}(T_{29}, t) & + (a''_{24})^{(4,4,4)}(T_{25}, t) \\ + (a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t) \end{array} \right] G_{32} \quad 109$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t) & + (a''_{29})^{(5,5,5)}(T_{29}, t) & + (a''_{25})^{(4,4,4)}(T_{25}, t) \\ + (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t) \end{array} \right] G_{33} \quad 110$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t) & + (a''_{30})^{(5,5,5)}(T_{29}, t) & + (a''_{26})^{(4,4,4)}(T_{25}, t) \\ + (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t) & + (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t) & + (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t) \end{array} \right] G_{34} \quad 111$$

$(a''_{32})^{(6)}(T_{33}, t), (a''_{33})^{(6)}(T_{33}, t), (a''_{34})^{(6)}(T_{33}, t)$ are first augmentation coefficients for category 1, 2 and 3 112

$(a''_{28})^{(5,5,5)}(T_{29}, t), (a''_{29})^{(5,5,5)}(T_{29}, t), (a''_{30})^{(5,5,5)}(T_{29}, t)$ are second augmentation coefficients for category 1, 2 and 3

$(a''_{24})^{(4,4,4)}(T_{25}, t), (a''_{25})^{(4,4,4)}(T_{25}, t), (a''_{26})^{(4,4,4)}(T_{25}, t)$ are third augmentation coefficients for category 1, 2 and 3

$(a''_{13})^{(1,1,1,1,1,1)}(T_{14}, t), (a''_{14})^{(1,1,1,1,1,1)}(T_{14}, t), (a''_{15})^{(1,1,1,1,1,1)}(T_{14}, t)$ - are fourth augmentation coefficients

$(a''_{16})^{(2,2,2,2,2,2)}(T_{17}, t), (a''_{17})^{(2,2,2,2,2,2)}(T_{17}, t), (a''_{18})^{(2,2,2,2,2,2)}(T_{17}, t)$ - fifth augmentation coefficients

$(a''_{20})^{(3,3,3,3,3,3)}(T_{21}, t), (a''_{21})^{(3,3,3,3,3,3)}(T_{21}, t), (a''_{22})^{(3,3,3,3,3,3)}(T_{21}, t)$ sixth augmentation coefficients

113

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}, t) & - (b''_{28})^{(5,5,5)}(G_{31}, t) & - (b''_{24})^{(4,4,4)}(G_{27}, t) \\ - (b''_{13})^{(1,1,1,1,1,1)}(G, t) & - (b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t) \end{array} \right] T_{32} \quad 114$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[\begin{array}{ccc} (b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}, t) & - (b''_{29})^{(5,5,5)}(G_{31}, t) & - (b''_{25})^{(4,4,4)}(G_{27}, t) \\ - (b''_{14})^{(1,1,1,1,1,1)}(G, t) & - (b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t) \end{array} \right] T_{33} \quad 115$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[\begin{array}{ccc} (b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}, t) & - (b''_{30})^{(5,5,5)}(G_{31}, t) & - (b''_{26})^{(4,4,4)}(G_{27}, t) \\ - (b''_{15})^{(1,1,1,1,1,1)}(G, t) & - (b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t) & - (b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t) \end{array} \right] T_{34} \quad 116$$

$-(b''_{32})^{(6)}(G_{35}, t), -(b''_{33})^{(6)}(G_{35}, t), -(b''_{34})^{(6)}(G_{35}, t)$ are first detrition coefficients for category 1,2 and 3 117

$-(b''_{28})^{(5,5,5)}(G_{31}, t), -(b''_{29})^{(5,5,5)}(G_{31}, t), -(b''_{30})^{(5,5,5)}(G_{31}, t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4)}(G_{27}, t), -(b''_{25})^{(4,4,4)}(G_{27}, t), -(b''_{26})^{(4,4,4)}(G_{27}, t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{13})^{(1,1,1,1,1)}(G, t), -(b''_{14})^{(1,1,1,1,1)}(G, t), -(b''_{15})^{(1,1,1,1,1)}(G, t)$ are fourth detrition coefficients for category 1, 2, and 3

$-(b''_{16})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{17})^{(2,2,2,2,2,2)}(G_{19}, t), -(b''_{18})^{(2,2,2,2,2,2)}(G_{19}, t)$ are fifth detrition coefficients for category 1, 2, and 3

$-(b''_{20})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{21})^{(3,3,3,3,3,3)}(G_{23}, t), -(b''_{22})^{(3,3,3,3,3,3)}(G_{23}, t)$ are sixth detrition coefficients for category 1, 2, and 3

118

Where we suppose

119

(A) $(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0,$ 120

$i, j = 13,14,15$

(B) The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}, (r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14}, t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

121

$$(b''_i)^{(1)}(G, t) \leq (r_i)^{(1)} \leq (b'_i)^{(1)} \leq (\hat{B}_{13})^{(1)}$$

(C) $\lim_{T_2 \rightarrow \infty} (a''_i)^{(1)}(T_{14}, t) = (p_i)^{(1)}$ 122

$$\lim_{G \rightarrow \infty} (b''_i)^{(1)}(G, t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}$ are positive constants and $i = 13,14,15$

They satisfy Lipschitz condition: 123

$$|(a''_i)^{(1)}(T'_{14}, t) - (a''_i)^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)} |T'_{14} - T_{14}| e^{-(\hat{M}_{13})^{(1)}t}$$

124

$$|(b''_i)^{(1)}(G', t) - (b''_i)^{(1)}(G, t)| < (\hat{k}_{13})^{(1)} ||G - G'| e^{-(\hat{M}_{13})^{(1)}t}$$

125

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(1)}(T'_{14}, t)$ and $(a''_i)^{(1)}(T_{14}, t)$. (T'_{14}, t) and (T_{14}, t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a''_i)^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a''_i)^{(1)}(T_{14}, t)$, the first augmentation coefficient WOULD be absolutely continuous. 126

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 127

(D) $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 128

(E) There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ which together
 with $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants
 $(a_i)^{(1)}, (a'_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13,14,15,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(a_i)^{(1)} + (a'_i)^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}} [(b_i)^{(1)} + (b'_i)^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

(F) $(a_i)^{(2)}, (a'_i)^{(2)}, (a''_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (b''_i)^{(2)} > 0, \quad i, j = 16,17,18$

(G) The functions $(a''_i)^{(2)}, (b''_i)^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}, (r_i)^{(2)}$:

$$(a''_i)^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)}$$

$$(b''_i)^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b'_i)^{(2)} \leq (\hat{B}_{16})^{(2)}$$

(H) $\lim_{T_2 \rightarrow \infty} (a''_i)^{(2)}(T_{17}, t) = (p_i)^{(2)}$

$$\lim_{G \rightarrow \infty} (b''_i)^{(2)}(G_{19}, t) = (r_i)^{(2)}$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$:

Where $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}$ are positive constants and $i = 16,17,18$

They satisfy Lipschitz condition:

$$|(a''_i)^{(2)}(T'_{17}, t) - (a''_i)^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T'_{17} - T_{17}| e^{-(\hat{M}_{16})^{(2)}t}$$

$$|(b''_i)^{(2)}((G'_{19}), t) - (b''_i)^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} |(G'_{19}) - (G_{19})'| e^{-(\hat{M}_{16})^{(2)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(2)}(T'_{17}, t)$ and $(a''_i)^{(2)}(T_{17}, t)$. (T'_{17}, t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a''_i)^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a''_i)^{(2)}(T_{17}, t)$, the SECOND augmentation coefficient would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

(I) $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together
 with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)}$ and $(\hat{B}_{16})^{(2)}$ and the constants
 $(a_i)^{(2)}, (a'_i)^{(2)}, (b_i)^{(2)}, (b'_i)^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(a_i)^{(2)} + (a'_i)^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 150$$

$$\frac{1}{(\hat{M}_{16})^{(2)}} [(b_i)^{(2)} + (b'_i)^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 151$$

Where we suppose 152

$$(J) \quad (a_i)^{(3)}, (a'_i)^{(3)}, (a''_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (b''_i)^{(3)} > 0, \quad i, j = 20, 21, 22 \quad 153$$

The functions $(a''_i)^{(3)}, (b''_i)^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}, (r_i)^{(3)}$:

$$(a''_i)^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b''_i)^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b'_i)^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$\lim_{T_2 \rightarrow \infty} (a''_i)^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 154$$

$$\lim_{G \rightarrow \infty} (b''_i)^{(3)}(G_{23}, t) = (r_i)^{(3)} \quad 155$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$: 156

Where $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}$ are positive constants and $i = 20, 21, 22$

They satisfy Lipschitz condition: 157

$$|(a''_i)^{(3)}(T'_{21}, t) - (a''_i)^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)} |T_{21} - T'_{21}| e^{-(\hat{M}_{20})^{(3)}t} \quad 158$$

$$|(b''_i)^{(3)}(G_{23}', t) - (b''_i)^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)} ||G_{23} - G_{23}'|| e^{-(\hat{M}_{20})^{(3)}t} \quad 159$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a''_i)^{(3)}(T'_{21}, t)$ and $(a''_i)^{(3)}(T_{21}, t)$. (T'_{21}, t) and (T_{21}, t) are points belonging to the interval $[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}]$. It is to be noted that $(a''_i)^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a''_i)^{(3)}(T_{21}, t)$, the THIRD augmentation coefficient, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 161

(K) $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)}$ and $(\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a'_i)^{(3)}, (b_i)^{(3)}, (b'_i)^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20, 21, 22$, satisfy the inequalities 162
163
164

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(a_i)^{(3)} + (a'_i)^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1 \quad 165$$

$$\frac{1}{(\hat{M}_{20})^{(3)}} [(b_i)^{(3)} + (b'_i)^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1 \quad 166$$

Where we suppose 168

$$(a_i)^{(4)}, (a'_i)^{(4)}, (a''_i)^{(4)}, (b_i)^{(4)}, (b'_i)^{(4)}, (b''_i)^{(4)} > 0, \quad i, j = 24, 25, 26 \quad 169$$

(M) The functions $(a''_i)^{(4)}, (b''_i)^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}, (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

170

(N) $\lim_{T_2 \rightarrow \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)}$
 $\lim_{G \rightarrow \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}$ are positive constants and $i = 24, 25, 26$

They satisfy Lipschitz condition:

171

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25}' - T_{25}| e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} |(G_{27})' - (G_{27})| e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 4$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **FOURTH augmentation coefficient WOULD** be absolutely continuous.

172

173

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$:

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}} + \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$:

175

(Q) There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)}$ and $(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24, 25, 26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}} [(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)} (\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

176

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28, 29, 30$$

177

(S) The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}, (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}), t) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

178

$$(T) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}$ are positive constants and $i = 28, 29, 30$

They satisfy Lipschitz condition:

179

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)} |T_{29}' - T_{29}| e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}((G_{31}), t)| < (\hat{k}_{28})^{(5)} |(G_{31})' - (G_{31})| e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 5$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the FIFTH **augmentation coefficient** attributable would be absolutely continuous.

180

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$:

181

$$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, \text{ are positive constants}$$

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$:

182

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)}$ and $(\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28, 29, 30$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}} [(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

183

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32, 33, 34$$

184

(W) The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

185

$$(X) \quad \lim_{T_2 \rightarrow \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$

$$\lim_{G \rightarrow \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}$ are positive constants and $i = 32, 33, 34$

They satisfy Lipschitz condition:

186

$$|(a_i'')^{(6)}(T_{33}, t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T_{33}'| e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} \|(G_{35}) - (G_{35})'\| e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}, t)$ and $(a_i'')^{(6)}(T_{33}, t) \cdot (T_{33}', t)$ and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 6$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the SIXTH **augmentation coefficient** would be absolutely continuous.

187

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$:

188

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$ are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$:

189

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)}$ and $(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32, 33, 34$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}} [(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

190

Theorem 1: if the conditions IN THE FOREGOING above are fulfilled, there exists a solution satisfying the conditions

191

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}, \quad T_i(0) = T_i^0 > 0$$

192

193

Definition of $G_i(0), T_i(0)$

$$G_i(t) \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t}, \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t} , \quad T_i(0) = T_i^0 > 0$$

194

195

$$G_i(t) \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t} , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t} , \quad T_i(0) = T_i^0 > 0$$

Definition of $G_i(0), T_i(0)$:

196

$$G_i(t) \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

197

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

198

Definition of $G_i(0), T_i(0)$:

199

$$G_i(t) \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t} , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t} , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

200

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{13})^{(1)}, T_i^0 \leq (\hat{Q}_{13})^{(1)},$$

201

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$$

202

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$$

203

By

204

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + (a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

205

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

206

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

207

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)} \quad 208$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)} \quad 209$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

210

Proof:

211

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{16})^{(2)}, T_i^0 \leq (\hat{Q}_{16})^{(2)}, \quad 212$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t} \quad 213$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)}t} \quad 214$$

By

215

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)} \quad 216$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)} \quad 217$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)} \quad 218$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)} \quad 219$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)} \quad 220$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

221

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{20})^{(3)}, T_i^0 \leq (\hat{Q}_{20})^{(3)}, \quad 222$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t} \quad 223$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t} \quad 224$$

By

225

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)} \quad 226$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)} \quad 227$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)} \quad 228$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)} \quad 229$$

$$\bar{T}_{22}(t) = T_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)} \quad 230$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy 231

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{24})^{(4)}, T_i^0 \leq (\hat{Q}_{24})^{(4)}, \quad 232$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad 233$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad 234$$

By 235

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)} \quad 236$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)} \quad 237$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)} \quad 238$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)} \quad 239$$

$$\bar{T}_{26}(t) = T_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)} \quad 240$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy 241

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{28})^{(5)}, T_i^0 \leq (\hat{Q}_{28})^{(5)}, \quad 243$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad 244$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad 245$$

By 246

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)} \quad 247$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)} \quad 248$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)} \quad 249$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)} \quad 250$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)} \quad 251$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

252

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions $G_i, T_i: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0, T_i(0) = T_i^0, G_i^0 \leq (\hat{P}_{32})^{(6)}, T_i^0 \leq (\hat{Q}_{32})^{(6)}, \quad 253$$

$$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t} \quad 254$$

$$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t} \quad 255$$

By 256

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32}{}^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)} \quad 257$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)} \quad 258$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)} \quad 259$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)} \quad 260$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)} (G(s_{(32)}), s_{(32)}) \right) T_{34}(s_{(32)}) \right] ds_{(32)} \quad 261$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

262

(a) The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying GLOBAL EQUATIONS into itself 263
 .Indeed it is obvious that

$$G_{13}(t) \leq G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}s_{(13)}} \right) \right] ds_{(13)} = \\
 \left(1 + (a_{13})^{(1)} t \right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)}t} - 1 \right)$$

From which it follows that 264

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)}t} \leq \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0 \right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0} \right)} + (\hat{P}_{13})^{(1)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$ 265

(b) The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying GLOBAL EQUATIONS into itself 266
 .Indeed it is obvious that

$$G_{16}(t) \leq G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}} \right) \right] ds_{(16)} = (1 + (a_{16})^{(2)} t) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1 \right) \quad 267$$

From which it follows that 268

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \leq \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0 \right) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0} \right)} + (\hat{P}_{16})^{(2)} \right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$ 269

(a) The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying GLOBAL EQUATIONS into itself 270
 .Indeed it is obvious that

$$G_{20}(t) \leq G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}} \right) \right] ds_{(20)} = (1 + (a_{20})^{(3)} t) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1 \right)$$

From which it follows that 271

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \leq \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^0 \right) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0} \right)} + (\hat{P}_{20})^{(3)} \right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$ 272

(b) The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying GLOBAL EQUATIONS into itself 273
 .Indeed it is obvious that

$$G_{24}(t) \leq G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}} \right) \right] ds_{(24)} = (1 + (a_{24})^{(4)} t) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1 \right)$$

From which it follows that 274

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \leq \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left[\left((\hat{P}_{24})^{(4)} + G_{25}^0 \right) e^{-\left(\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0} \right)} + (\hat{P}_{24})^{(4)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

(c) The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying GLOBAL EQUATIONS into itself 275
 .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} \left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}} \right) \right] ds_{(28)} = (1 + (a_{28})^{(5)} t) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}} \left(e^{(\hat{M}_{28})^{(5)} t} - 1 \right)$$

From which it follows that 276

$$(G_{28}(t) - G_{28}^0)e^{-(\bar{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\bar{M}_{28})^{(5)}} \left[((\hat{P}_{28})^{(5)} + G_{29}^0)e^{-\left(\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)} \right]$$

(G_i^0) is as defined in the statement of theorem 1

(d) The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying GLOBAL EQUATIONS into itself 277
 .Indeed it is obvious that

$$G_{32}(t) \leq G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}s_{(32)}} \right) \right] ds_{(32)} =$$

$$\left(1 + (a_{32})^{(6)}t \right) G_{33}^0 + \frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\bar{M}_{32})^{(6)}} \left(e^{(\bar{M}_{32})^{(6)}t} - 1 \right)$$

From which it follows that

278

$$(G_{32}(t) - G_{32}^0)e^{-(\bar{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\bar{M}_{32})^{(6)}} \left[((\hat{P}_{32})^{(6)} + G_{33}^0)e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0}\right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

279

280

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\bar{M}_{13})^{(1)}}, \frac{(b_j)^{(1)}}{(\bar{M}_{13})^{(1)}} < 1$ and to choose

281

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

282

$$\frac{(a_i)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0 \right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{13})^{(1)}$$

283

$$\frac{(b_j)^{(1)}}{(\bar{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0 \right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)} \right] \leq (\hat{Q}_{13})^{(1)}$$

284

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying GLOBAL EQUATIONS into itself 285

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric

286

$$d \left((G^{(1)}, T^{(1)}), (G^{(2)}, T^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{13})^{(1)}t} \}$$

Indeed if we denote

287

Definition of \tilde{G}, \tilde{T} :

$$(\tilde{G}, \tilde{T}) = \mathcal{A}^{(1)}(G, T)$$

It results

$$\begin{aligned}
 |\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{13})^{(1)} |G_{14}^{(1)} - G_{14}^{(2)}| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} ds_{(13)} + \\
 &\int_0^t \{(a'_{13})^{(1)} |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} + \\
 &(a''_{13})^{(1)}(T_{14}^{(1)}, s_{(13)}) |G_{13}^{(1)} - G_{13}^{(2)}| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}} + \\
 &G_{13}^{(2)} |(a'_{13})^{(1)}(T_{14}^{(1)}, s_{(13)}) - (a'_{13})^{(1)}(T_{14}^{(2)}, s_{(13)})| e^{-(\widehat{M}_{13})^{(1)}s_{(13)}} e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\} ds_{(13)}
 \end{aligned}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned}
 |G^{(1)} - G^{(2)}| e^{-(\widehat{M}_{13})^{(1)}t} &\leq \\
 \frac{1}{(\widehat{M}_{13})^{(1)}} &\left((a_{13})^{(1)} + (a'_{13})^{(1)} + (\widehat{A}_{13})^{(1)} + (\widehat{P}_{13})^{(1)} (\widehat{k}_{13})^{(1)} \right) d\left((G^{(1)}, T^{(1)}; G^{(2)}, T^{(2)}) \right)
 \end{aligned} \tag{288}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a'_{13})^{(1)}$ and $(b'_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ and $(\widehat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t}$ respectively of \mathbb{R}_+ . 289

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(1)}$ and $(b'_i)^{(1)}$, $i = 13, 14, 15$ depend only on T_{14} and respectively on G (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 290

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(1)} - (a''_i)^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}} \geq 0 \tag{291}$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(1)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{13})^{(1)})_1$, and $((\widehat{M}_{13})^{(1)})_3$: 292

Remark 3: if G_{13} is bounded, the same property have also G_{14} and G_{15} . indeed if

$$G_{13} < (\widehat{M}_{13})^{(1)} \text{ it follows } \frac{dG_{14}}{dt} \leq ((\widehat{M}_{13})^{(1)})_1 - (a'_{14})^{(1)} G_{14} \text{ and by integrating}$$

$$G_{14} \leq ((\widehat{M}_{13})^{(1)})_2 = G_{14}^0 + 2(a_{14})^{(1)} ((\widehat{M}_{13})^{(1)})_1 / (a'_{14})^{(1)}$$

In the same way, one can obtain

$$G_{15} \leq ((\widehat{M}_{13})^{(1)})_3 = G_{15}^0 + 2(a_{15})^{(1)} ((\widehat{M}_{13})^{(1)})_2 / (a'_{15})^{(1)}$$

If G_{14} or G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} is bounded, from below, the same property holds for G_{14} and G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 293

Remark 5: If T_{13} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(1)}(G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \rightarrow \infty$. 294

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to 295

$$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1} \right) (1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$$

If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2} \right)$, $t = \log \frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t \rightarrow \infty} (b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

296

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} < 1$ and to choose

297

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[(\hat{P}_{16})^{(2)} + ((\hat{P}_{16})^{(2)} + G_j^0) e^{-\left(\frac{(\hat{P}_{16})^{(2)} + G_j^0}{G_j^0} \right)} \right] \leq (\hat{P}_{16})^{(2)}$$

298

$$\frac{(b_i)^{(2)}}{(\bar{M}_{16})^{(2)}} \left[((\hat{Q}_{16})^{(2)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{16})^{(2)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{16})^{(2)} \right] \leq (\hat{Q}_{16})^{(2)}$$

299

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying

300

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric

301

$$d \left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{16})^{(2)}t} \}$$

Indeed if we denote

302

Definition of $\tilde{G}_{19}, \tilde{T}_{19} : (\tilde{G}_{19}, \tilde{T}_{19}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results

303

$$\begin{aligned} |\tilde{G}_{16}^{(1)} - \tilde{G}_{16}^{(2)}| &\leq \int_0^t (a_{16})^{(2)} |G_{17}^{(1)} - G_{17}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} ds_{(16)} + \\ &\int_0^t \{ (a'_{16})^{(2)} |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{-(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &(a''_{16})^{(2)}(T_{17}^{(1)}, s_{(16)}) |G_{16}^{(1)} - G_{16}^{(2)}| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} + \\ &G_{16}^{(2)} |(a'_{16})^{(2)}(T_{17}^{(1)}, s_{(16)}) - (a'_{16})^{(2)}(T_{17}^{(2)}, s_{(16)})| e^{-(\bar{M}_{16})^{(2)}s_{(16)}} e^{(\bar{M}_{16})^{(2)}s_{(16)}} \} ds_{(16)} \end{aligned}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$

304

From the hypotheses it follows

$$\begin{aligned} & |(G_{19})^{(1)} - (G_{19})^{(2)}| e^{-(\widehat{M}_{16})^{(2)}t} \leq \quad 305 \\ & \frac{1}{(\widehat{M}_{16})^{(2)}} \left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + \right. \\ & \left. (\widehat{P}_{16})^{(2)} (\widehat{k}_{16})^{(2)} \right) d \left(((G_{19})^{(1)}, (T_{19})^{(1)}); ((G_{19})^{(2)}, (T_{19})^{(2)}) \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows 306

Remark 1: The fact that we supposed $(a'_{16})^{(2)}$ and $(b'_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$ respectively of \mathbb{R}_+ . 307

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(2)}$ and $(b'_i)^{(2)}$, $i = 16, 17, 18$ depend only on T_{17} and respectively on (G_{19}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 308

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a'_i)^{(2)} - (a''_i)^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b'_i)^{(2)}t} > 0 \quad \text{for } t > 0$$

Definition of $((\widehat{M}_{16})^{(2)})_1$, $((\widehat{M}_{16})^{(2)})_2$ and $((\widehat{M}_{16})^{(2)})_3$: 309

Remark 3: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$$G_{16} < (\widehat{M}_{16})^{(2)} \text{ it follows } \frac{dG_{17}}{dt} \leq ((\widehat{M}_{16})^{(2)})_1 - (a'_{17})^{(2)} G_{17} \text{ and by integrating}$$

$$G_{17} \leq ((\widehat{M}_{16})^{(2)})_2 = G_{17}^0 + 2(a_{17})^{(2)} ((\widehat{M}_{16})^{(2)})_1 / (a'_{17})^{(2)}$$

In the same way, one can obtain

$$G_{18} \leq ((\widehat{M}_{16})^{(2)})_3 = G_{18}^0 + 2(a_{18})^{(2)} ((\widehat{M}_{16})^{(2)})_2 / (a'_{18})^{(2)} \quad 310$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 4: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 311

Remark 5: If T_{16} is bounded from below and $\lim_{t \rightarrow \infty} ((b'_i)^{(2)}((G_{19})(t), t)) = (b'_{17})^{(2)}$ then $T_{17} \rightarrow \infty$. 312

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b'_i)^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 313

$$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2} \right) (1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_2 t} = \frac{1}{2} \text{ it results}$$

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2} \right)$, $t = \log \frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t \rightarrow \infty} (b'_{18})^{(2)}((G_{19})(t), t) = (b'_{18})^{(2)}$ 314

We now state a more precise theorem about the behaviors at infinity of the solutions

315

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} < 1$ and to choose

316

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[(\hat{P}_{20})^{(3)} + ((\hat{P}_{20})^{(3)} + G_j^0) e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{20})^{(3)}$$

317

$$\frac{(b_i)^{(3)}}{(\bar{M}_{20})^{(3)}} \left[((\hat{Q}_{20})^{(3)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)} \right] \leq (\hat{Q}_{20})^{(3)}$$

318

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i into itself

319

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric

320

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{20})^{(3)}t} \}$$

Indeed if we denote

321

$$\text{Definition of } \widetilde{G}_{23}, \widetilde{T}_{23} : (\widetilde{G}_{23}, \widetilde{T}_{23}) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$$

It results

322

$$\begin{aligned} |\widetilde{G}_{20}^{(1)} - \widetilde{G}_{20}^{(2)}| &\leq \int_0^t (a_{20})^{(3)} |G_{21}^{(1)} - G_{21}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} ds_{(20)} + \\ &\int_0^t \{ (a'_{20})^{(3)} |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{-(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) |G_{20}^{(1)} - G_{20}^{(2)}| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} + \\ &G_{20}^{(2)} |(a''_{20})^{(3)} (T_{21}^{(1)}, s_{(20)}) - (a''_{20})^{(3)} (T_{21}^{(2)}, s_{(20)})| e^{-(\bar{M}_{20})^{(3)}s_{(20)}} e^{(\bar{M}_{20})^{(3)}s_{(20)}} \} ds_{(20)} \end{aligned}$$

323

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |G^{(1)} - G^{(2)}| e^{-(\bar{M}_{20})^{(3)}t} &\leq \\ &\frac{1}{(\bar{M}_{20})^{(3)}} \left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\hat{A}_{20})^{(3)} + \right. \\ &\left. (\hat{P}_{20})^{(3)} (\hat{K}_{20})^{(3)} \right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)}) \right) \end{aligned}$$

324

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ and $(\hat{Q}_{20})^{(3)} e^{(\bar{M}_{20})^{(3)}t}$ respectively of \mathbb{R}_+ .

325

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}$, $i = 20, 21, 22$ depend only on T_{21} and respectively on

(G_{23}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 326

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s(20)), s(20))\} ds(20)} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(3)}t} > 0 \quad \text{for } t > 0$$

Definition of $((\widehat{M}_{20})^{(3)})_1, ((\widehat{M}_{20})^{(3)})_2$ and $((\widehat{M}_{20})^{(3)})_3$: 327

Remark 3: if G_{20} is bounded, the same property holds also for G_{21} and G_{22} . indeed if

$$G_{20} < ((\widehat{M}_{20})^{(3)}) \text{ it follows } \frac{dG_{21}}{dt} \leq ((\widehat{M}_{20})^{(3)})_1 - (a_{21}')^{(3)}G_{21} \text{ and by integrating}$$

$$G_{21} \leq ((\widehat{M}_{20})^{(3)})_2 = G_{21}^0 + 2(a_{21}')^{(3)}((\widehat{M}_{20})^{(3)})_1 / (a_{21}')^{(3)}$$

In the same way, one can obtain

$$G_{22} \leq ((\widehat{M}_{20})^{(3)})_3 = G_{22}^0 + 2(a_{22}')^{(3)}((\widehat{M}_{20})^{(3)})_2 / (a_{22}')^{(3)}$$

If G_{21} or G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 4: If G_{20} is bounded, from below, the same property holds for G_{21} and G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 328

Remark 5: If T_{20} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(3)}((G_{23})(t), t)) = (b_{21}')^{(3)}$ then $T_{21} \rightarrow \infty$. 329

Definition of $(m)^{(3)}$ and ε_3 : 330

Indeed let t_3 be so that for $t > t_3$

$$(b_{21}')^{(3)} - (b_i'')^{(3)}((G_{23})(t), t) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21}')^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 331

$$T_{21} \geq \left(\frac{(a_{21}')^{(3)}(m)^{(3)}}{\varepsilon_3} \right) (1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_3 t} = \frac{1}{2} \text{ it results}$$

$T_{21} \geq \left(\frac{(a_{21}')^{(3)}(m)^{(3)}}{2} \right)$, $t = \log \frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t \rightarrow \infty} (b_{22}'')^{(3)}((G_{23})(t), t) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

332

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose

333

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} \left[(\widehat{P}_{24})^{(4)} + ((\widehat{P}_{24})^{(4)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{24})^{(4)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{24})^{(4)} \quad 334$$

$$\frac{(b_i)^{(4)}}{(\bar{M}_{24})^{(4)}} \left[\left((\hat{Q}_{24})^{(4)} + T_j^0 \right) e^{-\left(\frac{(\hat{Q}_{24})^{(4)} + T_j^0}{T_j^0} \right)} + (\hat{Q}_{24})^{(4)} \right] \leq (\hat{Q}_{24})^{(4)} \quad 335$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying IN to itself 336

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 337

$$d \left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) = \sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{24})^{(4)}t} \}$$

Indeed if we denote

Definition of $(\widetilde{G}_{27}), (\widetilde{T}_{27}) : (\widetilde{G}_{27}), (\widetilde{T}_{27}) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\begin{aligned} |\widetilde{G}_{24}^{(1)} - \widetilde{G}_i^{(2)}| &\leq \int_0^t (a_{24})^{(4)} |G_{25}^{(1)} - G_{25}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} ds_{(24)} + \\ &\int_0^t \{ (a'_{24})^{(4)} |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{-(\bar{M}_{24})^{(4)}s_{(24)}} + \\ &(a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) |G_{24}^{(1)} - G_{24}^{(2)}| e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} + \\ &G_{24}^{(2)} | (a''_{24})^{(4)} (T_{25}^{(1)}, s_{(24)}) - (a''_{24})^{(4)} (T_{25}^{(2)}, s_{(24)}) | e^{-(\bar{M}_{24})^{(4)}s_{(24)}} e^{(\bar{M}_{24})^{(4)}s_{(24)}} \} ds_{(24)} \end{aligned}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\begin{aligned} |(G_{27})^{(1)} - (G_{27})^{(2)}| e^{-(\bar{M}_{24})^{(4)}t} &\leq \frac{1}{(\bar{M}_{24})^{(4)}} \left((a_{24})^{(4)} + (a'_{24})^{(4)} + (\hat{A}_{24})^{(4)} + \right. \\ &\left. (\hat{P}_{24})^{(4)} (\hat{k}_{24})^{(4)} \right) d \left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)}) \right) \end{aligned} \quad 338, 339$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ and $(\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)}t}$ respectively of \mathbb{R}_+ . 340

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(4)}$ and $(b'_i)^{(4)}$, $i = 24, 25, 26$ depend only on T_{25} and respectively on (G_{27}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 341

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(4)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{24})^{(4)})_1, ((\widehat{M}_{24})^{(4)})_2$ and $((\widehat{M}_{24})^{(4)})_3$: 342

Remark 3: if G_{24} is bounded, the same property have also G_{25} and G_{26} . indeed if

$$G_{24} < (\widehat{M}_{24})^{(4)} \text{ it follows } \frac{dG_{25}}{dt} \leq ((\widehat{M}_{24})^{(4)})_1 - (a'_{25})^{(4)}G_{25} \text{ and by integrating}$$

$$G_{25} \leq ((\widehat{M}_{24})^{(4)})_2 = G_{25}^0 + 2(a_{25})^{(4)}((\widehat{M}_{24})^{(4)})_1 / (a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq ((\widehat{M}_{24})^{(4)})_3 = G_{26}^0 + 2(a_{26})^{(4)}((\widehat{M}_{24})^{(4)})_2 / (a'_{26})^{(4)}$$

If G_{25} or G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 4: If G_{24} is bounded, from below, the same property holds for G_{25} and G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 343

Remark 5: If T_{24} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \rightarrow \infty$. 344

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 345

$$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4} \right) (1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_4 t} = \frac{1}{2} \text{ it results}$$

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2} \right)$, $t = \log \frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t \rightarrow \infty} (b''_{26})^{(4)}((G_{27})(t), t) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions ANALOGOUS inequalities hold also for G_{29} , G_{30} , T_{28} , T_{29} , T_{30}

346

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 347

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} \left[(\widehat{P}_{28})^{(5)} + ((\widehat{P}_{28})^{(5)} + G_j^0) e^{-\left(\frac{(\widehat{P}_{28})^{(5)} + G_j^0}{G_j^0} \right)} \right] \leq (\widehat{P}_{28})^{(5)} \quad 348$$

$$\frac{(b_i)^{(5)}}{(\bar{M}_{28})^{(5)}} \left[((\hat{Q}_{28})^{(5)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{28})^{(5)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)} \right] \leq (\hat{Q}_{28})^{(5)} \quad 349$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i into itself 350

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 351

$$d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{28})^{(5)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{28})^{(5)}t} \}$$

Indeed if we denote

Definition of $(\bar{G}_{31}), (\bar{T}_{31}) : ((\bar{G}_{31}), (\bar{T}_{31})) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$\begin{aligned} |\bar{G}_{28}^{(1)} - \bar{G}_{28}^{(2)}| &\leq \int_0^t (a_{28})^{(5)} |G_{29}^{(1)} - G_{29}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}} ds_{(28)} + \\ &\int_0^t \{ (a'_{28})^{(5)} |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{-(\bar{M}_{28})^{(5)}s_{(28)}} + \\ &(a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) |G_{28}^{(1)} - G_{28}^{(2)}| e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}} + \\ &G_{28}^{(2)} | (a''_{28})^{(5)} (T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)} (T_{29}^{(2)}, s_{(28)}) | e^{-(\bar{M}_{28})^{(5)}s_{(28)}} e^{(\bar{M}_{28})^{(5)}s_{(28)}} \} ds_{(28)} \end{aligned}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

352

$$\begin{aligned} |(G_{31})^{(1)} - (G_{31})^{(2)}| e^{-(\bar{M}_{28})^{(5)}t} &\leq \\ \frac{1}{(\bar{M}_{28})^{(5)}} \{ (a_{28})^{(5)} + (a'_{28})^{(5)} + (\hat{A}_{28})^{(5)} + \\ (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)} \} d \left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)}) \right) \end{aligned} \quad 353$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis (35,35,36) the result follows

Remark 1: The fact that we supposed $(a'_{28})^{(5)}$ and $(b'_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}$ and $(\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)}t}$ respectively of \mathbb{R}_+ . 354

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a'_i)^{(5)}$ and $(b'_i)^{(5)}$, $i = 28, 29, 30$ depend only on T_{29} and respectively on (G_{31}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 355

From GLOBAL EQUATIONS it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(5)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{28})^{(5)})_1, ((\widehat{M}_{28})^{(5)})_2$ and $((\widehat{M}_{28})^{(5)})_3$: 356

Remark 3: if G_{28} is bounded, the same property have also G_{29} and G_{30} . indeed if

$$G_{28} < (\widehat{M}_{28})^{(5)} \text{ it follows } \frac{dG_{29}}{dt} \leq ((\widehat{M}_{28})^{(5)})_1 - (a'_{29})^{(5)}G_{29} \text{ and by integrating}$$

$$G_{29} \leq ((\widehat{M}_{28})^{(5)})_2 = G_{29}^0 + 2(a_{29})^{(5)}((\widehat{M}_{28})^{(5)})_1 / (a'_{29})^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq ((\widehat{M}_{28})^{(5)})_3 = G_{30}^0 + 2(a_{30})^{(5)}((\widehat{M}_{28})^{(5)})_2 / (a'_{30})^{(5)}$$

If G_{29} or G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 4: If G_{28} is bounded, from below, the same property holds for G_{29} and G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 357

Remark 5: If T_{28} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(5)}((G_{31})(t), t)) = (b'_{29})^{(5)}$ then $T_{29} \rightarrow \infty$. 358

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)} \quad \text{359}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 360

$$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right) (1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_5 t} = \frac{1}{2} \text{ it results}$$

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), t = \log \frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t \rightarrow \infty} (b''_{30})^{(5)}((G_{31})(t), t) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

361

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 362

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\bar{M}_{32})^{(6)}} \left[(\hat{P}_{32})^{(6)} + ((\hat{P}_{32})^{(6)} + G_j^0) e^{-\left(\frac{(\hat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)} \right] \leq (\hat{P}_{32})^{(6)} \quad 363$$

$$\frac{(b_i)^{(6)}}{(\bar{M}_{32})^{(6)}} \left[((\hat{Q}_{32})^{(6)} + T_j^0) e^{-\left(\frac{(\hat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)} \right] \leq (\hat{Q}_{32})^{(6)} \quad 364$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i into itself 365

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 366

$$d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)}) \right) =$$

$$\sup_i \{ \max_{t \in \mathbb{R}_+} |G_i^{(1)}(t) - G_i^{(2)}(t)| e^{-(\bar{M}_{32})^{(6)}t}, \max_{t \in \mathbb{R}_+} |T_i^{(1)}(t) - T_i^{(2)}(t)| e^{-(\bar{M}_{32})^{(6)}t} \}$$

Indeed if we denote

$$\underline{\text{Definition of}} (\overline{G_{35}}, \overline{T_{35}}) : (\overline{G_{35}}, \overline{T_{35}}) = \mathcal{A}^{(6)}((G_{35}), (T_{35}))$$

It results

$$\begin{aligned} |\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}| &\leq \int_0^t (a_{32})^{(6)} |G_{33}^{(1)} - G_{33}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}} ds_{(32)} + \\ &\int_0^t \{ (a'_{32})^{(6)} |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{-(\bar{M}_{32})^{(6)}s_{(32)}} + \\ &(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) |G_{32}^{(1)} - G_{32}^{(2)}| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}} + \\ &G_{32}^{(2)} |(a''_{32})^{(6)} (T_{33}^{(1)}, s_{(32)}) - (a''_{32})^{(6)} (T_{33}^{(2)}, s_{(32)})| e^{-(\bar{M}_{32})^{(6)}s_{(32)}} e^{(\bar{M}_{32})^{(6)}s_{(32)}} \} ds_{(32)} \end{aligned}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$ 367

From the hypotheses it follows

$$\begin{aligned} |(G_{35})^{(1)} - (G_{35})^{(2)}| e^{-(\bar{M}_{32})^{(6)}t} &\leq \quad 368 \\ \frac{1}{(\bar{M}_{32})^{(6)}} &((a_{32})^{(6)} + (a'_{32})^{(6)} + (\bar{A}_{32})^{(6)} + \\ &(\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}) d \left(((G_{35})^{(1)}, (T_{35})^{(1)}), (G_{35})^{(2)}, (T_{35})^{(2)} \right) \end{aligned}$$

And analogous inequalities for G_i and T_i . Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis, in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}$ and $(\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)}t}$ respectively of \mathbb{R}_+ . 369

If instead of proving the existence of the solution on \mathbb{R}_+ , we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(6)}$ and $(b''_i)^{(6)}$, $i = 32, 33, 34$ depend only on T_{33} and respectively on (G_{35}) (and not on t) and hypothesis can be replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 370

From 69 to 32 it results

$$G_i(t) \geq G_i^0 e^{-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}} \geq 0$$

$$T_i(t) \geq T_i^0 e^{-(b_i')^{(6)}t} > 0 \text{ for } t > 0$$

Definition of $((\widehat{M}_{32})^{(6)})_1, ((\widehat{M}_{32})^{(6)})_2$ and $((\widehat{M}_{32})^{(6)})_3$: 371

Remark 3: if G_{32} is bounded, the same property have also G_{33} and G_{34} . indeed if

$$G_{32} < (\widehat{M}_{32})^{(6)} \text{ it follows } \frac{dG_{33}}{dt} \leq ((\widehat{M}_{32})^{(6)})_1 - (a'_{33})^{(6)}G_{33} \text{ and by integrating}$$

$$G_{33} \leq ((\widehat{M}_{32})^{(6)})_2 = G_{33}^0 + 2(a_{33})^{(6)}((\widehat{M}_{32})^{(6)})_1 / (a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq ((\widehat{M}_{32})^{(6)})_3 = G_{34}^0 + 2(a_{34})^{(6)}((\widehat{M}_{32})^{(6)})_2 / (a'_{34})^{(6)}$$

If G_{33} or G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 4: If G_{32} is bounded, from below, the same property holds for G_{33} and G_{34} . The proof is 372
 analogous with the preceding one. An analogous property is true if G_{33} is bounded from below.

Remark 5: If T_{32} is bounded from below and $\lim_{t \rightarrow \infty} ((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then 373
 $T_{33} \rightarrow \infty$.

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}((G_{35})(t), t) < \varepsilon_6, T_{32}(t) > (m)^{(6)} \tag{374}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 375

$$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6} \right) (1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t} \text{ If we take } t \text{ such that } e^{-\varepsilon_6 t} = \frac{1}{2} \text{ it results}$$

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2} \right)$, $t = \log \frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is
 unbounded. The same property holds for T_{34} if $\lim_{t \rightarrow \infty} (b''_{34})^{(6)}((G_{35})(t), t(t), t) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions

376

Behavior of the solutions

377

If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

(a) $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying

$$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$$

$$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 378

(b) By $(v_1)^{(1)} > 0, (v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0, (u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 379

By $(\bar{v}_1)^{(1)} > 0, (\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0, (\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}(v^{(1)})^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}(u^{(1)})^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 380

(c) If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$$(m_2)^{(1)} = (v_0)^{(1)}, (m_1)^{(1)} = (v_1)^{(1)}, \text{ if } (v_0)^{(1)} < (v_1)^{(1)}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (\bar{v}_1)^{(1)}, \text{ if } (v_1)^{(1)} < (v_0)^{(1)} < (\bar{v}_1)^{(1)},$$

$$\text{and } (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}$$

$$(m_2)^{(1)} = (v_1)^{(1)}, (m_1)^{(1)} = (v_0)^{(1)}, \text{ if } (\bar{v}_1)^{(1)} < (v_0)^{(1)}$$

and analogously 381

$$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \text{ if } (u_0)^{(1)} < (u_1)^{(1)}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, \text{ if } (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$$

$$\text{and } (u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}$$

$$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, \text{ if } (\bar{u}_1)^{(1)} < (u_0)^{(1)} \text{ where } (u_1)^{(1)}, (\bar{u}_1)^{(1)} 382$$

are defined respectively

Then the solution satisfies the inequalities 383

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{13}(t) \leq G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \leq G_{14}(t) \leq \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)} ((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})} \left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t} \right] + G_{15}^0 e^{-(S_2)^{(1)}t} \right) \leq G_{15}(t) \leq 384$$

$$\frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)} ((S_1)^{(1)} - (a'_{15})^{(1)})} \left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t} \right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}$$

$$T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} 385$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \leq T_{13}(t) \leq \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} 386$$

$$\frac{(b_{15})^{(1)}T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})} \left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t} \right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \leq T_{15}(t) \leq \quad 387$$

$$\frac{(a_{15})^{(1)}T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})} \left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t} \right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 388

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions 389

If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 390

(d) $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \leq -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \leq -(\sigma_1)^{(2)} \quad 391$$

$$-(\tau_2)^{(2)} \leq -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}((G_{19}), t) - (b''_{17})^{(2)}((G_{19}), t) \leq -(\tau_1)^{(2)} \quad 392$$

Definition of $(v_1)^{(2)}, (v_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 393

By $(v_1)^{(2)} > 0, (v_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0, (u_2)^{(2)} < 0$ the roots 394

(e) of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 395

$$\text{and } (b_{14})^{(2)}(u^{(2)})^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \text{ and} \quad 396$$

Definition of $(\bar{v}_1)^{(2)}, (\bar{v}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 397

By $(\bar{v}_1)^{(2)} > 0, (\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0, (\bar{u}_2)^{(2)} < 0$ the 398

roots of the equations $(a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 399

$$\text{and } (b_{17})^{(2)}(u^{(2)})^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0 \quad 400$$

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 401

(f) If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 402

$$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)}, \text{ if } (v_0)^{(2)} < (v_1)^{(2)} \quad 403$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \text{ if } (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)}, \quad 404$$

$$\text{and } (v_0)^{(2)} = \frac{a_{16}^0}{a_{17}^0}$$

$$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)}, \text{ if } (\bar{v}_1)^{(2)} < (v_0)^{(2)} \quad 405$$

and analogously 406

$$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}, \text{ if } (u_0)^{(2)} < (u_1)^{(2)}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \text{ if } (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$$

$$\text{and } (u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}$$

$$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}, \text{ if } (\bar{u}_1)^{(2)} < (u_0)^{(2)} \quad 407$$

Then the solution satisfies the inequalities 408

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{16}(t) \leq G_{16}^0 e^{(S_1)^{(2)}t}$$

$(p_i)^{(2)}$ is defined 409

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \leq G_{17}(t) \leq \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t} \quad 410$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)} ((S_1)^{(2)} - (p_{16})^{(2)}) - (S_2)^{(2)}} \right) \left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t} \right] + G_{18}^0 e^{-(S_2)^{(2)}t} \leq G_{18}(t) \leq \quad 411$$

$$\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)} ((S_1)^{(2)} - (a'_{18})^{(2)})} \left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t} \right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 412$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \leq T_{16}(t) \leq \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 413$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)} ((R_1)^{(2)} - (b'_{18})^{(2)})} \left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t} \right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \leq T_{18}(t) \leq \quad 414$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)} ((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})} \left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t} \right] + T_{18}^0 e^{-(R_2)^{(2)}t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 415

$$\text{Where } (S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)} \quad 416$$

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 417$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

418

Behavior of the solutions 419

If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

(a) $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \leq -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \leq -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \leq -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G, t) - (b''_{21})^{(3)}((G_{23}), t) \leq -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 420

(b) By $(v_1)^{(3)} > 0, (v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0, (u_2)^{(3)} < 0$ the roots of the

equations $(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}(u^{(3)})^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0, (\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0, (\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}(u^{(3)})^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 421

(c) If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}, \text{ if } (v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}, \text{ if } (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)},$

and $(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}, \text{ if } (\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 422

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}, \text{ if } (u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}, \text{ if } (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}, \text{ and } (u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}, \text{ if } (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined 423

$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t}$ 424

$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)} ((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})} \left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t} \right] + G_{22}^0 e^{-(S_2)^{(3)}t} \right) \leq G_{22}(t) \leq \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)} ((S_1)^{(3)} - (a'_{22})^{(3)})} \left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t} \right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}$ 425

$T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$ 426

$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$ 427

$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)} ((R_1)^{(3)} - (b'_{22})^{(3)})} \left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t} \right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \leq T_{22}(t) \leq$ 428

$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)} ((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})} \left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t} \right] + T_{22}^0 e^{-(R_2)^{(3)}t}$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 429

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

430

431

Behavior of the solutions

432

If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

(d) $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \leq -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \leq -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}((G_{27}), t) - (b''_{25})^{(4)}((G_{27}), t) \leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$:

433

(e) By $(v_1)^{(4)} > 0, (v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0, (u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}(u^{(4)})^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)}, (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$:

434

By $(\bar{v}_1)^{(4)} > 0, (\bar{v}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0, (\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

$$\text{and } (b_{25})^{(4)}(u^{(4)})^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$$

436

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (v_0)^{(4)}$:-

(f) If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (v_0)^{(4)}, (m_1)^{(4)} = (v_1)^{(4)}, \text{ if } (v_0)^{(4)} < (v_1)^{(4)}$$

$$(m_2)^{(4)} = (v_1)^{(4)}, (m_1)^{(4)} = (\bar{v}_1)^{(4)}, \text{ if } (v_4)^{(4)} < (v_0)^{(4)} < (\bar{v}_1)^{(4)},$$

$$\text{and } (v_0)^{(4)} = \frac{a_{24}^0}{a_{25}^0}$$

$$(m_2)^{(4)} = (v_4)^{(4)}, (m_1)^{(4)} = (v_0)^{(4)}, \text{ if } (\bar{v}_4)^{(4)} < (v_0)^{(4)}$$

and analogously

437

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \text{ if } (u_0)^{(4)} < (u_1)^{(4)}$$

438

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \text{ if } (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

$$\text{and } (u_0)^{(4)} = \frac{r_{24}^0}{r_{25}^0}$$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \text{ if } (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$ are defined by 59 and 64 respectively

Then the solution satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)} ((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t} \right] + G_{26}^0 e^{-(S_2)^{(4)}t} \right) \leq G_{26}(t) \leq \left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)} ((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)}t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)} ((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \leq T_{26}(t) \leq$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)} ((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t} \right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:-

$$\text{Where } (S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$$

$$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$$

$$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$$

$$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$$

Behavior of the solutions

If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

(g) $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \leq -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \leq -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \leq -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \leq -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$:

(h) By $(v_1)^{(5)} > 0, (v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0, (u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$ and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$:

By $(\bar{v}_1)^{(5)} > 0, (\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0, (\bar{u}_2)^{(5)} < 0$ the

roots of the equations $(a_{29})^{(5)}(v^{(5)})^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
 and $(b_{29})^{(5)}(u^{(5)})^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

(i) If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}, \text{ if } (v_0)^{(5)} < (v_1)^{(5)}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}, \text{ if } (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)},$$

$$\text{and } (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}$$

$$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}, \text{ if } (\bar{v}_1)^{(5)} < (v_0)^{(5)}$$

and analogously

457

$$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}, \text{ if } (u_0)^{(5)} < (u_1)^{(5)}$$

$$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}, \text{ if } (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$$

$$\text{and } (u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}$$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \text{ if } (\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$ are defined respectively

Then the solution satisfies the inequalities

458

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t}$$

459

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)} ((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})} \left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t} \right] + G_{30}^0 e^{-(S_2)^{(5)}t} \right) \leq G_{30}(t) \leq$$

$$\frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)} ((S_1)^{(5)} - (a'_{30})^{(5)})} \left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t} \right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}$$

460

461

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}}$$

462

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}$$

463

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)} ((R_1)^{(5)} - (b'_{30})^{(5)})} \left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t} \right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \leq T_{30}(t) \leq$$

464

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)} ((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})} \left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t} \right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:-

465

$$\text{Where } (S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$$

$$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$$

$$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$$

$$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$$

Behavior of the solutions

466

If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

(j) $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \leq -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \leq -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$:

467

(k) By $(v_1)^{(6)} > 0, (v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0, (u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$ and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$:

468

By $(\bar{v}_1)^{(6)} > 0, (\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0, (\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$ and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

(l) If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}, \text{ if } (v_0)^{(6)} < (v_1)^{(6)}$$

470

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}, \text{ if } (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)},$$

$$\text{and } (v_0)^{(6)} = \frac{a_{32}^0}{a_{33}^0}$$

$$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}, \text{ if } (\bar{v}_1)^{(6)} < (v_0)^{(6)}$$

and analogously

471

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \text{ if } (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \text{ if } (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

$$\text{and } (u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}$$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}$, if $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$ are defined respectively

Then the solution satisfies the inequalities

472

$$G_{32}^0 e^{((s_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{32}(t) \leq G_{32}^0 e^{(s_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((s_1)^{(6)} - (p_{32})^{(6)})t} \leq G_{33}(t) \leq \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(s_1)^{(6)}t}$$

473

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)} (S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)}} \left[e^{(S_1)^{(6)} - (p_{32})^{(6)} t} - e^{-(S_2)^{(6)} t} \right] + G_{34}^0 e^{-(S_2)^{(6)} t} \leq G_{34}(t) \leq \right. \tag{474}$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)} (S_1)^{(6)} - (a'_{34})^{(6)}} \left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t} \right] + G_{34}^0 e^{-(a'_{34})^{(6)} t} \right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)}) t}} \tag{475}$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \leq T_{32}(t) \leq \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)}) t} \tag{476}$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)} (R_1)^{(6)} - (b'_{34})^{(6)}} \left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t} \right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \leq T_{34}(t) \leq \tag{477}$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)} (R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)}} \left[e^{((R_1)^{(6)} + (r_{32})^{(6)}) t} - e^{-(R_2)^{(6)} t} \right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 478

Where $(S_1)^{(6)} = (a_{32})^{(6)} (m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)} (\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

479

Proof : From GLOBAL EQUATIONS we obtain 480

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)} (T_{14}, t) \right) - (a''_{14})^{(1)} (T_{14}, t) v^{(1)} - (a_{14})^{(1)} v^{(1)}$$

Definition of $v^{(1)}$:- 481

$$\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$$

It follows

$$- \left((a_{14})^{(1)} (v^{(1)})^2 + (\sigma_2)^{(1)} v^{(1)} - (a_{13})^{(1)} \right) \leq \frac{dv^{(1)}}{dt} \leq - \left((a_{14})^{(1)} (v^{(1)})^2 + (\sigma_1)^{(1)} v^{(1)} - (a_{13})^{(1)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(1)}, (v_0)^{(1)}$:-

(a) For $0 < \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (v_1)^{(1)} < (\bar{v}_1)^{(1)}$

$$v^{(1)}(t) \geq \frac{(v_1)^{(1)} + (C)^{(1)} (v_2)^{(1)} e^{-(a_{14})^{(1)} (v_1)^{(1)} - (v_0)^{(1)} t}}{1 + (C)^{(1)} e^{-(a_{14})^{(1)} (v_1)^{(1)} - (v_0)^{(1)} t}}, \quad \boxed{(C)^{(1)} = \frac{(v_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (v_2)^{(1)}}$$

it follows $(v_0)^{(1)} \leq v^{(1)}(t) \leq (v_1)^{(1)}$

In the same manner , we get

482

$$v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)} (\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)} (\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}] t}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} (\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}] t}}, \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{v}_1)^{(1)} - (v_0)^{(1)}}{(v_0)^{(1)} - (\bar{v}_2)^{(1)}}$$

From which we deduce $(v_0)^{(1)} \leq v^{(1)}(t) \leq (\bar{v}_1)^{(1)}$

(b) If $0 < (v_1)^{(1)} < (v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{v}_1)^{(1)}$ we find like in the previous case, 483

$$(v_1)^{(1)} \leq \frac{(v_1)^{(1)} + (C)^{(1)} (v_2)^{(1)} e^{[-(a_{14})^{(1)} ((v_1)^{(1)} - (v_2)^{(1)}) t]}}{1 + (C)^{(1)} e^{[-(a_{14})^{(1)} ((v_1)^{(1)} - (v_2)^{(1)}) t]}} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)} (\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)} (\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}] t}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} (\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}] t}} \leq (\bar{v}_1)^{(1)}$$

(c) If $0 < (v_1)^{(1)} \leq (\bar{v}_1)^{(1)} \leq \boxed{(v_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 484

$$(v_1)^{(1)} \leq v^{(1)}(t) \leq \frac{(\bar{v}_1)^{(1)} + (\bar{C})^{(1)} (\bar{v}_2)^{(1)} e^{[-(a_{14})^{(1)} (\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}] t}}{1 + (\bar{C})^{(1)} e^{[-(a_{14})^{(1)} (\bar{v}_1)^{(1)} - (\bar{v}_2)^{(1)}] t}} \leq (v_0)^{(1)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in GLOBAL E486QUATIONS we get easily the result stated in the theorem.

Particular case : 485

If $(a'_{13})^{(1)} = (a'_{14})^{(1)}$, then $(\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(v_1)^{(1)} = (\bar{v}_1)^{(1)}$ if in addition $(v_0)^{(1)} = (v_1)^{(1)}$ then $v^{(1)}(t) = (v_0)^{(1)}$ and as a consequence $G_{13}(t) = (v_0)^{(1)} G_{14}(t)$ this also defines $(v_0)^{(1)}$ for the special case

Analogously if $(b'_{13})^{(1)} = (b'_{14})^{(1)}$, then $(\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(v_1)^{(1)}$ and $(\bar{v}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

486

we obtain

487

$$\frac{dv^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)} (T_{17}, t) \right) - (a'_{17})^{(2)} (T_{17}, t) v^{(2)} - (a_{17})^{(2)} v^{(2)}$$

Definition of $v^{(2)}$:-
$$v^{(2)} = \frac{G_{16}}{G_{17}}$$
 488

It follows 489

$$-\left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{dv^{(2)}}{dt} \leq -\left((a_{17})^{(2)}(v^{(2)})^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 490

Definition of $(\bar{v}_1)^{(2)}, (v_0)^{(2)}$:-

(d) For $0 < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (v_1)^{(2)} < (\bar{v}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}(v_1)^{(2)} - (v_0)^{(2)}]t}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}(v_1)^{(2)} - (v_0)^{(2)}]t}}, \quad (C)^{(2)} = \frac{(v_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (v_2)^{(2)}}$$

it follows $(v_0)^{(2)} \leq v^{(2)}(t) \leq (v_1)^{(2)}$

In the same manner, we get 491

$$v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}(\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}]t}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}(\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}]t}}, \quad (\bar{C})^{(2)} = \frac{(\bar{v}_1)^{(2)} - (v_0)^{(2)}}{(v_0)^{(2)} - (\bar{v}_2)^{(2)}}$$

From which we deduce $(v_0)^{(2)} \leq v^{(2)}(t) \leq (\bar{v}_1)^{(2)}$ 492

(e) If $0 < (v_1)^{(2)} < (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{v}_1)^{(2)}$ we find like in the previous case, 493

$$(v_1)^{(2)} \leq \frac{(v_1)^{(2)} + (C)^{(2)}(v_2)^{(2)}e^{[-(a_{17})^{(2)}(v_1)^{(2)} - (v_2)^{(2)}]t}}{1 + (C)^{(2)}e^{[-(a_{17})^{(2)}(v_1)^{(2)} - (v_2)^{(2)}]t}} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}(\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}]t}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}(\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}]t}} \leq (\bar{v}_1)^{(2)}$$

(f) If $0 < (v_1)^{(2)} \leq (\bar{v}_1)^{(2)} \leq (v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 494

$$(v_1)^{(2)} \leq v^{(2)}(t) \leq \frac{(\bar{v}_1)^{(2)} + (\bar{C})^{(2)}(\bar{v}_2)^{(2)}e^{[-(a_{17})^{(2)}(\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}]t}}{1 + (\bar{C})^{(2)}e^{[-(a_{17})^{(2)}(\bar{v}_1)^{(2)} - (\bar{v}_2)^{(2)}]t}} \leq (v_0)^{(2)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(2)}(t)$:- 495

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}$$

In a completely analogous way, we obtain 496

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}$$

. 497

Particular case : 498

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}$, then $(\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(v_1)^{(2)} = (\bar{v}_1)^{(2)}$ if in addition $(v_0)^{(2)} = (v_1)^{(2)}$ then $v^{(2)}(t) = (v_0)^{(2)}$ and as a consequence $G_{16}(t) = (v_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}$, then $(\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$ if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(v_1)^{(2)}$ and $(\bar{v}_1)^{(2)}$

499

From GLOBAL EQUATIONS we obtain

500

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t) \right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:-
$$v^{(3)} = \frac{G_{20}}{G_{21}}$$

501

It follows

$$- \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} \right) \leq \frac{dv^{(3)}}{dt} \leq - \left((a_{21})^{(3)}(v^{(3)})^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} \right)$$

502

From which one obtains

(a) For $0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \geq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}(v_1)^{(3)} - (v_0)^{(3)}]t}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}(v_1)^{(3)} - (v_0)^{(3)}]t}}, \quad (C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}$$

it follows $(v_0)^{(3)} \leq v^{(3)}(t) \leq (v_1)^{(3)}$

In the same manner, we get

503

$$v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}(\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}]t}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}(\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}]t}}, \quad (\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \leq v^{(3)}(t) \leq (\bar{v}_1)^{(3)}$

(b) If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case,

504

$$(v_1)^{(3)} \leq \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)} e^{[-(a_{21})^{(3)}(v_1)^{(3)} - (v_2)^{(3)}]t}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}(v_1)^{(3)} - (v_2)^{(3)}]t}} \leq v^{(3)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}(\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}]t}}{1 + (\bar{C})^{(3)} e^{[-(a_{21})^{(3)}(\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}]t}} \leq (\bar{v}_1)^{(3)}$$

(c) If $0 < (v_1)^{(3)} \leq (\bar{v}_1)^{(3)} \leq (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain

505

$$(v_1)^{(3)} \leq v^{(3)}(t) \leq \frac{(\bar{v}_1)^{(3)} + (C)^{(3)}(\bar{v}_2)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}}{1 + (C)^{(3)} e^{[-(a_{21})^{(3)}((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)})t]}} \leq (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \leq v^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}$, then $(\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(v_1)^{(3)} = (\bar{v}_1)^{(3)}$ if in addition $(v_0)^{(3)} = (v_1)^{(3)}$ then $v^{(3)}(t) = (v_0)^{(3)}$ and as a consequence $G_{20}(t) = (v_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}$, then $(\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(v_1)^{(3)}$ and $(\bar{v}_1)^{(3)}$

506

: From GLOBAL EQUATIONS we obtain

507

$$\frac{dv^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t) \right) - (a''_{25})^{(4)}(T_{25}, t)v^{(4)} - (a_{25})^{(4)}v^{(4)}$$

Definition of $v^{(4)}$:-

$$\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$$

508

It follows

$$- \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} \right) \leq \frac{dv^{(4)}}{dt} \leq - \left((a_{25})^{(4)}(v^{(4)})^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

(d) For $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}}{4 + (C)^{(4)} e^{[-(a_{25})^{(4)}((v_1)^{(4)} - (v_0)^{(4)})t]}} , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get

509

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{c})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} (\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}] t}}{4 + (\bar{c})^{(4)} e^{[-(a_{25})^{(4)} (\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}] t}}, \quad \boxed{(\bar{c})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

(e) If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 510

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (c)^{(4)} (v_2)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}}{1 + (c)^{(4)} e^{[-(a_{25})^{(4)} ((v_1)^{(4)} - (v_2)^{(4)}) t]}} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{c})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} (\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}] t}}{1 + (\bar{c})^{(4)} e^{[-(a_{25})^{(4)} (\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}] t}} \leq (\bar{v}_1)^{(4)}$$

511
512

(f) If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{c})^{(4)} (\bar{v}_2)^{(4)} e^{[-(a_{25})^{(4)} (\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}] t}}{1 + (\bar{c})^{(4)} e^{[-(a_{25})^{(4)} (\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}] t}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c), we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}$, then $(\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(v_1)^{(4)} = (\bar{v}_1)^{(4)}$ if in addition $(v_0)^{(4)} = (v_1)^{(4)}$ then $v^{(4)}(t) = (v_0)^{(4)}$ and as a consequence $G_{24}(t) = (v_0)^{(4)} G_{25}(t)$ **this also defines $(v_0)^{(4)}$ for the special case .** 513

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}$, then $(\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(v_1)^{(4)}$ and $(\bar{v}_1)^{(4)}$, **and definition of $(u_0)^{(4)}$.**

514
515

From GLOBAL EQUATIONS we obtain

$$\frac{dv^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)} (T_{29}, t) \right) - (a''_{29})^{(5)} (T_{29}, t) v^{(5)} - (a_{29})^{(5)} v^{(5)}$$

Definition of $v^{(5)}$:- $\boxed{v^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$- \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_2)^{(5)} v^{(5)} - (a_{28})^{(5)} \right) \leq \frac{dv^{(5)}}{dt} \leq - \left((a_{29})^{(5)} (v^{(5)})^2 + (\sigma_1)^{(5)} v^{(5)} - (a_{28})^{(5)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(5)}, (v_0)^{(5)}$:-

(g) For $0 < \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (v_1)^{(5)} < (\bar{v}_1)^{(5)}$

$$v^{(5)}(t) \geq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}}{5 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_0)^{(5)})t]}} , \quad \boxed{(C)^{(5)} = \frac{(v_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (v_2)^{(5)}}$$

it follows $(v_0)^{(5)} \leq v^{(5)}(t) \leq (v_1)^{(5)}$

In the same manner , we get

516

$$v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{5 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{v}_1)^{(5)} - (v_0)^{(5)}}{(v_0)^{(5)} - (\bar{v}_2)^{(5)}}$$

From which we deduce $(v_0)^{(5)} \leq v^{(5)}(t) \leq (\bar{v}_5)^{(5)}$

(h) If $0 < (v_1)^{(5)} < (v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{v}_1)^{(5)}$ we find like in the previous case,

517

$$(v_1)^{(5)} \leq \frac{(v_1)^{(5)} + (C)^{(5)}(v_2)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}}{1 + (C)^{(5)} e^{[-(a_{29})^{(5)}((v_1)^{(5)} - (v_2)^{(5)})t]}} \leq v^{(5)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (\bar{v}_1)^{(5)}$$

(i) If $0 < (v_1)^{(5)} \leq (\bar{v}_1)^{(5)} \leq \boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

518

$$(v_1)^{(5)} \leq v^{(5)}(t) \leq \frac{(\bar{v}_1)^{(5)} + (\bar{C})^{(5)}(\bar{v}_2)^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}}{1 + (\bar{C})^{(5)} e^{[-(a_{29})^{(5)}((\bar{v}_1)^{(5)} - (\bar{v}_2)^{(5)})t]}} \leq (v_0)^{(5)}$$

519

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(5)}(t)$:-

$$(m_2)^{(5)} \leq v^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{v^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

Particular case :

If $(a''_{28})^{(5)} = (a''_{29})^{(5)}$, then $(\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(v_1)^{(5)} = (\bar{v}_1)^{(5)}$ if in addition $(v_0)^{(5)} = (v_5)^{(5)}$ then $v^{(5)}(t) = (v_0)^{(5)}$ and as a consequence $G_{28}(t) = (v_0)^{(5)}G_{29}(t)$ **this also defines $(v_0)^{(5)}$ for the special case .**

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}$, then $(\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(v_1)^{(5)}$ and $(\bar{v}_1)^{(5)}$, **and definition of $(u_0)^{(5)}$.**

we obtain

520
521

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t) \right) - (a''_{33})^{(6)}(T_{33}, t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:-
$$v^{(6)} = \frac{G_{32}}{G_{33}}$$

It follows

$$- \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} \right) \leq \frac{dv^{(6)}}{dt} \leq - \left((a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} \right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(6)}, (v_0)^{(6)}$:-

(j) For $0 < \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (v_1)^{(6)} < (\bar{v}_1)^{(6)}$

$$v^{(6)}(t) \geq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_0)^{(6)})t]}} , \quad \boxed{(C)^{(6)} = \frac{(v_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (v_2)^{(6)}}$$

it follows $(v_0)^{(6)} \leq v^{(6)}(t) \leq (v_1)^{(6)}$

In the same manner , we get

522

$$v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{v}_1)^{(6)} - (v_0)^{(6)}}{(v_0)^{(6)} - (\bar{v}_2)^{(6)}}$$

523

From which we deduce $(v_0)^{(6)} \leq v^{(6)}(t) \leq (\bar{v}_1)^{(6)}$

(k) If $0 < (v_1)^{(6)} < (v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{v}_1)^{(6)}$ we find like in the previous case,

524

$$(v_1)^{(6)} \leq \frac{(v_1)^{(6)} + (C)^{(6)}(v_2)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}}{1 + (C)^{(6)} e^{[-(a_{33})^{(6)}((v_1)^{(6)} - (v_2)^{(6)})t]}} \leq v^{(6)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (\bar{v}_1)^{(6)}$$

(l) If $0 < (v_1)^{(6)} \leq (\bar{v}_1)^{(6)} \leq \boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain

525

$$(v_1)^{(6)} \leq v^{(6)}(t) \leq \frac{(\bar{v}_1)^{(6)} + (\bar{C})^{(6)}(\bar{v}_2)^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}}{1 + (\bar{C})^{(6)} e^{[-(a_{33})^{(6)}((\bar{v}_1)^{(6)} - (\bar{v}_2)^{(6)})t]}} \leq (v_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in GLOBAL EQUATIONS we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}$, then $(\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(v_1)^{(6)} = (\bar{v}_1)^{(6)}$ if in addition $(v_0)^{(6)} = (v_1)^{(6)}$ then $v^{(6)}(t) = (v_0)^{(6)}$ and as a consequence $G_{32}(t) = (v_0)^{(6)}G_{33}(t)$ **this also defines $(v_0)^{(6)}$ for the special case .**

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}$, then $(\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of $(u_0)^{(6)}$.**

527

526
527

We can prove the following

528

Theorem 3: If $(a'_i)^{(1)}$ and $(b'_i)^{(1)}$ are independent on t , and the conditions

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0 ,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

with $(p_{13})^{(1)}, (r_{14})^{(1)}$ as defined, then the system

529

If $(a'_i)^{(2)}$ and $(b'_i)^{(2)}$ are independent on t , and the conditions

530.

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$$

531

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$$

532

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0 ,$$

533

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$$

534

with $(p_{16})^{(2)}, (r_{17})^{(2)}$ as defined are satisfied , then the system

If $(a'_i)^{(3)}$ and $(b'_i)^{(3)}$ are independent on t , and the conditions

535

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0 ,$$

$$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$$

with $(p_{20})^{(3)}, (r_{21})^{(3)}$ as defined are satisfied , then the system

If $(a''_i)^{(4)}$ and $(b'_i)^{(4)}$ are independent on t , and the conditions 536

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$$

$$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0 ,$$

$$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$$

with $(p_{24})^{(4)}, (r_{25})^{(4)}$ as defined are satisfied , then the system

If $(a''_i)^{(5)}$ and $(b'_i)^{(5)}$ are independent on t , and the conditions 537

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$$

$$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0 ,$$

$$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$$

with $(p_{28})^{(5)}, (r_{29})^{(5)}$ as defined satisfied , then the system

If $(a''_i)^{(6)}$ and $(b'_i)^{(6)}$ are independent on t , and the conditions 538

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$$

$$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0 ,$$

$$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$$
539

with $(p_{32})^{(6)}, (r_{33})^{(6)}$ as defined are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - [(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})]G_{13} = 0$$
540

$$(a_{14})^{(1)}G_{13} - [(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})]G_{14} = 0$$
541

$$(a_{15})^{(1)}G_{14} - [(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})]G_{15} = 0$$
542

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)]T_{13} = 0$$
543

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)]T_{14} = 0$$
544

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)]T_{15} = 0$$
545

has a unique positive solution , which is an equilibrium solution for the system 546

$$(a_{16})^{(2)}G_{17} - [(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})]G_{16} = 0$$
547

$$(a_{17})^{(2)}G_{16} - [(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})]G_{17} = 0$$
548

$$(a_{18})^{(2)}G_{17} - [(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})]G_{18} = 0 \quad 549$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})]T_{16} = 0 \quad 550$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})]T_{17} = 0 \quad 551$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})]T_{18} = 0 \quad 552$$

has a unique positive solution , which is an equilibrium solution for 553

$$(a_{20})^{(3)}G_{21} - [(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})]G_{20} = 0 \quad 554$$

$$(a_{21})^{(3)}G_{20} - [(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})]G_{21} = 0 \quad 555$$

$$(a_{22})^{(3)}G_{21} - [(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})]G_{22} = 0 \quad 556$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})]T_{20} = 0 \quad 557$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})]T_{21} = 0 \quad 558$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})]T_{22} = 0 \quad 559$$

has a unique positive solution , which is an equilibrium solution 560

$$(a_{24})^{(4)}G_{25} - [(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})]G_{24} = 0 \quad 561$$

$$(a_{25})^{(4)}G_{24} - [(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})]G_{25} = 0 \quad 563$$

$$(a_{26})^{(4)}G_{25} - [(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})]G_{26} = 0 \quad 564$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27})]T_{24} = 0 \quad 565$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27})]T_{25} = 0 \quad 566$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27})]T_{26} = 0 \quad 567$$

has a unique positive solution , which is an equilibrium solution for the system 568

$$(a_{28})^{(5)}G_{29} - [(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})]G_{28} = 0 \quad 569$$

$$(a_{29})^{(5)}G_{28} - [(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})]G_{29} = 0 \quad 570$$

$$(a_{30})^{(5)}G_{29} - [(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})]G_{30} = 0 \quad 571$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 572$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 573$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 574$$

has a unique positive solution , which is an equilibrium solution for the system 575

$$(a_{32})^{(6)}G_{33} - [(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})]G_{32} = 0 \quad 576$$

$$(a_{33})^{(6)}G_{32} - [(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})]G_{33} = 0 \quad 577$$

$$(a_{34})^{(6)}G_{33} - [(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})]G_{34} = 0 \quad 578$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 579$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 580$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 584$$

has a unique positive solution , which is an equilibrium solution for the system 582

583

584

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

585

(a) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$F(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

586

587

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

588

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

589

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

560

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Definition and uniqueness of T_{14}^* :- 561

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value, we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)} + (a'_{13})^{(1)}(T_{14}^*)]} , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)} + (a'_{15})^{(1)}(T_{14}^*)]}$$

Definition and uniqueness of T_{17}^* :- 562

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T_{17}^* for which $f(T_{17}^*) = 0$. With this value, we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)} + (a'_{16})^{(2)}(T_{17}^*)]} , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)} + (a'_{18})^{(2)}(T_{17}^*)]}$$

Definition and uniqueness of T_{21}^* :- 564

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(3)}(T_{21})$ being increasing, it follows that there exists a unique T_{21}^* for which $f(T_{21}^*) = 0$. With this value, we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)} + (a'_{20})^{(3)}(T_{21}^*)]} , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)} + (a'_{22})^{(3)}(T_{21}^*)]}$$

565

Definition and uniqueness of T_{25}^* :- 566

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T_{25}^* for which $f(T_{25}^*) = 0$. With this value, we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)} + (a'_{24})^{(4)}(T_{25}^*)]} , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)} + (a'_{26})^{(4)}(T_{25}^*)]}$$

Definition and uniqueness of T_{29}^* :- 567

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T_{29}^* for which $f(T_{29}^*) = 0$. With this value, we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)} + (a'_{28})^{(5)}(T_{29}^*)]} , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)} + (a'_{30})^{(5)}(T_{29}^*)]}$$

Definition and uniqueness of T_{33}^* :- 568

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value, we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]}$$

(e) By the same argument, the equations 92,93 admit solutions G_{13}, G_{14} if 569

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - \\ [(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

(f) By the same argument, the equations 92,93 admit solutions G_{16}, G_{17} if 570

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - \\ [(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{17}^* such that $\varphi((G_{19})^*) = 0$ 571

(g) By the same argument, the concatenated equations admit solutions G_{20}, G_{21} if 572

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - \\ [(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$ 573

(h) By the same argument, the equations of modules admit solutions G_{24}, G_{25} if 574

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - \\ [(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^* such that $\varphi((G_{27})^*) = 0$

(i) By the same argument, the equations (modules) admit solutions G_{28}, G_{29} if 575

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - \\ [(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^* such that $\varphi((G_{31})^*) = 0$

(j) By the same argument, the equations (modules) admit solutions G_{32}, G_{33} if 578

579

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - \quad 580$$

$$[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0 \quad 581$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^* such that $\varphi(G^*) = 0$

Finally we obtain the unique solution of 89 to 94 582

G_{14}^* given by $\varphi(G^*) = 0, T_{14}^*$ given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} , G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} , T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 583

G_{17}^* given by $\varphi((G_{19})^*) = 0, T_{17}^*$ given by $f(T_{17}^*) = 0$ and 584

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} , G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 585$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} , T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 586$$

Obviously, these values represent an equilibrium solution 587

Finally we obtain the unique solution 588

G_{21}^* given by $\varphi((G_{23})^*) = 0, T_{21}^*$ given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} , G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}^*)]} , T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 589

G_{25}^* given by $\varphi(G_{27}) = 0, T_{25}^*$ given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)}G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} , G_{26}^* = \frac{(a_{26})^{(4)}G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)}T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} , T_{26}^* = \frac{(b_{26})^{(4)}T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 590$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 591

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)}G_{29}^*}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^*)]} , G_{30}^* = \frac{(a_{30})^{(5)}G_{29}^*}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^*)]} \tag{592}$$

$$T_{28}^* = \frac{(b_{28})^{(5)}T_{29}^*}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^*)]} , T_{30}^* = \frac{(b_{30})^{(5)}T_{29}^*}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^*)]} \tag{592}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 593

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)}G_{33}^*}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^*)]} , G_{34}^* = \frac{(a_{34})^{(6)}G_{33}^*}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^*)]} \tag{594}$$

$$T_{32}^* = \frac{(b_{32})^{(6)}T_{33}^*}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^*)]} , T_{34}^* = \frac{(b_{34})^{(6)}T_{33}^*}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^*)]} \tag{594}$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS 595

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ and $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i , T_i = T_i^* + \mathbb{T}_i \tag{596}$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} , \frac{\partial(b_{14}'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain 597

$$\frac{d\mathbb{G}_{13}}{dt} = -((a'_{13})^{(1)} + (p_{13})^{(1)})\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \tag{598}$$

$$\frac{d\mathbb{G}_{14}}{dt} = -((a'_{14})^{(1)} + (p_{14})^{(1)})\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \tag{599}$$

$$\frac{d\mathbb{G}_{15}}{dt} = -((a'_{15})^{(1)} + (p_{15})^{(1)})\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \tag{600}$$

$$\frac{d\mathbb{T}_{13}}{dt} = -((b'_{13})^{(1)} - (r_{13})^{(1)})\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}(s_{(13)(j)})T_{13}^*\mathbb{G}_j \tag{601}$$

$$\frac{d\mathbb{T}_{14}}{dt} = -((b'_{14})^{(1)} - (r_{14})^{(1)})\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}(s_{(14)(j)})T_{14}^*\mathbb{G}_j \tag{602}$$

$$\frac{d\mathbb{T}_{15}}{dt} = -((b'_{15})^{(1)} - (r_{15})^{(1)})\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}(s_{(15)(j)})T_{15}^*\mathbb{G}_j \tag{603}$$

If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ 604
 Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Denote 605

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i \tag{606}$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \quad \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \tag{607}$$

taking into account equations (global) and neglecting the terms of power 2, we obtain 608

$$\frac{d\mathbb{G}_{16}}{dt} = -((a'_{16})^{(2)} + (p_{16})^{(2)})\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \tag{609}$$

$$\frac{d\mathbb{G}_{17}}{dt} = -((a'_{17})^{(2)} + (p_{17})^{(2)})\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \tag{610}$$

$$\frac{d\mathbb{G}_{18}}{dt} = -((a'_{18})^{(2)} + (p_{18})^{(2)})\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \tag{611}$$

$$\frac{d\mathbb{T}_{16}}{dt} = -((b'_{16})^{(2)} - (r_{16})^{(2)})\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}(s_{(16)(j)})\mathbb{T}_{16}^*\mathbb{G}_j \tag{612}$$

$$\frac{d\mathbb{T}_{17}}{dt} = -((b'_{17})^{(2)} - (r_{17})^{(2)})\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}(s_{(17)(j)})\mathbb{T}_{17}^*\mathbb{G}_j \tag{613}$$

$$\frac{d\mathbb{T}_{18}}{dt} = -((b'_{18})^{(2)} - (r_{18})^{(2)})\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}(s_{(18)(j)})\mathbb{T}_{18}^*\mathbb{G}_j \tag{614}$$

If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable 615

Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \quad , \quad \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

616

Then taking into account equations (global) and neglecting the terms of power 2, we obtain 617

$$\frac{d\mathbb{G}_{20}}{dt} = -((a'_{20})^{(3)} + (p_{20})^{(3)})\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \tag{618}$$

$$\frac{d\mathbb{G}_{21}}{dt} = -((a'_{21})^{(3)} + (p_{21})^{(3)})\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \tag{619}$$

$$\frac{d\mathbb{G}_{22}}{dt} = -((a'_{22})^{(3)} + (p_{22})^{(3)})\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \tag{6120}$$

$$\frac{d\mathbb{T}_{20}}{dt} = -((b'_{20})^{(3)} - (r_{20})^{(3)})\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}(s_{(20)(j)})\mathbb{T}_{20}^*\mathbb{G}_j \tag{621}$$

$$\frac{d\mathbb{T}_{21}}{dt} = -((b'_{21})^{(3)} - (r_{21})^{(3)})\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}(s_{(21)(j)})\mathbb{T}_{21}^*\mathbb{G}_j \tag{622}$$

$$\frac{d\mathbb{T}_{22}}{dt} = -((b'_{22})^{(3)} - (r_{22})^{(3)})\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}(s_{(22)(j)})\mathbb{T}_{22}^*\mathbb{G}_j \tag{623}$$

If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable 624

Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

625

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \quad \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain

626

$$\frac{d\mathbb{G}_{24}}{dt} = -((a'_{24})^{(4)} + (p_{24})^{(4)})\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^*\mathbb{T}_{25} \quad 627$$

$$\frac{d\mathbb{G}_{25}}{dt} = -((a'_{25})^{(4)} + (p_{25})^{(4)})\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 628$$

$$\frac{d\mathbb{G}_{26}}{dt} = -((a'_{26})^{(4)} + (p_{26})^{(4)})\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 629$$

$$\frac{d\mathbb{T}_{24}}{dt} = -((b'_{24})^{(4)} - (r_{24})^{(4)})\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(24)(j)})T_{24}^*\mathbb{G}_j \quad 630$$

$$\frac{d\mathbb{T}_{25}}{dt} = -((b'_{25})^{(4)} - (r_{25})^{(4)})\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26} (s_{(25)(j)})T_{25}^*\mathbb{G}_j \quad 631$$

$$\frac{d\mathbb{T}_{26}}{dt} = -((b'_{26})^{(4)} - (r_{26})^{(4)})\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26} (s_{(26)(j)})T_{26}^*\mathbb{G}_j \quad 632$$

633

If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

634

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \quad \frac{\partial(b_i'')^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations (global) and neglecting the terms of power 2, we obtain

635

$$\frac{d\mathbb{G}_{28}}{dt} = -((a'_{28})^{(5)} + (p_{28})^{(5)})\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 636$$

$$\frac{d\mathbb{G}_{29}}{dt} = -((a'_{29})^{(5)} + (p_{29})^{(5)})\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 637$$

$$\frac{d\mathbb{G}_{30}}{dt} = -((a'_{30})^{(5)} + (p_{30})^{(5)})\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 638$$

$$\frac{d\mathbb{T}_{28}}{dt} = -((b'_{28})^{(5)} - (r_{28})^{(5)})\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(28)(j)})T_{28}^*\mathbb{G}_j \quad 639$$

$$\frac{d\mathbb{T}_{29}}{dt} = -((b'_{29})^{(5)} - (r_{29})^{(5)})\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30} (s_{(29)(j)})T_{29}^*\mathbb{G}_j \quad 640$$

$$\frac{d\mathbb{T}_{30}}{dt} = -((b'_{30})^{(5)} - (r_{30})^{(5)})\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30} (s_{(30)(j)})T_{30}^*\mathbb{G}_j \quad 641$$

642

If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , \quad T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{33}''^{(6)})}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \quad , \quad \frac{\partial(b_i''^{(6)})}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations(global) and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -((a'_{32})^{(6)} + (p_{32})^{(6)})\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33}$$

$$\frac{d\mathbb{G}_{33}}{dt} = -((a'_{33})^{(6)} + (p_{33})^{(6)})\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33}$$

$$\frac{d\mathbb{G}_{34}}{dt} = -((a'_{34})^{(6)} + (p_{34})^{(6)})\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33}$$

$$\frac{d\mathbb{T}_{32}}{dt} = -((b'_{32})^{(6)} - (r_{32})^{(6)})\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(32)(j)})T_{32}^*\mathbb{G}_j$$

$$\frac{d\mathbb{T}_{33}}{dt} = -((b'_{33})^{(6)} - (r_{33})^{(6)})\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}(s_{(33)(j)})T_{33}^*\mathbb{G}_j$$

$$\frac{d\mathbb{T}_{34}}{dt} = -((b'_{34})^{(6)} - (r_{34})^{(6)})\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}(s_{(34)(j)})T_{34}^*\mathbb{G}_j$$

The characteristic equation of this system is

$$((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)})\{((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)})$$

$$\left[((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)})(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^* \right]$$

$$\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^* \right)$$

$$+ \left(((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)})(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^* \right)$$

$$\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^* \right)$$

$$\left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right)$$

$$\left(((\lambda)^{(1)})^2 + ((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}) (\lambda)^{(1)} \right)$$

$$+ \left(((\lambda)^{(1)})^2 + ((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}) (\lambda)^{(1)} \right) (q_{15})^{(1)}G_{15}$$

$$+ ((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}) \left((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^* \right)$$

$$\left(((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)})s_{(14),(15)}T_{14}^* + (b_{14})^{(1)}s_{(13),(15)}T_{13}^* \right) \} = 0$$

+

$$((\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)})\{((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)})$$

643

644

645

646

647

648

649

650

651

652

653

$$\begin{aligned}
 & \left[\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)} \right) (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (q_{16})^{(2)} G_{16}^* \right] \\
 & \left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)} \right) s_{(17),(17)} T_{17}^* + (b_{17})^{(2)} s_{(16),(17)} T_{17}^* \\
 & + \left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)} \right) (q_{16})^{(2)} G_{16}^* + (a_{16})^{(2)} (q_{17})^{(2)} G_{17}^* \\
 & \left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)} \right) s_{(17),(16)} T_{17}^* + (b_{17})^{(2)} s_{(16),(16)} T_{16}^* \\
 & \left((\lambda)^{(2)} \right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & \left((\lambda)^{(2)} \right)^2 + \left((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)} \right) (\lambda)^{(2)} \\
 & + \left((\lambda)^{(2)} \right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)} \right) (\lambda)^{(2)} (q_{18})^{(2)} G_{18} \\
 & + \left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)} \right) \left((a_{18})^{(2)} (q_{17})^{(2)} G_{17}^* + (a_{17})^{(2)} (a_{18})^{(2)} (q_{16})^{(2)} G_{16}^* \right) \\
 & \left. \left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)} \right) s_{(17),(18)} T_{17}^* + (b_{17})^{(2)} s_{(16),(18)} T_{16}^* \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & \left((\lambda)^{(3)} + (b'_{22})^{(3)} - (r_{22})^{(3)} \right) \left\{ \left((\lambda)^{(3)} + (a'_{22})^{(3)} + (p_{22})^{(3)} \right) \right. \\
 & \left. \left[\left((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)} \right) (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (q_{20})^{(3)} G_{20}^* \right] \right. \\
 & \left((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) s_{(21),(21)} T_{21}^* + (b_{21})^{(3)} s_{(20),(21)} T_{21}^* \\
 & + \left((\lambda)^{(3)} + (a'_{21})^{(3)} + (p_{21})^{(3)} \right) (q_{20})^{(3)} G_{20}^* + (a_{20})^{(3)} (q_{21})^{(1)} G_{21}^* \\
 & \left((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) s_{(21),(20)} T_{21}^* + (b_{21})^{(3)} s_{(20),(20)} T_{20}^* \\
 & \left((\lambda)^{(3)} \right)^2 + \left((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & \left((\lambda)^{(3)} \right)^2 + \left((b'_{20})^{(3)} + (b'_{21})^{(3)} - (r_{20})^{(3)} + (r_{21})^{(3)} \right) (\lambda)^{(3)} \\
 & + \left((\lambda)^{(3)} \right)^2 + \left((a'_{20})^{(3)} + (a'_{21})^{(3)} + (p_{20})^{(3)} + (p_{21})^{(3)} \right) (\lambda)^{(3)} (q_{22})^{(3)} G_{22} \\
 & + \left((\lambda)^{(3)} + (a'_{20})^{(3)} + (p_{20})^{(3)} \right) \left((a_{22})^{(3)} (q_{21})^{(3)} G_{21}^* + (a_{21})^{(3)} (a_{22})^{(3)} (q_{20})^{(3)} G_{20}^* \right) \\
 & \left. \left((\lambda)^{(3)} + (b'_{20})^{(3)} - (r_{20})^{(3)} \right) s_{(21),(22)} T_{21}^* + (b_{21})^{(3)} s_{(20),(22)} T_{20}^* \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & \left((\lambda)^{(4)} + (b'_{26})^{(4)} - (r_{26})^{(4)} \right) \left\{ \left((\lambda)^{(4)} + (a'_{26})^{(4)} + (p_{26})^{(4)} \right) \right. \\
 & \left. \left[\left((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)} \right) (q_{25})^{(4)} G_{25}^* + (a_{25})^{(4)} (q_{24})^{(4)} G_{24}^* \right] \right.
 \end{aligned}$$

$$\begin{aligned}
 & \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)})s_{(25),(25)}T_{25}^* + (b_{25})^{(4)}s_{(24),(25)}T_{25}^* \right) \\
 & + \left(((\lambda)^{(4)} + (a'_{25})^{(4)} + (p_{25})^{(4)})(q_{24})^{(4)}G_{24}^* + (a_{24})^{(4)}(q_{25})^{(4)}G_{25}^* \right) \\
 & \quad \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)})s_{(25),(24)}T_{25}^* + (b_{25})^{(4)}s_{(24),(24)}T_{24}^* \right) \\
 & \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & \quad \left(((\lambda)^{(4)})^2 + ((b'_{24})^{(4)} + (b'_{25})^{(4)} - (r_{24})^{(4)} + (r_{25})^{(4)}) (\lambda)^{(4)} \right) \\
 & + \left(((\lambda)^{(4)})^2 + ((a'_{24})^{(4)} + (a'_{25})^{(4)} + (p_{24})^{(4)} + (p_{25})^{(4)}) (\lambda)^{(4)} \right) (q_{26})^{(4)}G_{26} \\
 & + ((\lambda)^{(4)} + (a'_{24})^{(4)} + (p_{24})^{(4)}) ((a_{26})^{(4)}(q_{25})^{(4)}G_{25}^* + (a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G_{24}^*) \\
 & \left. \left(((\lambda)^{(4)} + (b'_{24})^{(4)} - (r_{24})^{(4)})s_{(25),(26)}T_{25}^* + (b_{25})^{(4)}s_{(24),(26)}T_{24}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(5)} + (b'_{30})^{(5)} - (r_{30})^{(5)}) \{ ((\lambda)^{(5)} + (a'_{30})^{(5)} + (p_{30})^{(5)}) \\
 & \left[\left(((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)})(q_{29})^{(5)}G_{29}^* + (a_{29})^{(5)}(q_{28})^{(5)}G_{28}^* \right) \right] \\
 & \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})s_{(29),(29)}T_{29}^* + (b_{29})^{(5)}s_{(28),(29)}T_{29}^* \right) \\
 & + \left(((\lambda)^{(5)} + (a'_{29})^{(5)} + (p_{29})^{(5)})(q_{28})^{(5)}G_{28}^* + (a_{28})^{(5)}(q_{29})^{(5)}G_{29}^* \right) \\
 & \quad \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})s_{(29),(28)}T_{29}^* + (b_{29})^{(5)}s_{(28),(28)}T_{28}^* \right) \\
 & \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & \quad \left(((\lambda)^{(5)})^2 + ((b'_{28})^{(5)} + (b'_{29})^{(5)} - (r_{28})^{(5)} + (r_{29})^{(5)}) (\lambda)^{(5)} \right) \\
 & + \left(((\lambda)^{(5)})^2 + ((a'_{28})^{(5)} + (a'_{29})^{(5)} + (p_{28})^{(5)} + (p_{29})^{(5)}) (\lambda)^{(5)} \right) (q_{30})^{(5)}G_{30} \\
 & + ((\lambda)^{(5)} + (a'_{28})^{(5)} + (p_{28})^{(5)}) ((a_{30})^{(5)}(q_{29})^{(5)}G_{29}^* + (a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G_{28}^*) \\
 & \left. \left(((\lambda)^{(5)} + (b'_{28})^{(5)} - (r_{28})^{(5)})s_{(29),(30)}T_{29}^* + (b_{29})^{(5)}s_{(28),(30)}T_{28}^* \right) \right\} = 0
 \end{aligned}$$

+

$$\begin{aligned}
 & ((\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)}) \{ ((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)}) \\
 & \left[\left(((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)})(q_{33})^{(6)}G_{33}^* + (a_{33})^{(6)}(q_{32})^{(6)}G_{32}^* \right) \right] \\
 & \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(33)}T_{33}^* + (b_{33})^{(6)}s_{(32),(33)}T_{33}^* \right)
 \end{aligned}$$

$$\begin{aligned}
 &+ \left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)}) (q_{32})^{(6)} G_{32}^* + (a_{32})^{(6)} (q_{33})^{(6)} G_{33}^* \right) \\
 &\quad \left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(32)} T_{33}^* + (b_{33})^{(6)} s_{(32),(32)} T_{32}^* \right) \\
 &\left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 &\quad \left(((\lambda)^{(6)})^2 + ((b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)}) (\lambda)^{(6)} \right) \\
 &+ \left(((\lambda)^{(6)})^2 + ((a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)}) (\lambda)^{(6)} \right) (q_{34})^{(6)} G_{34} \\
 &+ ((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}) ((a_{34})^{(6)} (q_{33})^{(6)} G_{33}^* + (a_{33})^{(6)} (a_{34})^{(6)} (q_{32})^{(6)} G_{32}^*) \\
 &\left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)}) s_{(33),(34)} T_{33}^* + (b_{33})^{(6)} s_{(32),(34)} T_{32}^* \right) \} = 0
 \end{aligned}$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

Acknowledgments:

The introduction is a collection of information from various articles, Books, News Paper reports, Home Pages Of authors, Journal Reviews, Nature 's Letters, Article Abstracts, Research papers, Abstracts Of Research Papers, Stanford Encyclopedia, Web Pages, Ask a Physicist Column, Deliberations with Professors, the internet including Wikipedia. We acknowledge all authors who have contributed to the same. In the eventuality of the fact that there has been any act of omission on the part of the authors, we regret with great deal of compunction, contrition, regret, trepidation and remorse. As Newton said, it is only because erudite and eminent people allowed one to piggy ride on their backs; probably an attempt has been made to look slightly further. Once again, it is stated that the references are only illustrative and not comprehensive

REFERENCES

1. Dr K N Prasanna Kumar, Prof B S Kiranagi, Prof C S Bagewadi - [MEASUREMENT DISTURBS EXPLANATION OF QUANTUM MECHANICAL STATES-A HIDDEN VARIABLE THEORY](#) - published at: "*International Journal of Scientific and Research Publications, Volume 2, Issue 5, May 2012 Edition*".
2. DR K N PRASANNA KUMAR, PROF B S KIRANAGI and PROF C S BAGEWADI - [CLASSIC 2 FLAVOUR COLOR SUPERCONDUCTIVITY AND ORDINARY NUCLEAR MATTER-A NEW PARADIGM STATEMENT](#) - published at: "*International Journal of Scientific and Research Publications, Volume 2, Issue 5, May 2012 Edition*".
3. A HAIMOVICI: "On the growth of a two species ecological system divided on age groups". Tensor, Vol 37 (1982), Commemoration volume dedicated to Professor Akitsugu Kawaguchi on his 80th birthday
4. FRTJOF CAPRA: "The web of life" Flamingo, Harper Collins See "Dissipative structures" pages 172-188
5. HEYLIGHEN F. (2001): "The Science of Self-organization and Adaptivity", in L. D. Kiel, (ed) . Knowledge Management, Organizational Intelligence and Learning, and Complexity, in: The

Encyclopedia of Life Support Systems ((EOLSS), (Eolss Publishers, Oxford)
[<http://www.eolss.net>

6. MATSUI, T, H. Masunaga, S. M. Kreidenweis, R. A. Pielke Sr., W.-K. Tao, M. Chin, and Y. J. Kaufman (2006), "Satellite-based assessment of marine low cloud variability associated with aerosol, atmospheric stability, and the diurnal cycle", *J. Geophys. Res.*, 111, D17204, doi:10.1029/2005JD006097
7. STEVENS, B, G. Feingold, W.R. Cotton and R.L. Walko, "Elements of the microphysical structure of numerically simulated nonprecipitating stratocumulus" *J. Atmos. Sci.*, 53, 980-1006
8. FEINGOLD, G, Koren, I; Wang, HL; Xue, HW; Brewer, WA (2010), "Precipitation-generated oscillations in open cellular cloud fields" *Nature*, 466 (7308) 849-852, doi: 10.1038/nature09314, Published 12-Aug 2010

- (9)^{a b c} Einstein, A. (1905), "Ist die Trägheit eines Körpers von seinem Energieinhalt abhängig?", *Annalen der Physik* **18**: 639 Bibcode 1905AnP...323..639E, DOI:10.1002/andp.19053231314. See also the English translation.
- (10)^{a b} Paul Allen Tipler, Ralph A. Llewellyn (2003-01), *Modern Physics*, W. H. Freeman and Company, pp. 87–88, ISBN 0-7167-4345-0
- (11)^{a b} Rainville, S. et al. World Year of Physics: A direct test of $E=mc^2$. *Nature* 438, 1096-1097 (22 December 2005) | doi: 10.1038/4381096a; Published online 21 December 2005.
- (12)[^] In F. Fernflores. The Equivalence of Mass and Energy. Stanford Encyclopedia of Philosophy
- (13)[^] Note that the relativistic mass, in contrast to the rest mass m_0 , is not a relativistic invariant, and that the velocity is not a Minkowski four-vector, in contrast to the quantity dx^μ , where dx^μ is the differential of the proper time. However, the energy-momentum four-vector is a genuine Minkowski four-vector, and the intrinsic origin of the square-root in the definition of the relativistic mass is the distinction between $d\mathbf{x}$ and dt .
- (14)[^] Relativity DeMystified, D. McMahon, Mc Graw Hill (USA), 2006, ISBN 0-07-145545-0
- (15)[^] Dynamics and Relativity, J.R. Forshaw, A.G. Smith, Wiley, 2009, ISBN 978-0-470-01460-8
- (16)[^] Hans, H. S.; Puri, S. P. (2003). *Mechanics* (2 ed.). Tata McGraw-Hill. p. 433. ISBN 0-07-047360-9., Chapter 12 page 433
- (17)[^] E. F. Taylor and J. A. Wheeler, **Spacetime Physics**, W.H. Freeman and Co., NY. 1992. ISBN 0-7167-2327-1, see pp. 248-9 for discussion of mass remaining constant after detonation of nuclear bombs, until heat is allowed to escape.
- (18)[^] Mould, Richard A. (2002). *Basic relativity* (2 ed.). Springer. p. 126. ISBN 0-387-95210-1., Chapter 5 page 126
- (19)[^] Chow, Tail L. (2006). *Introduction to electromagnetic theory: a modern perspective*. Jones & Bartlett Learning. p. 392. ISBN 0-7637-3827-1., Chapter 10 page 392
- (20)[^] [2] Cockcroft-Walton experiment
- (21)^{a b c} Conversions used: 1956 International (Steam) Table (IT) values where one calorie $\equiv 4.1868$ J and one BTU $\equiv 1055.05585262$ J. Weapons designers' conversion value of one gram TNT

≡ 1000 calories used.

(22)[^] Assuming the dam is generating at its peak capacity of 6,809 MW.

(23)[^] Assuming a 90/10 alloy of Pt/Ir by weight, a C_p of 25.9 for Pt and 25.1 for Ir, a Pt-dominated average C_p of 25.8, 5.134 moles of metal, and 132 J.K^{-1} for the prototype. A variation of ± 1.5 picograms is of course, much smaller than the actual uncertainty in the mass of the international prototype, which are ± 2 micrograms.

(24)[^] [3] Article on Earth rotation energy. Divided by c^2 .

(25)^{^ a b} Earth's gravitational self-energy is 4.6×10^{-10} that of Earth's total mass, or 2.7 trillion metric tons. Citation: *The Apache Point Observatory Lunar Laser-Ranging Operation (APOLLO)*, T. W. Murphy, Jr. *et al.* University of Washington, Dept. of Physics (132 kB PDF, [here](#).)

(26)[^] There is usually more than one possible way to define a field energy, because any field can be made to couple to gravity in many different ways. By general scaling arguments, the correct answer at everyday distances, which are long compared to the quantum gravity scale, should be *minimal coupling*, which means that no powers of the curvature tensor appear. Any non-minimal couplings, along with other higher order terms, are presumably only determined by a theory of quantum gravity, and within string theory, they only start to contribute to experiments at the string scale.

(27)[^] G. 't Hooft, "Computation of the quantum effects due to a four-dimensional pseudoparticle", *Physical Review D* 14:3432–3450 (1976).

(28)[^] A. Belavin, A. M. Polyakov, A. Schwarz, Yu. Tyupkin, "Pseudoparticle Solutions to Yang Mills Equations", *Physics Letters* 59B:85 (1975).

(29)[^] F. Klinkhammer, N. Manton, "A Saddle Point Solution in the Weinberg Salam Theory", *Physical Review D* 30:2212.

(30)[^] Rubakov V. A. "Monopole Catalysis of Proton Decay", *Reports on Progress in Physics* 51:189–241 (1988).

(31)[^] S.W. Hawking "Black Holes Explosions?" *Nature* 248:30 (1974).

(32)[^] Einstein, A. (1905), "Zur Elektrodynamik bewegter Körper." (PDF), *Annalen der Physik* 17: 891–921, Bibcode 1905AnP...322...891E, DOI:10.1002/andp.19053221004. English translation.

- (33)[^] See e.g. Lev B. Okun, *The concept of Mass*, Physics Today **42** (6), June 1969, p. 31–36, http://www.physicstoday.org/vol-42/iss-6/vol42no6p31_36.pdf
- (34)[^] Max Jammer (1999), *Concepts of mass in contemporary physics and philosophy*, Princeton University Press, p. 51, ISBN 0-691-01017-X
- (35)[^] Eriksen, Erik; Vøyenli, Kjell (1976), "The classical and relativistic concepts of mass", *Foundations of Physics* (Springer) **6**: 115–124, Bibcode 1976FoPh...6..115E, DOI:10.1007/BF00708670
- (36)[^] ^{a b} Janssen, M., Mecklenburg, M. (2007), *From classical to relativistic mechanics: Electromagnetic models of the electron.*, in V. F. Hendricks, et al., , *Interactions: Mathematics, Physics and Philosophy* (Dordrecht: Springer): 65–134
- (37)[^] ^{a b} Whittaker, E.T. (1951–1953), *2. Edition: A History of the theories of aether and electricity, vol. 1: The classical theories / vol. 2: The modern theories 1900–1926*, London: Nelson
- (38)[^] Miller, Arthur I. (1981), *Albert Einstein's special theory of relativity. Emergence (1905) and early interpretation (1905–1911)*, Reading: Addison–Wesley, ISBN 0-201-04679-2
- (39)[^] ^{a b} Darrigol, O. (2005), "The Genesis of the theory of relativity." (PDF), *Séminaire Poincaré* **1**: 1–22
- (40)[^] Philip Ball (Aug 23, 2011). "Did Einstein discover $E = mc^2$?" Physics World.
- (41)[^] Ives, Herbert E. (1952), "Derivation of the mass-energy relation", *Journal of the Optical Society of America* **42** (8): 540–543, DOI:10.1364/JOSA.42.000540
- (42)[^] Jammer, Max (1961/1997). *Concepts of Mass in Classical and Modern Physics*. New York: Dover. ISBN 0-486-29998-8.
- (43)[^] Stachel, John; Torretti, Roberto (1982), "Einstein's first derivation of mass-energy equivalence", *American Journal of Physics* **50** (8): 760–763, Bibcode1982AmJPh..50..760S, DOI:10.1119/1.12764
- (44)[^] Ohanian, Hans (2008), "Did Einstein prove $E=mc^2$ ", *Studies In History and Philosophy of Science Part B* **40** (2): 167–173, arXiv:0805.1400, DOI:10.1016/j.shpsb.2009.03.002
- (45)[^] Hecht, Eugene (2011), "How Einstein confirmed $E_0=mc^2$ ", *American Journal of Physics* **79** (6): 591–600, Bibcode 2011AmJPh..79..591H, DOI:10.1119/1.3549223

(46)^ Rohrlich, Fritz (1990), "An elementary derivation of $E=mc^2$ ", *American Journal of Physics* **58** (4): 348–349, Bibcode 1990AmJPh..58..348R, DOI:10.1119/1.16168

(47) (1996). *Lise Meitner: A Life in Physics*. California Studies in the History of Science. **13**. Berkeley: University of California Press. pp. 236–237. ISBN 0-520-20860-

(48)^ UIBK.ac.at

(49)^ J. J. L. Morton; *et al.* (2008). "Solid-state quantum memory using the ^{31}P nuclear spin". *Nature* **455** (7216): 1085–1088. Bibcode 2008Natur.455.1085M.DOI:10.1038/nature07295.

(50)^ S. Weisner (1983). "Conjugate coding". *Association of Computing Machinery, Special Interest Group in Algorithms and Computation Theory* **15**: 78–88.

(51)^ A. Zelinger, *Dance of the Photons: From Einstein to Quantum Teleportation*, Farrar, Straus & Giroux, New York, 2010, pp. 189, 192, ISBN 0374239665

(52)^ B. Schumacher (1995). "Quantum coding". *Physical Review A* **51** (4): 2738–2747. Bibcode 1995PhRvA..51.2738S. DOI:10.1103/PhysRevA.51.2738.

(53)^ ^{a b} Straumann, N (2000). "On Pauli's invention of non-abelian Kaluza-Klein Theory in 1953". ArXiv: gr-qc/0012054 [gr-qc].

(54)^ See Abraham Pais' account of this period as well as L. Susskind's "Superstrings, Physics World on the first non-abelian gauge theory" where Susskind wrote that Yang–Mills was "rediscovered" only because Pauli had chosen not to publish.

(55)^ Reifler, N (2007). "Conditions for exact equivalence of Kaluza-Klein and Yang-Mills theories". ArXiv: gr-qc/0707.3790 [gr-qc].

(56)^ Yang, C. N.; Mills, R. (1954). "Conservation of Isotopic Spin and Isotopic Gauge Invariance". *Physical Review* **96** (1): 191–195. Bibcode 1954PhRv...96...191Y.DOI:10.1103/PhysRev.96.191.

(57)^ Caprini, I.; Colangelo, G.; Leutwyler, H. (2006). "Mass and width of the lowest resonance in QCD". *Physical Review Letters* **96** (13): 132001. ArXiv: hep-ph/0512364. Bibcode 2006PhRvL..96m2001C. DOI:10.1103/PhysRevLett.96.132001.

(58)[^] Yndurain, F. J.; Garcia-Martin, R.; Pelaez, J. R. (2007). "Experimental status of the $\pi\pi$ isoscalar S wave at low energy: $f_0(600)$ pole and scattering length". *Physical Review D* **76** (7): 074034. ArXiv:hep-ph/0701025. Bibcode 2007PhRvD..76g4034G.DOI:10.1103/PhysRevD.76.074034.

(59)[^] Novikov, V. A.; Shifman, M. A.; A. I. Vainshtein, A. I.; Zakharov, V. I. (1983). "Exact Gell-Mann-Low Function of Supersymmetric Yang-Mills Theories From Instanton Calculus". *Nuclear* **229** (2): 381–393. Bibcode 1983NuPhB.229..381N.DOI:10.1016/0550-3213(83)90338-3.

(60)[^] Rytov, T.; Sannino, F. (2008). "Super symmetry Inspired QCD Beta Function". *Physical Review D* **78** (6): 065001. Bibcode 2008PhRvD..78f5001R.DOI:10.1103/PhysRevD.78.065001

(61)[^] Bogolubsky, I. L.; Ilgenfritz, E.-M.; A. I. Müller-Preussker, M.; Sternbeck, A. (2009). "Lattice gluodynamics computation of Landau-gauge Green's functions in the deep infrared". *Physics Letters B* **676** (1-3): 69–73. Bibcode 2009PhLB..676...69B.DOI:10.1016/j.physletb.2009.04.076.

First Author: ¹**Mr. K. N.Prasanna Kumar** has three doctorates one each in Mathematics, Economics, Political Science. Thesis was based on Mathematical Modeling. He was recently awarded D.litt. for his work on 'Mathematical Models in Political Science'--- Department of studies in Mathematics, Kuvempu University, Shimoga, Karnataka, India Corresponding Author:drknpkumar@gmail.com

Second Author: ²**Prof. B.S Kiranagi** is the Former Chairman of the Department of Studies in Mathematics, Manasa Gangotri and present Professor Emeritus of UGC in the Department. Professor Kiranagi has guided over 25 students and he has received many encomiums and laurels for his contribution to Co homology Groups and Mathematical Sciences. Known for his prolific writing, and one of the senior most Professors of the country, he has over 150 publications to his credit. A prolific writer and a prodigious thinker, he has to his credit several books on Lie Groups, Co Homology Groups, and other mathematical application topics, and excellent

publication history.-- UGC Emeritus Professor (Department of studies in Mathematics), Manasagangotri, University of Mysore, Karnataka, India

Third Author: ³**Prof. C.S. Bagewadi** is the present Chairman of Department of Mathematics and Department of Studies in Computer Science and has guided over 25 students. He has published articles in both national and international journals. Professor Bagewadi specializes in Differential Geometry and its wide-ranging ramifications. He has to his credit more than 159 research papers. Several Books on Differential Geometry, Differential Equations are coauthored by him--- Chairman, Department of studies in Mathematics and Computer science, Jnanasahyadri Kuvempu University, Shankarghatta, Shimoga district, Karnataka, India
